

Chemical Composition and Biological Activities of Essential Oils from the Oleogum Resins of Three Endemic Soqotraen *Boswellia* Species

Nasser A. Awadh Ali^{*1}, Martina Wurster³, Norbert Arnold², Axel Teichert²,
Jürgen Schmidt², Ulrike Lindequist³ and Ludger Wessjohann²

¹Department of Pharmacognosy, Faculty of Pharmacy, Sana'a University, P.O. Box 13150, Yemen

²Leibniz Institute of Plant Biochemistry, Department of Bio-organic Chemistry, Weinberg 3, 06120 Halle/Saale, Germany

³Department of Pharmaceutical Biology, Institute of Pharmacy, Ernst-Moritz-Arndt-University Greifswald, Friedrich-Ludwig-Jahn-Str. 17, 17487 Greifswald Germany

(Received December, 16, 2007; Revised January, 8, 2008, Accepted January, 12, 2008)

Abstract: The chemical composition, antioxidant and anticholinesterase activity of three essential oils (EOs) obtained from the oleogum resin of three endemic Soqotraen *Boswellia* species, *Boswellia socotrana* Balf. f, *Boswellia ameero* Balf. f, and *Boswellia elongata* Balf. f were determined. GC-MS technique was used for the analysis of the oils. Oils of *B. socotrana* and *B. ameero* were characterized by a high content of monoterpenes. The main constituents of *B. socotrana* and *B. ameero* were (E)-2,3-epoxycarene (51.8%), 1,5-isopropyl-2-methylbicyclo[3.1.0]hex-3-en-2-ol (31.3%), and α -cymene (7.1%); (3E,5E)-2,6-dimethyl-1,3,5,7-octatetraene (34.9%), 1-(2,4-Dimethylphenyl)ethanol (20.3%), 3,4-dimethylstyrene (17.3%), α -campholenal (13.4%) and α -terpineol (12.4%) respectively. The composition of *B. elongata* oil was dominated by the diterpene verticilol (52.4%), the sesquiterpene caryophellene (39.1%) and methylcycloundecanecarboxylate (7.8%). The oils were screened for their antioxidant activity by using the DPPH free radical scavenger assay and their anticholinesterase activity on acetylcholinesterase enzyme by using in vitro Ellman method. The antioxidant activity of EOs from *B. socotrana* (IC₅₀ =121.4 μ g/mL) appeared to be more potent than that of *B. elongata* (IC₅₀ =211.2 μ g/mL) and *B. ameero* (IC₅₀ =175.2 μ g/mL). EO of *B. socotrana* showed the higher AChE inhibitory activity with 59.3% at concentration of 200 μ g/mL in comparison to EOs of *B. elongata* and *B. ameero* (29.6, 41.6 enzyme inhibition) respectively.

Key words: Essential oils, *Boswellia*, GC-MS, antioxidant, AChE inhibitor, Soqotra

* Corresponding author: E-Mail: alinasser57@yahoo.com

1. Introduction

The genus *Boswellia* (family Burseraceae) consists of many species widespread throughout the world. It includes approximately 23 species of small trees that grow mainly in Arabia, on eastern coast of Africa and in India. Olibanum is a natural oleo-gum-resin that exudes from tappings in the bark of *Boswellia* trees [1]. The genus is represented in Soqotra Island by 8 species. The oleogum resin is used traditionally by the native inhabitants for relieving the pain in cavities, sweetening the breath, or to sooth a disturbed stomach [2].

Considerable work on the composition of the essential oils (EOs) from different species of *Boswellia* is reported in literature. Monoterpenoids seem to be the dominant class of compounds found in the oils. The main compounds of *B. dalzielii* EO from Nigeria were α -pinene (45.7%), γ -terpinene (11.5%) and trans-sabinene hydrate [3]. The EO of *B. frereana* from Somalia was dominated by α -thujene (10.10%), p-cymene (4.3%) [4]. The EO of *B. serrata* from India was found to contain mainly α -thujene (61.0%), α -pinene (7.7%), and sabinene (5.1%) [5]. Since the chemical composition of EOs depends on various environmental factors, three studies on the EOs composition of *B. carteri* from Somalia revealed this composition variety: α -thujene (19.2%), sabinene (9.4%), limonene (7.8%) and α -pinene (7.2%) [4]; octyl acetate (60.0%), octanol (12.7%) and p-cymene (8.7%) [6]; α -pinene (41.0%), limonene (12.8%) [7]

Several studies investigated the anti-inflammatory, immunomodulatory, anti-leukotriene, antiacetylcholinesterase, and anticancer activity of the resin and especially its major components, boswellic acid derivatives [8-18]. In addition, the EO showed antibacterial, antifungal and immunostimulating activity [7, 19, 20].

To the best of our knowledge, no literature information is available on the chemistry and biology of the EOs of Soqotraen *Boswellia* species. In the present paper, we wish to report the chemical composition, antioxidant, and antiacetylcholinesterase activities of the EOs obtained from the olegum resins of three endemic Soqotraen *Boswellia* species.

2. Materials and Methods

2.1 Plant material

The plant material was collected in March 2006 from different locations in Soqotra Island. The plants were taxonomically identified at the Centre of Environment and Biodiversity, Aden University, Yemen. Species names are according to International Plant Name Index (IPNI) (<http://www.ipni.org>). Voucher specimens (*B. socotrana* (SP-Bu-03) *B. elongata* (SP-Bu-02) and *B. ameero* (SP-Bu-01) of the plant material are deposited at the Pharmacognosy Department, Aden University, Yemen.

2.2 Volatile oil extraction

Oleogum resins (12 g each) of *B. socotrana*, *B. elongata*, *B. ameero*, were hydrodistilled for 2-3 h in a Clevenger type apparatus according to European Pharmacopia to yield three oils. The obtained volatile oils dried over anhydrous Na₂SO₄ and stored at 4 °C until testing.

2.3 Determination of antioxidant activity

Estimation of a radical scavenging effect was carried out by using a DPPH free radical scavenger assay in 96 well micro titre plates (MTP) according to the modified method given in refs. [21, 22]. A solution of DPPH (Sigma Aldrich, Germany) was prepared by dissolving 5 mg DPPH in 2 mL of methanol, and the solution was kept in the dark at 4 °C until use. Stock solutions of the samples

were prepared at 4 mg/mL and diluted. 5 μ L of methanolic DPPH solution was added to each well. The plate was shaken to ensure thorough mixing before being wrapped with aluminium foil and stored in the dark. After 30 min the optical density (OD) of the solution was measured at the wavelength of 517 nm using a micro titre plate reader Tecan GeniosPro (Germany). A methanolic solution of DPPH served as control. All tests were carried out in duplicates. Ascorbic acid (Sigma Aldrich, Germany) was used as positive control.

2.4. Microplate assay for AChE activity

2.4.1. Chemicals

Acetylthiocholine iodide (ATCI), 5,5'-dithiobis-(2-nitrobenzoic acid) (DTNB), galanthamine, bovine serum albumine (BSA) and acetylcholinesterase (AChE) from horse serum (lyophilized, 500 U/vial solid, 65 U/mg) were purchased from Sigma (Germany). The following buffers were used: Buffer A: 50 mM Tris-HCl, pH 8, containing 0.1% BSA; Buffer B: 50 mM Tris-HCl, pH 8 containing 0.1 M NaCl, 0.02 M MgCl₂·6H₂O; Galanthamine (Sigma Aldrich, Germany).

2.4.2. AChE inhibitory activity

AChE inhibitory activity was detected by a microtiter plate assay based on Ellman's method [23, 24] by using acetylthiocholine as substrate. In 96-well plates, 25 μ L of 15 mM ATCI (43 mg/10 mL Millipore water), 125 μ L of 3 mM DTNB (11.9 mg/10 mL buffer B), 50 μ L of buffer A, 25 μ L of EO at concentration of 2 and 0.5 mg/mL (final concentration in the assay: 0.2, 0.05 mg/mL DMSO) were added and the absorbance was measured at 405 nm every 13 s for five times. After adding 25 μ L of 0.22 U/mL enzyme (0.34 mg AChE dissolved in 100 mL buffer A), the absorbance was read again every 13 s for five times. The absorbance was measured using a Tecan GeniosPro micro plate reader. Percentage of inhibition was calculated by comparing the rates for the sample to the blank (DMSO), control contained all components except the tested EOs. Galanthamine was used as positive control. All treatments were performed in tetraplicate on each well, and was done twice. (n= 2)

2.5. Gas chromatography-mass spectrometry

Analytical GC-MS system consisted of an Agilent 6890N gas chromatograph and a mass selective detector (Agilent@5973 Network MSD). Injection was done with Agilent@7683 Series Injector (Split 1:40 at 250 C, 2.0 μ L; carrier gas: helium 1.1 mL/min (60 kPa) at 110°C; pressure rise: 6 kPa/min). The MS operated in the electron impact mode with an ionization energy of 70eV. The oven program started with 1min at 70°C, the oven temperature was increased at 3°C/min to 220°C. Full scan mass spectra were acquired from 45-650 m/z at a rate of 4.5 scans/s and with a 5.00 min solvent delay. Chromatography was performed using a 30 m DB-5 column (J&W Scientific, Folsom, USA) with 0.25 mm i.d. and 0.25 μ m film thickness.

The detected compounds were identified by processing of the raw GC-MS data with ChemStation G1701CA software and comparing with NIST mass spectral database 2.0 d (National Institute of Standards and Technology, Gaithersburg, USA) and from retention times and mass spectra of standard compounds. Relative amounts of detected compounds were calculated based on GC peak areas.

Table 1. Main Components of EOs from Oleogum resins of *Boswellia socotrana* (A), *Boswellia elongata* (B), *Boswellia ameero* (C)

RI	Compounds ^a	A (%)	B (%)	C (%)
1020	α -Cymene	7.1	-	-
1039	1,5-Isopropyl-2-methylbicyclo[3.1.0]hex-3-en-2-ol	31.3	-	-
1082	4-Terpinenylacetate	3.9	-	-
1111	α -Campholenal	-	-	13.4
1130	(3E,5E)-2,6-Dimethyl-1,3,5,7-octatetraene	-	-	34.9
1153	(E)-2,3-Epoxycarene	51.8	-	-
1170	3,4-Dimethylstyrene	-	-	17.3
1177	α -Terpineol	-	-	12.4
1196	1-(2,4-Dimethylphenyl)ethanol	-	-	20.3
1213	4-methyl-Benzoic acid	2.7	-	-
1238	p-Menth-1(7)-en-2-one	2.6	-	-
1350	α -Terpinylacetate	0.1	-	-
1432	Caryophyllene	-	39.1	-
1536	Methylcycloundecanecarboxylate	-	7.9	-
2106	Verticicol	-	52.4	-
	Total identified	98.5	99.4	98.3

^aCompounds listed in order to their elution on the DB-5 column
Retention indices on the DB-5 column relative to C₁₀-C₂₀ *n*-alkanes

3. Results and Discussion

The yields of EOs were obtained by hydrodistillation of the crushed oleogum resins derived from *B. socotrana*, *B. ameero* and *B. elongata* (1.2%, 1.8%, and 2.3%) on dry weight basis respectively. The reported average yields of some *Boswellia* species were ranging from 1.3% to 3% [3, 20].

There have been no reports on GC-MS analysis of the three EOs. The GC-MS analysis of EOs from *B. socotrana*, *B. ameero* and *B. elongata* allowed the identification of 7, 5, and 3 components which accounted for 98.5%, 98.3% and 99.4% of the three EOs respectively (Table 1).

Oil of *B. socotrana* was characterized by a higher content of oxygenated monoterpenes (89.9%): (E)-2,3-Epoxycarene (51.8%), 1,5-isopropyl-2-methyl bicyclo[3.1.0]hex-3-en-2-ol (31.3%), 4-Terpinenylacetate (3.9%) and p-Menth-1(7)-en-2-on (2.6%), while the oil of *B. ameero* contained (46.2%) phenylpropan derivatives and oxygenated monoterpenes: 1-(2,4-Dimethylphenyl)ethanol (20.3%), α -campholenal (13.4%) and α -terpineol (12.4%). Both oils were characterized by bicyclo[3.1.0]hexane and bicyclo[3.1.1]heptane.

EO of *B. elongata* possessed a different terpenic composition from the two aforementioned oils. The oil was poor in monoterpenes, but the diterpene verticicol (52.4%) predominated over the sesquiterpene caryophellene (39.1%). The presence of such a high content of a cembranoid diterpene, verticicol, was detected for the first time in the EO of frankincense derived from *B. elongata*. The comparison of our results with those of the literature showed that the main components of chemical composition of the three endemic Soqotraen *Boswellia* EOs were markedly different from that of other known *Boswellia* species such as *B. carteri*, *B. sacra*, *B. frereana*, *B. serrata*, and *B. dalzielii*. [3-7, 25]

Table 2. Antioxidant, anticholinesterase activity of EOs of *B. socotrana*, *B. elongata* and *B. ameero*

Sample	DPPH	Enzyme inhibition in %	
	IC ₅₀ (µg/mL)	200 µg/mL	50 µg/mL
<i>B. socotrana</i>	121.4 (± 7.6)	59.3 (±7.8)	19.1 (±6.6)
<i>B. elongata</i>	211.2 (± 12.8)	29.6 (± 6.7)	11.5 (±8.1)
<i>B. ameero</i>	175.2 (± 10.3)	41.6 (±6.0)	10.7 (±6.7)
Ascorbic acid	96.5 (± 5.9)		
Galanthamine (8 µg/mL)		87.8 (±7.7)	

The free- radical scavenging activity of *Boswellia* EOs evaluated using the DPPH method is presented in Table 2. The model of scavenging stable DPPH free radicals can be used to evaluate the antioxidative activities in a relatively short time. The absorbance decreases as a result of a color change from purple to yellow as the radical is scavenged by antioxidants through donation of hydrogen to form the stable DPPH-H molecule [26]. The effect of antioxidants on DPPH radical scavenging was thought to be due to their hydrogen-donating ability. The EOs were able to reduce the stable free radical DPPH to the yellow colored 1, 1 –diphenyl-2-picrylhydrazyl with an IC₅₀ value of 121.4 µg/mL, 211.2 µg/mL and 175.2 µg/mL for *B. socotrana*, *B. elongata* and *B. ameero* respectively. The free- radical scavenging activity of *B. socotrana* EO showed higher activity than those of *elongata* and *B. ameero*. The positive factor for *B. socotrana* EO was the higher concentration of oxygenated monoterpenes [27, 28] but there are no data on the antioxidant activity of the oxygenated monoterpenes (E)-2,3-epoxycarene that was the main constituent in this oil. *Boswellia* EOs showed lower free- radical scavenging activity in comparison to other reported essential oils rich in oxygenated monoterpenes, such as *Melissa officinalis*, and *M. piperita* with IC₅₀= 7.58 and 2.53 µg/mL, respectively [29, 30].

Inhibition of the acetylcholinesterase, which is responsible for the hydrolysis of acetylcholine, represents the most effective approach for finding new AChE inhibitors from natural sources for treating Alzheimer's disease. Currently no AChE inhibitory activity has been reported from EOs of *B. species*. *B. species* contain in addition to essential oils, triterpenic boswellic acids, a group of compounds reported to have AChE inhibitory activity [17]

In the present study, some AChE inhibitory activity was detected in EOs of *Boswellia species* (Table 2). At a concentration of 200 µg/mL, EO of *B. socotrana* (59.3% enzyme inhibition) exhibited higher AChE inhibitory activity than the EOs of *B. elongata* and *B. ameero* (29.6 and 41.5% enzyme inhibition, respectively). The AChE inhibitory activity of *B. socotrana* oil may be due to the presence of (E)-2,3-epoxycarene, and to p-menth-1(7)-en-2-one, a group of monoterpenoid skeletons reported to have AChE inhibitory activity [31, 32]. Pulegone, a monoterpene with p-menthane skeleton in *Mentha* spp showed AChE inhibitory activity with IC₅₀ 890 µM [32]. The reported AChE inhibition for oils of *Melissa officinalis* and *Rosmarinus officinalis* [33, 34] was stronger than the inhibitory activity of *B. socotrana* oil. This may be due to the additive or may be even synergistic effect among the components of the oils, as it was reported for 1.8 cineole, (+) – α -pinene, camphor and bornyl acetate (IC₅₀: 670, 630, >4000, >4000 µM respectively) in *Salvia lavandulaefolia* [31].

In conclusion, based on the chemical profile of the EOs, the three oleogum resins represent a new brands which can be named Soqotraen brands that are specific in their main volatile components in comparison to the old brands such as Eritrean (*B. carteri*), Omani (*B. sacra*), Somali (*B. frereana*) and Indian brands (*B. serrata*) [1]. Bioactivity guided fractionation is in progress for isolating the active compound (s) from *B. socotrana* oil.

Acknowledgments

The authors would like to thank Deutscher Akademischer Austauschdienst (DAAD) for a grant enabling the stay of Dr. Nasser A. Awadh Ali at the Leibniz Institute of Plant Biochemistry. We are also indebted to Soqotra Archipelago Conservation and Development Program (SCDP) for facilitating our mission

References

- [1] S. J. Hamm, J. C. Bleton and A. Tchaplal (2005). A chemical investigation by headspace SPME and GC-MS of volatile and semi-volatile terpenes in various olibanum samples. *Phytochemistry*. **66**, 1499-514.
- [2] G. A. Miller. and M. Morris. (2004). Ethnoflora of the Soqotra Archipelag. Charlesworth Group, Huddersfield, UK. P.457-464.
- [3] D. Kubmarawa, I. A. Ogunwande, D. A. Okorie, N. O. Olawore and A. A. Kasali, (2006). Constituents of the essential oils of *Boswellia dalzielii* Hutch. from Nigeria. *J. Ess. Oil Res.* **18**, 19-120.
- [4] G. Chiavari, G. C. Galletti, R. Piccagali and M.A. Mohammed (1991). Differential between resins of *Boswellia carteri* and *Boswellia frereana* (Frankincense of Somali origin. *J. Ess. Oil Res.* **3**, 185-186.
- [5] A. K. Kasali, A. M. Adio, A. O. Oyedeji, A. O. Eshilokun and M. Adeferwa (2002). Volatile constituents of *Boswellia serrata* Roxb (Bursaceae) bark. *Flavour Frag. J.* **17**, 462-464.
- [6] W. Wang, Y. X. Zhu, L. Z. Liy, D. K. Link, X. L. Qin and J. G. Tian (1993). Analysis of the chemical constituents of essential oil of *Boswellia carteri* birdwood from Somali origin. *Yaowu Fenxi Zazhi*, 13 170-173
- [7] S. M. Abdulwahab, E. A. Aboutalb, S. M. El-Zalabani, H. A. Fouad, H. L. De Pooter and El-Fallaha (1987). The essential oil of *Olibanum*. *Planta med.* **53**, 382-384.
- [8] H. P. T. Ammon, T. Singh and H. Safayhi (1991). Inhibition of leukotriene B4 formation in rat peritoneal neutrophils by an ethanolic extract of the gum resin exudate of *Boswellia serrata*. *Planta Med.* **57**, 203-207.
- [9] H. P. T. Ammon, H. Safayhi, T. Mack and J. Sabieraj (1993). Mechanism of antiinflammatory actions of curcumin and boswellic acids. *J Ethnopharmacol.* **38**, 113-9.
- [10] I. Gupta, A. Parihar, P. Malhotra, G. B. Singh, R. Ludtke, H. Safayhi and H. P. T. Ammon (1997). Effects of *Boswellia serrata* gum resin in patients with ulcerative colitis. *Eur. J. Med. Res.* **2**, 37-43.
- [11] I. Gupta, V. Gupta, A. Parihar, S. Gupta, R. Ludtke, H. Safayhi and H. P. T. Ammon (1998). Effects of *Boswellia serrata* gum resin in patients with bronchial asthma: results of a double-blind, placebo-controlled, 6-week clinical study. *Eur. J. Med. Res.* **3**, 511-514.
- [12] H. Safayhi, T. Mack, J. Sabieraj, M. I. Anazodo, L. R. Subramanian and H. P. T. Ammon (1992). Boswellic acids: Novel, specific, nonredox inhibitors of 5-lipoxygenase. *J. Pharmacol. Exp. Therap.* **261**, 1143-1146.
- [13] H. Safayhi, E. R. Sailer and H. P. T. Ammon (1995). Mechanism of 5-lipoxygenase inhibition by acetyl-11-keto-Boswellic acid. *Mol. Pharmacol.* **47**, 1212-1216.
- [14] M. L. Sharma, A. Khajuria, A. Kaul, S. Singh, G. B. Singh and C. K. Atal (1988). Effect of salai guggal ex- *Boswellia serrata* on cellular and humoral immune responses and leucocyte migration. *Agents Actions.* **24**, 161-164.
- [15] G. B. Singh and C. K. Atal (1986). Pharmacology of an extract of salai guggal ex-*Boswellia serrata*, a new non-steroidal anti-inflammatory agent. *Agents Actions* **18**, 407-412.
- [16] Y. Shao, C. T. Ho, C. K. Chin, V. Badmaev, W. Ma and M. T. Haung (1997). Inhibitory Activity of Boswellic acids from *Boswellia serrata* against Human Leukemia HL-60 Cells in Culture. *Plant med* **64**, 328-331.
- [17] M. Ota and P. J. Houghton (2005). Boswellic acid with acetylcholinesterase inhibitory properties from frankincense" 53rd annual congress organized by society of medicinal plants, Societa Italiana di Fitochimica Florence, Italy 21st-25th (Book of abstracts, p. 339)
- [18] T. Akihisa, K. Tabata, N. Banno, H. Tokuda, R. Nishimura, Y. Nakamura, Y. Kimura, K. Yasukawa and T. Suzuki (2006). Cancer chemopreventive effects and cytotoxic activities of the triterpene acids from the resin of *Boswellia carteri*. *Biol Pharm Bull.* **29**, 1976-86.

- [19] M. L. Gangwal and D. K. Vardhan (1995). Antifungal studies of volatile constituents of *Boswellia serrata*. *Asian J. Chem.* **7**, 675-678.
- [20] B. R. Mikhail, G. T. Maatooq, F. A. Badria and M. M. A. Amer (2003). Chemistry and immunomodulatory Activity of Frankincense Oil. *Z. Naturforsch.* **58c**, 230-238.
- [21] S. K. Lee, Z. H. Mbwambo, H. S. Chung, L. Luyengi, E. J. C. Gamez, R. G. Mehta, A. D. Kinghorn, and J. M. Pezzuto (1998). Evaluation of the antioxidant potential of natural products. *Combinatorial Chemistry and High Throughput Screening* **1**, 35-46.
- [22] H. Mohamad, F. Abas, D. Permana, N. H. Lajis, A-M. Ali, M. A. Sukari, T. T. Y. Hin, H. Kikuzaki and N. Nakatani (2004). DPPH free radical scavenger components from the fruits of *Alpinia rafflesiana* Wall. Ex Bak. (Zingiberaceae). *Z. Naturforsch.* **59c**, 811-815.
- [23] G. L. Ellman, D. K. Lourtney, V. Andres and G. Gmelin (1961). A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochem. Pharmacology* **7**, 88-95.
- [24] I. K. Rhee, M. Van de Meent, K. Ingkaninan and R. Verpoorte (2001). Screening for acetylcholinesterase inhibitors from Amaryllidaceae using silica gel thin-layer chromatography in combination with bioactivity staining. *J. Chromatogr. A* **915**, 217-223.
- [25] N. Ammar, G. Fournier and S. El-deeb (1994). The volatile constituents of *Boswellia sacra* frankincense. *J. Drug Res. Egypt* **21**, 55-88.
- [26] W. Brand-Williams, M. E. Cuvelier and C. Berset (1995). Use of a free radical method to evaluate antioxidant activity. *Lebensm.-Wiss. U Technol.* **28**, 25-30.
- [27] M.T. Baratta, H. J. D. Dorman, S.G. Deans, D.M. Biondi and G. Ruberto (1998). Chemical composition, antimicrobial and antioxidative activity of laurel, sage, rosemary, oregano and coriander essential oils. *J. Ess. Oil Res.* **10**, 618-627.
- [28] G. Ruberto and M.T. Baratta (2000). Antioxidant activity of selected essential oil components in two lipid model systems. *Food Chem.* **69**, 167-174.
- [29] N. Mimica-Duke, B. Bozin, M. Sokovic and N. Simin (2004). Antimicrobial and antioxidant activities of *Melissa officinalis* L. (Lamiaceae) essential oil. *J. Agric Food Chem.* **52**, 2485-9
- [30] N. Mimica-Dukic, B. Bozin, M. Sokovic and M. Matavulj (2003). Antimicrobial and antioxidant activities of three *Mentha* species essential oils.. *Planta med.* **69**, 413-9.
- [31] M. Miyazawa and C. Yamafuji (2005). Inhibition of acetylcholinesterase activity by bicyclic monoterpenoid. *J Agric Food Chem.* **53**, 1765-8.
- [32] M. Miyazawa, H. Watanabe and H. Kameoka (1997). Inhibition of Acetylcholinesterase Activity by Monoterpenoids with a *p*-Menthane Skeleton. *J Agric Food Chem.* **45**, 677-679
- [33] N. Perry, P. Houghton P, A Theobald, P. Jenner and E. K. Perry (2000) In-vitro inhibition of human erythrocyte acetylcholinesterase by *Salvia lavandulaefolia* essential oil and constituent terpenes. *J Pharm Pharmacol.* **52**, 895-902
- [34] N. Perry, G. Court, N. Bidet, J. court and E. Perry (1996). European herbs with cholinergic activities: potential dementia therapy. *International J. of geriatric Psychiatry.* **11**, 1063-1069

A C G
publications

© 2008 Reproduction is free for scientific studies