

Supportive information

Rec. Nat. Prod. 8:2 (2014) 121-127

**Two new antioxidant Triterpenoids from *Lonicera
quinquelocularis***

Dilfaraz Khan¹, Masood Afzal¹, Simon Woodward² and Shafiullah Khan^{1*}

¹*Department of Chemistry Gomal University Dera Ismail Khan, KPK, Pakistan*

²*School of Chemistry, Department of Organic Chemistry, University of Nottingham UK*

| Table of Contents | Page |
|-----------------------------------|-------------|
| S1: Experimental Procedure | 2 |

Experimental

Melting points were determined by using Kofler hot-stage apparatus (Reichert, Vienna, Austria). Glass capillaries were used and melting points are uncorrected. Aluminium TLC plates (20 × 20, 0.5 mm thick) precoated with silica gel 60 F₂₅₄ (20x20 cm, 0.2 mm thick; E. Merck, Darmstadt, Germany) were used for TLC to check the purity of the compounds. Column chromatography (CC) was carried out using silica gel of 230-400 mesh (E. Merck, Darmstadt, Germany). Ceric sulphate and potassium permanganate solutions were used as visualization reagents. The UV spectra (λ_{max} nm) were recorded on Shimadzu UV-2700 spectrophotometer (Shimadzu, Japan) in EtOH. Mass Spectra were recorded on Bruker TOF Mass spectrometers (Billerica, USA) using electrospray ionisation (ESI). The ¹H NMR and ¹³C NMR spectra were recorded on a Bruker DPX-400 NMR spectrometer (Billerica, USA) (400 MHz for ¹H and 100 MHz for ¹³C-NMR), using CDCl₃ as solvent. Further assignments were made by DEPT, COSY, HMQC and HMBC experiments.