

Supporting Information
***Rec. Nat. Prod.* 8:2 (2014) 195-198**
Isolation and Absolute Configuration of Boehmenan
from *Durio affinis* Becc.


Rudiyansyah^{1*}, Masriani², I. Wayan Mudianta³ and Mary J. Garson³

¹*Department of Chemistry, Faculty of Mathematics and Natural Sciences, University of Tanjungpura, Ahmad Yani Street, 78124 West Kalimantan, Indonesia*


²*Department of Chemistry, Faculty of Education, University of Tanjungpura, Ahmad Yani Street, 78124 West Kalimantan, Indonesia*

³*School of Chemistry and Molecular Biosciences, The University of Queensland, Brisbane, QLD 4072, Australia*


Table of Contents	Page
S1: Cytotoxicity effect of boehmenan and <i>D. affinis</i> ethanol extract on a normal Vero cellline	2
S2: Cytotoxicity effect of <i>D. affinis</i> ethanol extract on T47D and HeLa cells	2
S3: Cytotoxicity effect of boehmenan on T47D and HeLa cells	2
S4: Cytotoxicity effect of doxorubicin on HeLa, T47D and Vero cells	3




S1: Cytotoxicity effect of boehmenan and *D. affinis* ethanol extract on a normal Vero cellline.



S2: Cytotoxicity effect of *D. affinis* ethanol extract on T47D and HeLa cells.



S3: Cytotoxicity effect of boehmenan on T47D and HeLa cells.



S4: Cytotoxicity effect of doxorubicin on HeLa, T47D and Vero cells.