

Components from the Essential oil of *Centaurea aeolica* Guss. and *C. diluta* Aiton from Sicily, Italy

Mariem Ben Jemia¹, Felice Senatore², Maurizio Bruno^{3*} and Svetlana Bancheva⁴

¹Laboratoire des Plantes Extremophiles - Biotechnologic Center Borj-Cedria Technopark, B.P. 901, 2050 Hammam-Lif, Tunisia

² Department of Pharmacy, University of Naples "Federico II", Via D. Montesano, 49 - 80131 Naples, Italy

³ Department of Biological, Chemical and Pharmaceutical Sciences and Technologies (STeBiCEF), University of Palermo, Viale delle Scienze, Parco d'Orleans II - 90128 Palermo, Italy

⁴Institute of Biodiversity and Ecosystem Research, Bulgarian Academy of Sciences, Acad. G. Bonchev Str. 23, 1113-Sofia, Bulgaria

(Received September 18, 2012; Revised March 05, 2015; Accepted March 05, 2015)

Abstract: Volatile components from florets, leaves and stems and branches of *Centaurea aeolica* Guss. harvested in Lipari, Sicily, Italy, were analysed by gas phase chromatography (GC) and gas chromatography mass spectrometry (GC-MS). The main constituents were β -eudesmol, caryophyllene oxide, (E)-12-norcaryophyll-5-en-2-one and hexahydrofarnesylacetone in flowers, hexahydrofarnesylacetone, 2-methyloctadecane and tricosane in the leaves and hexadecanoic acid, caryophyllene oxide and β -eudesmol in the stems and branches. The analysis of the essential oil of the aerial parts of *Centaurea diluta* Aiton gave mainly fatty acids and derivatives, the main ones being hexadecanoic acid and (Z,Z)-9,12-octadecadienoic acid methyl ester.

Keywords: *Centaurea aeolica*; *Centaurea diluta*; GC-MS; volatile components; β -eudesmol; caryophyllene oxide; hexahydrofarnesylacetone; hexadecanoic acid. © 2015 ACG Publications. All rights reserved.

1. Plant Source

Centaurea aeolica Guss. ex DC. (common name: Fiordaliso of the Aeolian islands), belonging to section Alolepidae (J. Arones) Dostal, is a perennial plant, with white-tomentose stems at the youth state, after glabrescent and more or less green, up to 30-40 cm height. Lower leaves are lanceolate and deeply divided, higher cauline ones are linear. Florets (June-July) are violet and united in head inflorescences (capitula), 10-13 mm in diameter, ovoid; the bracts are without appendage. Fruits are small achenes equipped with soft and plumpy appendix (pappus), wind disseminated. Usually, it grows on dry slopes and cliffs up to 400 a. s. l. The plant is an Aeolian endemic that can be considered symbol of the flora of the Aeolian Islands. In the island is spread more or less evenly along the coast, especially along the provincial road between Malfa and Pollara (Salina Island) and White Sand Beach (Lipari Island). It is included as rare in the list of species "at risk" in the vascular flora native of Sicily

* Corresponding author: E-Mail: maurizio.bruno@unipa.it; Phone: +39-091-23897531 Fax: +39- 091- 596825

[1] and is listed among the species at lower risk of extinction in the regional Red List of Plants in Italy. This taxon was described by the Neapolitan botanist Giovanni Gussone in 1837 [2].

Centaurea diluta Aiton. (North African Knapweed) belongs to section Solstictaria (Hill) Dobrocz and it is an annual or perennial plant, native to southwestern Europe and northern Africa. The stems are up to 200 cm, erect and branched, glabrous or hairy. The lower leaves are incise-dentate, the lowermost ones – lyrate, while the upper - entire, semiamplexicaul. Capitula are solitary, pedunculate with ovoid involucre, 8-12 mm in diameter, while the bracts are brown and appressed. Appendages are shortly decurrent, orbicular-ovate, with membranous margin, irregularly fimbriate-lacerate, the apex is emarginate with a rigid, filiform spine in the notch. Florets are purple, the marginal distinctly patent. Inner achenes are with pappus, as long as achene; outer with very short pappus [3].

In continuation of our studies on *Centaurea* species of the Mediterranean area [4-5] the aerial parts of *C. aeolica* were collected in June, 2011, in White Sand Beach (Lipari Island), Sicily, Italy and the aerial parts of *C. diluta* were collected in June, 2011, near Alimena, Sicily, Italy. Typical specimens were identified by Prof. S. Bancheva, Sofia, Bulgaria and have been deposited in the Department STeBiCEF of the University of Palermo, Palermo, Italy (PAL 11/7MB and PAL 11/8MB, respectively).

2. Previous Studies

The presence of polyacetylene compounds has been previously reported in *C. diluta* [6-9], whereas neither studies on its volatile components, nor any paper on *C. aeolica* have been published.

3. Present Study

The volatile components from air-dried and ground florets (AF), leaves (AL) and stems and branches (AS) of *C. aeolica* and from the aerial parts of *C. diluta* (DAP) were isolated by hydrodistillation for 3 h, using a Clevenger-type apparatus according to the method recommended in the European Pharmacopoeia [10]. The oil was dried over anhydrous sodium sulphate and stored under N₂ at +4°C in the dark until tested and analysed. The samples yielded 1.93% (AF), 0.02% (AL), 0.02% (AS) and 0.13% (DAP) of yellow oils (w/w). The yields are moisture free basis.

GC analysis: Analytical gas chromatography was carried out on a Perkin-Elmer Sigma 115 gas chromatograph fitted with a HP-5 MS capillary column (30 m x 0.25 mm i.d.; 0.25 µm film thickness). Helium was the carrier gas (1 mL min⁻¹). Column temperature was initially kept at 40°C for 5 min, then gradually increased to 250°C at 2°C min⁻¹, held for 15 min and finally raised to 270°C at 10°C min⁻¹. Diluted samples (1/100 v/v, in *n*-hexane) of 1 µL were injected manually at 250°C, and in the splitless mode. Flame ionization detection (FID) was performed at 280°C. Analysis was also run by using a fused silica HP Innowax polyethylenglycol capillary column (50 m x 0.20 mm i.d.; 0.20 µm film thickness).

GC-MS analysis: GC-MS analysis was performed on an Agilent 6850 Ser. II apparatus, fitted with a fused silica HP-1 capillary column (30 m x 0.25 mm i.d.; 0.25 µm film thickness), coupled to an Agilent Mass Selective Detector MSD 5973; ionization voltage 70 eV; electron multiplier energy 2000 V. Gas chromatographic conditions were as reported above; transfer line temperature, 295°C. Analysis was also run by using a fused silica HP Innowax polyethylenglycol capillary column (60 m x 0.25 mm i.d.; 0.25 µm film thickness) [11].

Qualitative and quantitative analyses: Most constituents were identified by gas chromatography by comparison of their retention indices (RI) with either those of the literature [12, 13] or with those of authentic compounds available in our laboratories. The retention indices were determined in relation to a homologous series of *n*-alkanes (C₈-C₃₂) under the same operating conditions. Further identification was made by comparison of their mass spectra on both columns with either those stored in NIST 02 and Wiley 275 libraries or with mass spectra from the literature [12, 14] and our home made library. Component relative concentrations were calculated based on GC peak areas without using correction factors.

Hydrodistillation of *C. aeolica* florets (**AF**), leaves (**AL**) and stems and branches (**AS**) gave overall 49 compounds were identified: 17 in (**AF**), 17 in (**AL**) and 37 in (**AS**), respectively. The components are listed in Table 1 according to their retention indices on a HP 5MS column and are classified on the basis of their chemical structures into 8 classes. The three oils presented a different composition and the identity of the most dominant components differed notably. Furthermore, the composition of *C. aeolica* is different from that studied so far [15, 16], in fact, it is to highlight the content in oxygenated sesquiterpenes that varies between 18.8% (**AL**) and 64.0% (**AF**), higher than that described for other *Centaurea* species. The most abundant component of this class in **AL** is β-eudesmol (26.3%), followed by caryophyllene oxide (12.6%) and (*E*)-12-norcaryophyll-5-en-2-one (10.8%).

A high content of β-eudesmol and caryophyllene oxide has been also reported for *C. mucronifera* (Sect. Psephelloides, 17.4% and 5.2%, respectively) [17], *C. aladaghensis* (Sect. Cynaroides, 11.8% and 7.5%, respectively) [18], *C. gracilenta* (Sect. Arenariae, 12.8% and 6.7%, respectively) [19] and *Centaurea ensiformis* (Sect. Cheirolepis, 29.8% and 7.6%, respectively) [20]. The last two, similarly to *C. aeolica*, are devoid of germacrene D that, instead, is the main compound of *C. mucronifera* and *C. aladaghensis*. Furthermore, caryophyllene oxide (18.2%) was detected as the main product in *C. helenoides* [21] and one of the main component (5.4%) in *C. pulcherrima* var. *pulcherrima* [22], and β-eudesmol was the principal metabolite of *C. sessilis* (11.1%) and *C. armena* (11.2%) [23]. On the other hand, it is worthy to point out that (*E*)-12-norcaryophyll-5-en-2-one has never been detected in other *Centaurea* species although it was identified in species belonging to other Families [24-27].

In the **AL** oil the main fractions were represented by hydrocarbons (35.1%) and carbonylic compounds (34.5%), the last one including just hexahydrofarnesylacetone while in **AS** oil fatty acids and derivatives (31.3%) was the main fraction with hexadecanoic acid (24.7%) as principal compound.

Thirty-five compounds were determined in the essential oil of the aerial parts of *C. diluta* (**DAP**), representing 90.5% of total oil content. By far the main class is fatty acids and derivatives with hexadecanoic acid (21.3%) and (*Z,Z*)-9,12-octadecadienoic acid methyl ester (12.2%) as most abundant compounds. Hydrocarbons accounted for the 15.3% being pentacosane the most abundant one (7.1%). Terpenoids were present in low amount (2.8%) and among the carbonylic compounds (8.7%) only (*E*)-β-damascenone (3.5%) was present in good quantity. The chemical profile of *C. diluta* is quite similar to those of *C. nicaensis* (sect. Solstictaria) and *C. solstitialis* ssp. *shouwii* (sect. Solstictaria) [28], in fact, in both cases the main compounds were hexadecanoic acid (33.5%-29.4%) and (*Z,Z*)-9,12-octadecadienoic acid (28.8%-4.9%) and this fact confirm that *C. diluta* has been placed in the same section Solstictaria.

Table 1. Composition of essential oils from of *Centaurea aeolica* and *C. diluta*.

Compounds	K _i ^a	K _i ^b	AF ^c	AL ^d	AS ^e	DAP ^f	Ident. ^g
Hydrocarbons			9.4	35.1	7.2	15.3	
1-Pentadecene	1492	1544			0.2		1, 2
Cadalene	1677	2256			1.0		1, 2
Octadecane	1800	1800			0.4		1,2,3
1-Nonadecene	1893	1942	0.8				1, 2
2-Methyloctadecane	1945	1854		15.5			1, 2
Eicosane	2000	2000	2.8	3.4			1,2,3
Heneicosane	2100	2100	3.7	1.3			1,2,3
Docosane	2200	2200	0.4	t	0.9		1, 2, 3
Tricosane	2300	2300	1.2	13.0	3.9	0.3	1, 2, 3
Tetracosane	2400	2400		1.9	0.8	0.7	1, 2, 3
Pentacosane	2500	2500	0.5			7.1	1, 2, 3
Hexacosane	2600	2600				0.6	1, 2, 3
Heptacosane	2700	2700				3.0	1, 2, 3
Octacosane	2800	2800				0.4	1, 2, 3
Nonacosane	2900	2900				2.8	1, 2, 3
Hentriacontane	3100	3100				0.1	1, 2, 3

Tritriacontane			0.3	2
Carbonylic compounds		9.2	34.5	13.3
(E,E)-2,4-Heptadienal	1015	1507	t	1, 2
Phenylacetaldehyde	1045	1663		0.8
(E,E)-2,4-Decadienal	1315	1827		t 1, 2
Nonanal	1102	1399		1.4
Decanal	1206	1510		t 1,2,3
(E)-β-Damascenone	1358	1787		3.5 1, 2
Tetradecanal	1611	1934	0.5	1, 2
Pentadecanal	1715	2038	2.7	1, 2
2-Hexadecanone	1805	2203	0.5	1, 2
Hexadecanal	1817	2121	0.9	1, 2
Hexahydrofarnesylacetone	1845	2131	9.2 34.5	6.8 1.9 1, 2
2-Octadecanone	1902	2402	0.1	1, 2
Heptadecanal	1920	2241	0.8	1, 2
Octadecanal	2042	2357	0.3	1, 2
Nonadecanal	2110	2464	0.2	1, 2
(E,E)-Farnesylacetone	1920	2387		1.1 1, 2
Eicosanal	2221	2571	0.2	1, 2
Heneicosanal	2329	2682	0.3	1, 2
Sesquiterpene hydrocarbons			3.3	1.8
Cyclosativene	1363	1492	0.2	1, 2
α-Copaene	1377	1497		1.6 1, 2
β-Elemene	1387	1600		0.2 1, 2
γ-Murolene	1478	1704	3.1	
Oxygenated monoterpenes				1.0
cis-Pinocamphone	1174	1565	t	1, 2
Pulegone	1236	1662		1.0 1, 2
Oxygenated sesquiterpenes		64.0	18.8	24.1
Ledol	1565	2057	0.9	1, 2
Spathulenol	1578	2150	1.1	1, 2, 3
Caryophyllene oxide	1579	2208	12.6	4.3 7.4 1, 2, 3
β-Oopopenone	1608	2093		0.4 1, 2
Caryophylla-2(12),6(13)-dien-5β -ol (=Caryophylladienol I)	1638	2317	7.1	t 1, 2
Caryophylla-3,8(13)-dien-5β -ol	1639	2371	1.3	
Caryophylla-2(12),6(13)-dien-5β -ol (=Caryophylladienol II)	1649	2396	3.9	1.1 1, 2
β-Eudesmol	1650	2258	26.3	4.4 6.9 1, 2
Selin-11-en-4α-ol	1656	2275		8.4 2.6 1, 2
(E)-12-Norcaryophyll-5-en-2-one	1668	1984	10.8	0.6 3.6 1, 2
Valerenal	1716	2238		2.5 1, 2
β-Costol	1780	2607		0.7 1, 2
Fatty acids and derivatives		11.1	3.6	42.2
Octanoic acid	1180	2053	0.1	1, 2, 3
Nonanoic acid	1278	2159	0.8	1, 2, 3
Decanoic acid	1367	2316	1.9	1, 2, 3
Undecanoic acid	1467	2419	0.5	1, 2, 3
Dodecanoic acid	1566	2503	7.6	2.6 1, 2, 3
Tetradecanoic acid	1758	2723	3.3	2.9 0.9 1, 2, 3
Benzyl benzoate	1760	2655		0.7 1, 2, 3
Pentadecanoic acid	1873	2825	t	0.9 1, 2, 3
Hexadecenoic acid	1909	2962		0.2 1, 2, 3
Hexadecanoic acid methyl ester	1925	2208		0.5 1, 2, 3
Hexadecanoic acid	1957	2932	7.8	3.6 24.7 21.3 1, 2, 3
Heptadecanoic acid	2074	3023		0.5 1, 2, 3
(Z,Z)-9,12-Octadecadienoic acid ethyl ester	2078	2532		7.3 1, 2, 3
(Z,Z)-9,12-Octadecadienoic acid methyl ester	2098	2507		12.2 1, 2, 3

(Z,Z)-9,12-Octadecadienoic acid	2104	3160	2.5	1.0	1, 2, 3
(Z,Z,Z)-9,12,15-Octadecatrienoic acid methyl ester	2107	2583		3.8	1, 2, 3
Octadecanoic acid	2172	3172		0.7	1,2,3
Phenolic compounds					6.4
4-Vinylguaiacol	1312	2180		6.4	1,2
Others					5.9
2-Pentylfuran	1002	1243		t	1, 2
Benzothiazole	1231	1943		0.7	1, 2,3
Hexadecanol	1878	2389		5.2	1, 2, 3
Total amount of compounds		93.7	92.0	90.1	90.5

^a: HP-5 MS column; ^b: HP Innowax column; ^c: *C. aeolica* florets; ^d: *C. aeolica* leaves; ^e: *C. aeolica* stems and branches; ^f: *C. diluta* aerial parts; ^g: 1: retention index, 2: mass spectrum, 3: co-injection with authentic compound; t: traces, less than 0.05%.

Acknowledgments

The GC and GC-MS spectra were performed at the ‘C.S.I.A.S.’ of the University ‘Federico II’, Napoli. The assistance of the staff is gratefully appreciated. Authors thanks Mrs. Liliana Borzoni and Mr. Franco Peluso for the collection of plant material

References

- [1] F. M. Raimondo, L. Gianguzzi and V. Ilardi (1994). Inventario delle specie “a rischio” nella flora vascolare nativa della Sicilia, *Quad. Bot. Ambientale Appl.* **3**, 65–132.
- [2] A. P. de Candolle (1838) [1837 publ. early Jan 1838], *Prodromus Systematis Naturalis Regni Vegetabilis* **6**, 584.
- [3] J. Dostál (1976). *Centaurea* L. in TG Tutin, VH Heywood, NA Burges, DM Moore, DH Valentine, SM Walters, DA Webb, eds. *Flora Europaea* **4**, 254–301. Cambridge University Press, London, New York, Melbourne.
- [4] M. Ben Jemia, C. Formisano, S. Bancheva, M. Bruno and F. Senatore (2012). Chemical Composition of the essential oils of *Centaurea formanekii* and *C. orphanidea* ssp. *thessala*, growing wild in Greece, *Nat. Prod. Commun.* **7**, 1083-1086.
- [5] C. Formisano, D. Rigano, F. Senatore, S. Celik, M. Bruno and S. Rosselli (2008). Volatile constituents of aerial parts of three endemic *Centaurea* species from Turkey: *C. amanicola* Hub. - Mor., *C. consanguinea* DC. and *C. ptoesimopappa* Hayek and their biological activity, *Nat. Prod. Res.* **22**, 833-839.
- [6] F. Bohlmann, S. Postulka and J. Ruhnke (1958). Polyacetylene compounds. XXIV. Polyyynes of the species *Centaurea*, *Chem. Ber.* **91**, 642-656.
- [7] F. Bohlmann, K.M. Rode and C. Zdero (1966). Polyacetylene compounds. CXVII. New polyynes of the genus *Centaurea*, *Chem. Ber.* **99**, 3544-3551.
- [8] F. Bohlmann, M. Wotschokowsky, J. Laser, C. Zdero and K. D. Bach (1968). Polyacetylene compounds. CLI. Biosynthesis of tri- and tetraacetylene compounds, *Chem. Ber.* **101**, 2056-61
- [9] L. P. Christensen and G. Lam (1990). Acetylenes from *Centaurea* species, *Phytochemistry* **29**, 2753-2785.
- [10] European Pharmacopoeia 6.0, (2008). Determination of essential oils in herbal drugs, 2.8.12, pp. 251–252.
- [11] F. Senatore, N. Apostolidis Arnold and M. Bruno (2005). Volatile components of *Centaurea eryngioides* Lam. and *Centaurea iberica* Trev. var. *hermonis* Boiss. Lam., two Asteraceae growing wild in Lebanon, *Nat. Prod. Res.* **19**, 749-754.
- [12] W. Jennings and T. Shibamoto (1980). Qualitative Analysis of Flavour and Fragrance Volatiles by Glass Capillary Gas Chromatography. Academic Press, New York.
- [13] N. W. Davies (1990). Gas chromatographic retention indexes of monoterpenes and sesquiterpenes on methyl silicone and Carbowax 20M phases, *J. Chromatogr. A* **503**, 1-24.
- [14] R. P. Adams, (2007). Identification of essential oil components by gas chromatography/mass spectrometry. (4th ed.), Allured Publishing Co., Carol Stream, IL.
- [15] Y. B. Kose, B. Demirci, K. H. C. Başer and E. Yucel (2008). Composition of the essential oil of three endemic *Centaurea* species from Turkey, *J. Ess. Oil Res.* **20**, 335-338.
- [16] C. Karamenderes, B. Demirci and K. H. C. Başer (2008). Composition of essential oils of ten *Centaurea* L. taxa from Turkey, *J. Ess. Oil Res.* **20**, 342-349.

- [17] H. Dural, Y. Bağci, K. Ertuğrul, H. Demirelma, G. Flamini, P. R. Cioni and I. Morelli (2003). Essential oil composition of two endemic *Centaurea* species from Turkey, *Centaurea mucronifera* and *Centaurea chrysanthia*, collected in the same habitat, *Biochem. Syst. Ecol.* **31**, 1417-1425.
- [18] G. Flamini, M. Tebano, P. L. Cioni, Y. Bağci, H. Dural, K. Ertuğrul, T. Uysal and A. Savran (2006). A multivariate statistical approach to *Centaurea* classification using essential oil composition data of some species from Turkey, *Plant Syst. Evol.* **261**, 217-228.
- [19] C. Formisano, D. Rigano, F. Senatore, S. Bancheva, M. Bruno, A. Maggio and S. Rosselli (2011). Volatile components from aerial parts of *Centaurea gracilenta* and *C. ovina* ssp. *besserana* growing wild in Bulgaria, *Nat. Prod. Commun.* **6**, 1339-1342.
- [20] Y. B. Kose, A. Altintas, O. Tugay, T. Uysal, B. Demirci, K. Ertuğrul and K. H. C. Başer (2010). Composition of the essential oils of *Centaurea sericeae* Wagenitz and *Centaurea ensiformis* P.H. Davis from Turkey, *Asian J. Chem.* **22**, 7159-7163.
- [21] N. Yaylı, A. Yaşar, N. Yaylı, C. Albay, Y. Aşamaz, K. Coşkunçelebi and Ş. Karaoğlu (2009). Chemical composition and antimicrobial activity of essential oils from *Centaurea appendicigera* and *Centaurea helenioides*, *Pharm. Biol.* **47**, 7-12.
- [22] N. Kahrimana, G. Tosuna, N. Y. Iskender, Ş. A. Karaoglu and N. Yayli (2012). Antimicrobial activity and a comparative essential oil analysis of *Centaurea pulcherrima* Willd. var. *pulcherrima* extracted by hydrodistillation and microwave distillation, *Nat. Prod. Res.* **26**, 703-712.
- [23] N. Yaylı, A. Yaşar, C. Gülec, A. Usta, S. Kolaylı, K. Coşkunçelebi and Ş. Karaoğlu (2005). Composition and antimicrobial activity of essential oils from *Centaurea sessilis* and *Centaurea armena*, *Phytochemistry* **66**, 1741-1745.
- [24] R. Laville, C. Castel, J.-J. Filippi, C. Delbecque, A. Audran, P.-P. Garry, L. Legendre and X. Fernandez (2012). Amphilectane diterpenes from *Salvia sclarea*: biosynthetic considerations, *J. Nat. Prod.* **75**, 121-126.
- [25] J. Y. Y. Lalli, A. M. Viljoen, K. H. C. Başer, B. Demirci and T. Ozek (2006). The essential oil composition and chemotaxonomical appraisal of South African Pelargoniums (Geraniaceae), *J. Ess. Oil Res.* **18** (Spec. Ed.), 89-105.
- [26] K. H. C. Başer, T. Ozek and Z. Aytac (2000). Essential oil of *Hippomarathrum boissieri* Reuter et Hausskn, *J. Ess. Oil Res.* **12**, 231-232.
- [27] N. X. Dung, T. D. Chinh, D. D. Rang and P. A. Leclercq (1993). The constituents of the rhizome oil of *Zingiber zerumbet* (L.) Sm. from Vietnam, *J. Ess. Oil Res.* **5**, 553-5.
- [28] F. Senatore, C. Formisano, A. Raio, G. Bellone and M. Bruno (2008). Volatile components from flowers of *Centaurea nicaeensis* All., *C. parlatoris* Helder and *C. solstitialis* L., ssp. *schouwii* (D.C) Dostál wild growing in Sicily and their biological activity, *Nat. Prod. Res.* **22**, 825-832.

ACG
publications

© 2015 ACG Publications