

Cytotoxic Activity of *Laserpitium latifolium* L. Extract and Its Daucane and Phenylpropanoid Constituents

Višnja Popović^{1*}, Arne Heyerick², Silvana Petrović¹, Serge Van Calenbergh³, Izet Karalić³, Marjan Niketić⁴ and Dieter Deforce²

¹University of Belgrade - Faculty of Pharmacy, Department of Pharmacognosy, Vojvode Stepe 450, 11221 Belgrade, Serbia

²Laboratory of Pharmacognosy and Phytochemistry, Faculty of Pharmaceutical Sciences, Ghent University, Harelbekestraat 72, 9000 Ghent, Belgium

³Laboratory for Medicinal Chemistry, Faculty of Pharmaceutical Sciences, Ghent University, Harelbekestraat 72, 9000 Ghent, Belgium

⁴Natural History Museum, Njegoševa 51, 11000 Belgrade, Serbia

(Received October 17, 2012; Revised March 11, 2013; Accepted April 9, 2013)

Abstract: In the search for novel cytotoxic agents, sesquiterpenoids and phenylpropanoids have provided interesting lead compounds. From the chloroform extract of the underground parts of *Laserpitium latifolium*, the daucane sesquiterpenoids laserpitin and acetyldesoxodehydrolaserpitin, and the phenylpropanoids laserin and latifolon were isolated as the major compounds. Acetyldesoxodehydrolaserpitin is identified for the first time in the genus *Laserpitium*. Using a MTT and a sulforhodamine B (SRB) assay, the cytotoxic and antiproliferative activity of the extract and compounds laserpitin, acetyldesoxodehydrolaserpitin and laserin was tested in two closely related human breast adenocarcinoma cell lines, ie. the invasive MCF 7/6 and the non-invasive MCF 7/AZ. The IC₅₀ values of the extract were in the range of 184.72 – 397.16 µg/mL, with the most potent effect observed in the MTT test on the MCF 7/6 line. Among the tested compounds, acetyldesoxodehydrolaserpitin exerted a most potent, concentration-dependent effect (IC₅₀ values of 0.60 and 0.51 µM in the MCF 7/6 cell line, and 2.29 µM and 31.87 µM in the MCF 7/AZ cell line in the MTT and SRB test, respectively). The effect of laserin was more pronounced in the MTT test (IC₅₀ of 4.57 and 2.46 µM in the MCF 7/6 and MCF 7/AZ, respectively).

Keywords: Daucane esters; phenylpropanoids; *Laserpitium latifolium*; cytotoxic activity.

1. Plant Source

Laserpitium latifolium L. (Apiaceae) is widely distributed European species, whose underground parts (roots and rhizomes), known as Radix Gentianae albae, were used in traditional medicine for their sharp and bitter taste for preparing tonics for refreshing and strengthening, as well as for their diuretic and carminative properties [1,2]. The underground parts of *L. latifolium*, used in this investigation were collected at Mt. Gučev (West-Serbia) in October 2008. The plant material was collected and identified by Dr. Marjan Niketić, and a voucher specimen was deposited at the Natural History Museum in Belgrade under the accession number ko03102008.

* Corresponding author: E-Mail: visnjap@pharmacy.bg.ac.rs; Phone +381 11 3951 319.

2. Previous Studies

Previous chemical investigations of the extracts of underground parts of *L. latifolium* revealed the presence of daucane derivatives such as laserpitin [3,4], phenylpropanoid derivatives such as latifolon [4], while from extracts of herb were isolated sesquiterpene lactones of the guaianolide type [5].

Daucane derivatives are a relatively small group of sesquiterpenoids commonly isolated from Apiaceae species [6]. Some daucane esters exert cytotoxic activity on several human cancer cell lines [7]. The phenylpropanoid laserin was isolated and identified as the bitter compound present in carrot, *Daucus carota* L., and has been shown to exhibit cytotoxic activity against HL-60 cells [8].

3. Present Study

The underground parts of *L. latifolium* (631.96 g) were extracted twice with CHCl₃ for 48 h at room temperature, at a ratio of 100 g of powdered material in 1 L of CHCl₃. The collected filtrates were concentrated under vacuum to obtain a dark brown, gummy extract (31.78 g). Part of the crude CHCl₃ extract (4.51 g) was solubilized in MeOH and filtrated in order to remove waxes and lipophilic compounds. The obtained MeOH soluble fraction was dried under reduced pressure (4.46 g) and further eluted with a stepwise gradient system with increasing polarity of a solvent mixture composed of hexane-EtOAc (from 5:1 to pure EtOAc), followed by EtOAc-MeOH (from 1:1 to pure MeOH). In total 120 fractions were collected, in a volume of 20 mL each. Separation was followed by TLC analyses using vanillin-sulphuric acid for derivatization. The total extract and selected fractions were further analyzed by HPLC using a gradient system optimized for terpenoids. The mobile phases consisted of 0.025% HCOOH in H₂O (solvent A) and 0.025% HCOOH in 50 : 50 v/v of H₃CN and MeOH (solvent B). The elution program was 0–3 min isocratic 35.0% A (65% of B), 3–25 min linear gradient from 35 to 15% of A (85% of B), 25–28 min linear gradient from 15 to 0% of A (100% of B), 28–30 min isocratic 0% A (100% of B), 30–32 min linear gradient 0 to 35% A (65% of B) and 32–35 min isocratic 35% A (65% of B). The flow rate was 0.7 mL/min; the temperature was set at 35 °C; an injection volume of 20 µL was applied and a detection wavelength of 225 nm was used.

From the CHCl₃ extract of *L. latifolium* underground parts, four compounds were isolated as the main constituents: daucane esters laserpitin and acetyldesoxodehydrolaserpitin and phenylpropanoids laserine and latifolone. The structure of all isolated compounds was elucidated by 1D (¹H NMR, ¹³C NMR) and 2D (gHSQC, gHMBC and COSY) NMR analyses, and further confirmed by high resolution MS.

Latifolon ($\lambda_{\text{max}}=219$; 298 nm; R_t=6.47 min) was isolated as an amorphous solid, with the molecular formula C₁₁H₁₂O₄ as assigned by HR-MS (*m/z* molecular ion peak at 209.0826 (calcd for [M + H]⁺, 209.0736). Also its ¹H and ¹³C NMR spectra were corresponding to the structure of latifolon, previously isolated from *L. latifolium* root [4].

Laserpitin ($\lambda_{\text{max}}=223$ nm; R_t=18.22 min) was isolated as a white crystalline powder and with molecular formula C₂₅H₃₈O₇, assigned on the basis of the *m/z* molecular ion peak at 451.2759 (calcd for [M + H]⁺, 451.2690) in HR-MS. 1D and 2D NMR experiments allowed to identify characteristic patterns of two angelic acid esters bounded to alcohol laserol (Figure 1). Interestingly, laserpitin was the first isolated daucane derivative [6] whose structure, due to the complexity, was finally established after several revisions [3].

Acetyldesoxodehydrolaserpitin ($\lambda_{\text{max}}=223$ nm; R_t=25.84 min) was isolated as a sticky, colourless liquid. The molecular formula C₂₇H₄₀O₇ was assigned by HR-MS with *m/z* molecular ion peak at 477.2843 (calcd for [M + H]⁺, 477.2847) and *m/z* for the pseudomolecular ion peak at 499.2658 (calcd for [M + Na]⁺, 499.2666). The NMR spectra allowed characteristic patterns for two angeloyl and one acetyl ester moieties, bound to the daucane skeleton. The ¹H NMR spectrum matches with that of the daucane ester acetyldesoxodehydrolaserpitin that was previously isolated only from the roots of *Ferula tingitana* L. [9].

Laserin ($\lambda_{\text{max}}=223$ nm; R_t=20.86 min) was retrieved as a sticky, colourless liquid, with the molecular formula C₂₁H₂₆O₇ derived from the pseudomolecular ion peaks at *m/z* 413.1584 and

429.1335 (calcd for $[M + Na]^+$, 413.1600 and for $[M + K]^+$, 429.1300). 1H and ^{13}C NMR data were matching the previously reported spectra for laserin [8], isolated from *L. archangelica* Wulf. [10] and *L. garganicum* (Ten.) Bertol. [11].

In the system for terpenoids, using isolated compounds as external standards, contents of compounds were quantified, and it was determined that concentrations of laserpitin, acetyldesoxodehydrolaserpitin, laserin and latifolon were respectively 17.69, 1.02, 0.72 and 0.64 mg per g of the dry weight of *L. latifolium* underground parts.

The cytotoxic and antiproliferative activities of $CHCl_3$ *L. latifolium* extract and isolated daucanes laserpitin and acetyldesoxodehydrolaserpitin and the phenylpropanoid laserin were investigated using two *in vitro* colorimetric assays namely: MTT [12] (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide), for detecting the number of living, metabolically active cells, and SRB [13] (sulforhodamine B) for measuring the total cellular protein content. Activity was tested on the two human breast adenocarcinoma cell lines: MCF 7/6 and MCF 7/AZ, invasive and non-invasive type, respectively. The *L. latifolium* $CHCl_3$ extract and the pure compounds were tested at five concentrations between the following ranges: 7.5-750 $\mu g/mL$ for the extract and 0.5-100 μM for compounds. Vinblastine sulphate was used as a reference compound at five concentrations ranging from 0.1 to 40 nM. The results expressed as IC_{50} values (extract concentration required to inhibit 50% of the cell growth) are given in Table 1. The negative controls were carried out by replacing the test compounds by corresponding volumes of medium and these results were considered to represent 100% of cell growth.

The extract showed a concentration dependent cytotoxic and/or antiproliferative activity with IC_{50} values ranging from 184.72 to 397.16 $\mu g/mL$, and the highest effect was achieved in the MTT test on the invasive MCF 7/6 cell line (Table 1).

Among the compounds tested, acetyldesoxodehydrolaserpitin (Figure 1) was the most active compound, especially on highly invasive MCF 7/6 cell line in both assays tested. Determined IC_{50} values were in the low micromolar range (0.51 and 0.60 μM at MCF 7/6 line in the SRB and the MTT test, respectively and 2.29 μM on MCF 7/AZ in the MTT test), except IC_{50} value of 31.87 μM in SRB test at MCF 7/AZ cell line, where exerted cytotoxic and/or antiproliferative activity was weaker. The cell growth inhibition caused by acetyldesoxodehydrolaserpitin (5 μM) and the reference compound vinblastine sulphate (10 nM) are given in the Figure 2. The results are expressed as the mean values \pm SD of four independent replicates. Some compounds of the class of daucanes were previously shown to, at least in part, contribute to the cytotoxic activity of several *Ferula* species, such as elaeochytrin A, 6-antraniloyl daucane ester isolated from *F. elaeochytris* Korovin that was the most active compound in a MTT assay against two resistant cell lines of human (K562R) and mouse (DA1-3b/M2^{BCR-ABL}) leukemia (IC_{50} values 12.4 and 7.8 μM , respectively) [14]. In a panel investigation of the cytotoxicity of 16 daucane esters isolated from two *Ferula* and one *Ferulago* species on seven human tumor cell lines, 14 of them showed activity against at least one of the tumor cell lines [7]. Pallinin, a 6 α ,10 α diangeloyl daucane ester, was the most active compound with IC_{50} values in the lower molecular range against six cell lines tested [7]. Structure-activity relationship investigation showed that the β -orientation of ester group at position C(2) strongly enhances the cytotoxic activity [7]. Furthermore, the presence of an α,β -unsaturated ketone ("enone") groups in side chains of sesquiterpenes and sesquiterpene lactones increases the cytotoxicity toward tumor cells [15]. Acetyldesoxodehydrolaserpitin is a tri-ester, with two angeloyloxi groups featuring an "enone" system one of which has β -orientation at position C(2), thereby combining several structural factors contributing to the observed potent cytotoxicity in human breast adenocarcinoma cell lines.

Laserin exerted a concentration dependent effect in both tests on both cell lines, with the most potent activity in the MTT test for both cell lines (IC_{50} values 4.57 and 2.46 μM on MCF 7/6 and MCF 7/AZ, respectively). This compound and specially its erythro isomer 2-epilaserin that was isolated from carrot were characterized as cytotoxic phenylpropanoids in the human acute promyelocytic leukemia cell line HL-60 [15].

Our results suggest that both the daucane esters and phenylpropanoid laserin may be, at least in part, responsible for the demonstrated cytotoxic and antiproliferative activity of the investigated *L. latifolium* extract. Among the isolated compounds, acetyldesoxodehydrolaserpitin, whose biological

activity had not been tested before, was identified as the most potent cytotoxic and/or antiproliferative agent, especially in the invasive MCF 7/6 cell line.




Figure 1. Structure of daucane esters laserpitin (**1**) acetyl desoxodehydrolaserpitin (**2**) and phenylpropanoids laserin (**3**) and latifolon (**4**) isolated from the CHCl₃ extract of *L. latifolium* underground parts.




Figure 2. Percent of inhibition of the cell growth of the two human breast carcinoma cell lines MCF 7/6 and MCF 7/AZ in MTT and SRB tests, in comparison to the negative controls after application of 5 µM of acetyl desoxodehydrolaserpitin (ADODHL) and 10 nM of vinblastine sulphate as a reference compound. Results are expressed as the mean values ± SD of four independent replicates. Statistical significance of inhibition was determined in comparison to control groups at significance level * P ≤ 0.05; ** P ≤ 0.01; *** P ≤ 0.001.

Table 1. *In vitro* cytotoxic activity expressed as IC₅₀ values on MCF 7/6 and MCF 7/AZ cell lines for *L. latifolium* CHCl₃ extract, compounds and reference compound (vinblastine sulphate) in MTT and SRB assays.

Compound	MTT		SRB	
	MCF 7/6	MCF 7/AZ	MCF 7/6	MCF 7/AZ
1^a	31.80	87.13	>100	>100
2^a	0.60	2.29	0.51	31.87
3^a	4.57	2.46	62.04	7.36
Vinblastine^a	5.98×10^{-3}	15.41×10^{-3}	4.11×10^{-3}	3.66×10^{-3}
Extract^b	184.72	208.94	208.85	397.16

^aIC₅₀ values for laserpitin (**1**), acetyl desoxodehydrolaserpitin (**2**), laserin (**3**) and , vinblastine (reference compound) are given in µM;

^bIC₅₀ values for *L. latifolium* CHCl₃ extract are given in µg/ml.

Acknowledgment

This research was supported by a Basileus grant (Application № 1011092), for a research stay of PhD student Višnja Popović in Laboratory for Pharmacognosy and Phytochemistry, Faculty of Pharmaceutical Sciences, Ghent University, as a part of the Erasmus Mundus Action 2 program of European Comission, and Ministry of Education and Science of Republic of Serbia (Grant № 173021). Statistical analyses were performed with the kind help of Prof. Dr. Dragić Banković, Faculty of Mathematics, University of Kragujevac, Serbia.

References

- [1] G. Hegi (1906). *Laserpitium* L., In: *Illustrierte Flora von Mitteleuropa*, Band V (2), ed: H. Beger, A. Pichler's Witwe & Sohn, Wien, Austria, pp.1472-1501. [In German]
- [2] W. Schneider (1974). Pfanzliche Drogen. Govi-Verlag GmbH-Pharmazeutischer Verlag, Frankfurt a. M., Germany. [In German]
- [3] M. Holub, Z. Samek, V. Herout and F. Šorm (1967). Constitution of laserpitine, a sesquiterpenic compound from *Laserpitium latifolium* root, *Collect. Czech. Chem. Commun.* **32**, 591-609.
- [4] P. Moldt, U. W. Smitt and S. Brøgger Christensen (1987). A new sesquiterpene from *Laserpitium latifolium*, *J. Nat. Prod.* **50**, 974-975.
- [5] M. Djermanović, M. Stefanović, V. Djermanović and M. Milovanović (1994). Sesquiterpene lactones from *Laserpitium latifolium*, *Fitoterapia*, **65**, 556.
- [6] E. L. Ghisalberti (1994). The daucane (carotene) class of sesquiterpenes, *Phytochemistry*. **37**, 597-623.
- [7] S. Dall'Acqua, M. A. Linardi, F. Maggi, M. Nicoletti, V. Petitto, G. Innocenti, G. Basso and G. Viola (2011). Natural daucane sesquiterpenes with antiproliferative and proapoptotic activity against human tumor cells, *Bioorgan. Med. Chem.* **19**, 5876-5885.
- [8] R. L. Yang, Z. H. Yan and Y. Lu (2008). Cytotoxic phenylpropanoids from carrot, *J. Agr. Food Chem.* **56**, 3024-3027.
- [9] M. Miski and T. J. Mabry (1986). New daucane esters from *Ferula tingitana*, *J. Nat. Prod.* **49**, 657-660.
- [10] M. Holub and Z. Samek (1973). Structure of archangenolide, a sesquiterpenic lactone from *Laserpitium archangelica*, *Collect. Czech. Chem. Commun.* **38**, 731-738.
- [11] G. Appendino, M. G. Valle, R. Caniato and E. M. Cappelletti (1986). Sesquiterpene lactones from *Laserpitium garganicum*, *Phytochemistry*. **25**, 1747-1749.
- [12] T. Mosmann (1983). Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays, *J. Immunol. Methods*. **65**, 55-63.
- [13] P. Skehan, R. Stroeng, D. Scudiero, A. Monks, J. McMahon, D. Vistica, J. T. Warren, H. Bokesch, S. Kenney and M. R. Boyd (1990). New colorimetric cytotoxicity assay for anticancer drug screening, *J. Natl. Cancer Inst.* **82**, 1107-1112.
- [14] R. Alkhateib, T. Hennebelle, S. Joha, T. Idziorek, C. Preudhomme, B. Quesnel, S. Sahpaz and F. Bailleul (2008). Activity of elaeochytrin A from *Ferula elaeochytris* on leukemia cell lines, *Phytochemistry*. **69**, 2979-2983.
- [15] A. Ghantous, H. Gali-Muhtasib, H. Vuorela, N. A. Saliba and N. Darwiche (2010). What made sesquiterpene lactones reach cancer clinical trials, *Drug Discov. Today*. **15**, 668-678.

A C G
publications

© 2013 Reproduction is free for scientific studies