

## New Cytotoxic Metabolites from Pathogenic Fungus *Cylindrocarpon destructans* Associated with *Meconopsis grandis*

Dilfaraz Khan <sup>1,2\*</sup>, Shah Iram Niaz <sup>1</sup>, Kamran Tahir <sup>2</sup>, Hidayat Ullah Khan <sup>3</sup>, Muhammad Haroon <sup>4</sup>, Shahid Ullah Khan <sup>5</sup>, Syed Badshah <sup>2</sup>, Shafiullah Khan <sup>2</sup>, Muafia Jabeen <sup>6</sup> and Liu Lan <sup>1</sup>

<sup>1</sup>School of Marine Sciences, Sun Yat-sen University, Guangzhou 510006, China

<sup>2</sup>Institute of Chemical Sciences, Gomal University Dera Ismail Khan 29050, KPK, Pakistan

<sup>3</sup>Department of Chemistry, University of Science and Technology, Bannu 28100 KPK, Pakistan

<sup>4</sup>College of Chemical and Biological Engineering Zhejiang University, Hangzhou 310027, China

<sup>5</sup>National Key laboratory of Crop Genetics and Improvement, Huazhong Agriculture University, Wuhan, China

<sup>6</sup>Department of Pharmacy, COMSATS, Lahore, Pakistan

(Received January 13, 2018; Revised March 28, 2018; Accepted April 01, 2018)

**Abstract:** Phytochemical investigation on pathogenic fungus *Cylindrocarpon destructans* isolated from *Meconopsis grandis* plant led to the isolation of two new isochromene derivatives namely, 6,8-dimethoxy-3-methyl-3,4-dihydrobenzoisochromene-4,9,10-triol (**1**) and 3,5,6-trihydroxy-4-methylbenzoisochromene-9,11-dione (**2**) along with four known compounds (**3-6**). The structures of these compounds were elucidated by 1D and 2D NMR and mass spectroscopic data analysis. The isolated compounds were evaluated for cytotoxic activity. The compounds **1-4** showed good inhibition against the growth of cell lines MCF-7 and PC-3. Compounds **5-6** showed minimum inhibitory effect of cancerous cell lines growth.

**Keywords:** Pathogenic fungus; *Cylindrocarpon destructans*; *Meconopsis grandis*; cytotoxic activity © 2018 ACG Publications. All rights reserved.

### 1. Introduction

Species of *Cylindrocarpon* Wollenw. are common and may be isolated as soil inhabitants, saprobes on dead plant material, root colonizers or pathogens, or weak pathogens of various herbaceous and woody plants [1]. *Cylindrocarpon destructans* (Zinns.) [anamorph of *Neonectriaradicicola*] and *C. obtusisporum* have been reported to cause the root rots of various hosts [2, 3], and a black foot disease of grapevines [4-6]. *C. destructans* (*C. radicola*) has frequently been reported to cause decay of woody seedlings, especially conifers, and many other hosts as well [2]. Generally this fungus is not severe in its pathogenicity and has been regarded in many cases as the wound infectious fungus or the secondary invader.

\* Corresponding author: E-Mail: [dilfarazkhan@gu.edu.pk](mailto:dilfarazkhan@gu.edu.pk)

*Cylindrocarpon* species have been rarely associated with human disease. They are known to cause post traumatic keratitis [7, 8] and, have been implicated in mycetoma following injury [9, 10], athlete's foot [11], peritonitis in a case of continuous ambulatory peritoneal dialysis [10], localized invasive lesion in a case of AML [12], disseminate disinfection in neutropenic patients [13]. The human infecting species include *C. cyaneascens*, *C. destructans*, *C. lichenicola* and *C. vaginae* [14].

Previous phytochemical studies on *Cylindrocarpon* species have resulted in the isolation of Orsenol, Orsellinic acid, Ilicicolin (A-F), Ascochlorin, fatty acids [15], Cylindrols [16], Cylindrocyclin [17], Colletorine, Colletochlorin and Curvularine derivatives [18].

Only few metabolites such as radicicol and radicicolin [19] have been reported from *Cylindrocarpon destructans* (*C. radicola*). Herein we reported the isolation and structure elucidation of two new compounds (**1-2**) and four known compounds (**3-6**) from *Cylindrocarpon destructans* (*C. radicola*). The isolated compounds were screened for cytotoxic activities.

## 2. Materials and Methods

### 2.1. General

Optical rotations were measured with an Abbemat 300 spectrometer. NMR spectra were recorded with a Bruker Avance (400 MHz for  $^1\text{H}$  and 100 MHz for  $^{13}\text{C}$  spectrometer; Bruker Corporation, Switzerland). HRESIMS spectra were recorded using an LTQ-Orbitrap LC-MS spectrometer (Thermo Corporation, USA). UV spectra were recorded on a Blue Star A spectrophotometer. Thin layer chromatography silica gel GF<sub>254</sub> (Qingdao Marine Chemical, Factory PR China) were used for TLC. Sephadex LH-20 (Amersham Pharmacia) and Silica gel (100-200, 200-300, and 300-400 mesh, Qingdao Marine Chemical Factory, Qingdao, PR China) were used for column chromatography. The reagents in the research process were analytical grade from Guangzhou chemical reagent factory.

### 2.2. Fungal Material

The fungal strain *Cylindrocarpon destructans* (or *C. radicola*) (Gen accession number KC904953) was isolated from fresh roots of *Meconopsis grandis* (Tibetan Blue puppy) in Tibetan Plateau near to Damxung, Tibet, China. The fungus was identified as using morphological characteristics and ITS region. A voucher specimen (DH 29) has been preserved on PDA at 4°C at the school of marine science, Sun Yat-Sen University.

### 2.3. Fermentation, Extraction and Fractionation

The fungal strain was cultured for 6 days at 28 °C in Petri dishes containing Potato dextrose agar. The agar supported mycelia were then cut and transferred to 1000 mL Erlenmeyer flasks containing 500 mL potato dextrose broth (12 gm of PDB dissolved in 500 mL of 3% saline water) and then incubated at 28 °C for 5 days with continuous shaking on shaker at 150 rpm. Then 10 mL of the fungal broth were added into rice medium (110 bottles each 1000 mL Erlenmeyer flasks, each containing 60 g rice in 80 mL of 3% saline water) and were incubated for 30 days under static conditions and light. After incubation, the mycelia cultivated rice medium were crushed and extracted three times with methanol. The methanol extract was concentrated with rotary evaporator to get methanolic crude which was then suspended in 20 % MeOH-H<sub>2</sub>O. The suspension was then fractionated with n-hexane, chloroform, ethyl acetate to get the corresponding n-hexane (15 g), Chloroform (26 g) and ethyl acetate (44 g) fractions. The chloroform extract was separated into 7 sub-fractions (J-P) by silica gel column chromatography by gradient elution of petroleum ether/CH<sub>2</sub>Cl<sub>2</sub> (from 90:10 to 00:100) and then CH<sub>2</sub>Cl<sub>2</sub>/MeOH (from 100:00 to 00:100). Ethyl acetate fraction was further fractionated over silica gel column chromatography with gradient elution of petroleum ether/EtOAc (100:00 – 00:100) and then EtOAc/MeOH (100:00 to 00:100) to get 9 fractions (Q-Y).

### 3.4. Purification of the Compounds

Fraction **M** was further separated over silica gel CC eluted with pet ether/CH<sub>2</sub>Cl<sub>2</sub> (90:10 to 00:100) and then CH<sub>2</sub>Cl<sub>2</sub>/MeOH (5 and 10% MeOH in CH<sub>2</sub>Cl<sub>2</sub>) to afford 5 subfractions (M-1 to M-5). Fraction M-4 was purified by sephadex LH-20 with CH<sub>2</sub>Cl<sub>2</sub>: MeOH (v/v; 1:1) and then by semipreparative HPLC (65 % MeOH-H<sub>2</sub>O flow rate 1.5 ml /min; C<sub>18</sub>, 10×250 mm, 5 μm) to afford two compounds (**5** and **6**). Fraction **P** was further separated over silica gel CC eluted with pet ether/CH<sub>2</sub>Cl<sub>2</sub> (90:10 to 00:100) and then CH<sub>2</sub>Cl<sub>2</sub>/MeOH (5 and 15% MeOH in CH<sub>2</sub>Cl<sub>2</sub>) to afford 4 subfractions (P-1 to P-4). Fraction P-3 was purified by HPLC (70 % MeOH-H<sub>2</sub>O flow rate 1.5 ml /min; C<sub>18</sub>, 10×250 mm, 5 μm) to afford compounds **2** (13 mg).

Fraction **S** was subjected to series of silica gel chromatographic separation by gradient elution with pet ether/EtOAc and then EtOAc/MeOH and then to HPLC (70% MeOH-H<sub>2</sub>O, flow rate 1.0 ml/min; C<sub>18</sub>; 10×250 mm, 5 μm) to get a pure compound **1**. Fraction **T** was rechromatographed over Silica gel CC by gradient elution with EtOAc/MeOH (100:00 to 00:100) to get 7 sub-fractions (T-1 to T-7). Fraction T-6 was applied to reverse phase silica gel column MeOH/H<sub>2</sub>O (70:30) and then to semipreparative HPLC with (80% CH<sub>3</sub>CN-H<sub>2</sub>O flow rate 1.5 ml /min; C<sub>18</sub>, 10×250 mm, 5 μm) to yield two pure compounds (**3** and **4**).

3.4.1. *6,8-dimethoxy-3-methyl-3,4-dihydrobenzoisochromene-4,9,10-triol (1)*: Yellow powder; UV (MeOH) $\lambda_{\max}$  (log $\epsilon$ ): 230 (4.78), 284 (4.12), 304 (3.98), 338 (3.86) nm; HRESIMS *m/z*: 307.1140 [M+H]<sup>+</sup> (calcd for C<sub>16</sub>H<sub>16</sub>O<sub>6</sub>, 307.1136). <sup>1</sup>H NMR and <sup>13</sup>C NMR data see Table 1.

3.4.2. *3,5,6-trihydroxy-4-methylbenzoisochromene-9,11-dione (2)*: White amorphous powder;  $[\alpha]_D^{20}$  +65, (c 0.025, MeOH); UV (MeOH) $\lambda_{\max}$  (log $\epsilon$ ): 278 (4.12), 243 (3.41), 304 (3.98), 338 (3.86) nm; HRESIMS *m/z*: 261.0363 [M+H]<sup>+</sup> (calcd for C<sub>16</sub>H<sub>16</sub>O<sub>6</sub>, 261.0354). <sup>1</sup>H NMR and <sup>13</sup>C NMR data see Table 1.

### 3.5. Cytotoxicity MTT Assay

#### 3.5.1. Preparation of Cell Lines and Cell Cultures


MCF-7, PC-3 and WI-38 cells were seeded in culture plates containing MEM, DMEM Media, glutamine (2mM), penicillin (100 Units/mL) and streptomycin (100 μg/mL) accompanied with 10 % heat-inactivated fetal bovine serum under humidified atmosphere at constant temperature of 37 °C in 5% CO<sub>2</sub> incubator. After achieving a monolayer with 80% confluence of both adherent cell lines, the cells were cultured in 96-well plates at a seeding density of 5000-10000/well to start cytotoxicity experiments.

#### 3.5.2. MTT Assays Procedure

The method is based on the principal that was previously described by [20] with some modifications to perform cytotoxicity assay cells. The MCF-7, PC-3 and WI-38 cells were cultured for 24 thin Micro plate Elisa Reader for 96 Well Plate (ELX 800) BioTek USA. Different concentrations of test compounds (10, 1, 0.1, 0.01, 0.001 μM) were inoculated in test wells while control and blank wells were also prepared using cell with media and dimethyl sulfoxide (DMSO), respectively. The plates were then incubated for 48 h at 37 °C. After that cells were fixed with 50 μL of MTT solution at 37 °C for 1 h. The plates were washed 5 times with PBS (Phosphate-Buffered Saline). After that the MTT solution was removed and added 100 μL of DMSO The absorbance was measured at 570 nm subtracting the background measurement. Results were reported as mean of three independent experiments (± SEM) and expressed as percent inhibitions calculated by the formula. Inhibition (%) = [100 - (abs of test comp/abs of control) × 100]. IC<sub>50</sub> values of selected compounds exhibiting >50% activity at 0.5 mM were calculated after suitable dilutions.

### 3. Results and Discussion

The fungus *Cylindrocarpon* sp. DH 29 was cultured in rice solid medium for 30 days. The  $\text{CHCl}_3$  and EtOAc fraction were repeatedly fractionated and purified by using silica gel column chromatography, sephadex LH-20, reverse phase silica column and HPLC to obtain two new compound (**1-2**) and four known compounds (**3-6**) Figure 1.



**Figure 1.** Structures of compounds **1-6**

Compound **1** was isolated as yellow amorphous powder. Its molecular formula was deduced on the basis of HRESIMS as  $\text{C}_{16}\text{H}_{18}\text{O}_6$  ( $m/z$ : 307.1140  $[\text{M}+\text{H}]^+$ ) with eight degrees of unsaturation. The  $^1\text{H}$ NMR spectrum of compound **1** displayed a couple of doublets resonated at  $\delta$  6.10 (1H,  $J = 2.1$ ) and 6.56 (1H,  $J = 2.1$ ) assignable to aromatic protons and a pair of doublets resonated at  $\delta$  4.78 (1H,  $J = 15.6$ ) and 5.20 (1H,  $J = 15.6$ ) were attributed to two geminal oxygenated methylenic protons. The quartet of doublet at  $\delta$  3.71 (1H,  $J = 6.3, 1.7$ ) and a doublet resonated at  $\delta$  3.83 (1H,  $J = 1.7$ ) were characteristics of two vicinal methine protons. COSY correlation (Figure S6) between the two H-1 proton established their geminal relationship while the correlation between H-3 and H-4 suggested their vicinal relationship. The two methoxy groups exhibited signals at  $\delta$  3.43 (3H, s) and 4.09 (3H, s) and the doublet at  $\delta$  1.19 (3H,  $J = 6.3$ ) was due to methyl protons. The  $^{13}\text{C}$ NMR spectrum along with DEPT experiment showed the presence of four tertiary carbons that include two aromatic and two oxygenated methine carbons, one oxygenated methylene carbon, three primary carbons including two methoxy and one methyl carbons and eight quaternary carbons. The HMBC (Figure S8) correlations from H-5 to C-6, C-8a, C-10a and from H-7 to C-8 and C-8a were in consistent with the position of these protons. Finally the structure of compound **1** was assigned as 6,8-dimethoxy-3-methyl-3,4-dihydrobenzoisochromene-4,9,10-triol on the basis of the spectral data that share common structural unit

with the compounds available in literature [21, 22). The structure of compound **1** was further confirmed by 2D experiments and mass spectra (Figure **S6- S8**).

**Table 1.**  $^1\text{H}$  (400 MHz) and  $^{13}\text{C}$  (100MHz) NMR data for Compound **1** and Compound **2** (MeOD)

Position	1		2		
	$\delta\text{ H (J in Hz)}$		$\delta\text{ C}$	$\delta\text{ H (J in Hz)}$	$\delta\text{ C}$
1	4.78 (1H, d, $J = 15.6$ ), 5.20 (1H, d, $J = 15.6$ )		66.52		97.0
2				6.81 (1H, s)	117.3
3	3.71 (1H, qd, $J = 6.3, 1.7$ )		73.97		166.7
4	3.83 (1H, d, $J = 1.7$ )		66.02		112.4
4a			134.48		151.7
5	6.56 (1H, d, $J = 2.1$ )		99.13		165.4
6			149.2		164.7
7	6.10 (1H, d, $J = 2.1$ )		97.62	6.42 (1H, s)	99.5
8			136.44		91.7
8a			109.95		136.9
9			157.14		167.9
9a			113.54		
10			157.10		
10a			124.88		
11	1.19 (3H, d, $J = 6.3$ )		15.02		167.6
12				2.87 (3H, s)	25.05
6-OCH <sub>3</sub>	4.09 (3H, s)		55.46		
8-OCH <sub>3</sub>	3.43 (3H, s)		53.98		

Compound **2** was isolated as white amorphous powder. The HREIMS spectra gave  $[\text{M}+\text{H}]^+$  peak at  $m/z$  261.0363 consistent with the molecular formula of  $\text{C}_{13}\text{H}_8\text{O}_6$  corresponding to nine degrees of unsaturation. The  $^1\text{H}$ NMR spectrum of compound **2** displayed a couple of singlets resonated at 6.81 and 6.42 assignable to aromatic protons. A singlet resonated at 2.87 was characteristics of methyl protons.  $^{13}\text{C}$  NMR spectrum exhibited signals for 13 carbon that were assigned by DEPT as two aromatic methine carbons, one methyl carbon and ten quaternary carbons that include two carbonyl carbon and eight fully substituted aromatic carbons. The aromatic protons were found to show a long range HMBC correlations with the carbonyl carbons resonated at 167.9 and 167.6 indicated the connectivity of these protons at C-2 and C-7 respectively. On the basis of these analysis the structure of compound **2** was assigned as 3,5,6-trihydroxy-4-methylbenzoisochromene-9,11-dione that has close resemblance with the available literature (Lin *et al.* 2012). The known compounds **3-6** were identified by comparison of their spectral data with the literature available and include *Preussiafuran A* (**3**), *Preussiafuran B* (**4**) [24], *4,5,6-trihydroxy-7-methyl-1,3-dihydroisobenzofuran* (**5**), *4,5,6-trihydroxy-7-methylphthalide* (**6**) [25].

The compounds **1-6** were evaluated for their *in vitro* cytotoxicity (Table 2). All the compounds were analyzed against two different cancer cell lines cell lines MCF-7 and PC-3 was determined by MTT assay. Tamoxifen was used as standard drug. The compounds **1, 2, 3** and **4** inhibit MCF-7 with  $\text{IC}_{50}$  values of  $05.37 \pm 1.03$ ,  $09.02 \pm 1.20$ ,  $08.04 \pm 2.10$ ,  $10.04 \pm 0.20$  and Cell viability (%)  $03.02 \pm 2.11$ ,  $6.07 \pm 0.30$ ,  $04.14 \pm 0.81$ ,  $05.21 \pm 0.01$  ( $\mu\text{M}$ ). The PC-3 with  $\text{IC}_{50}$  values of  $04.12 \pm 0.38$ ,  $12.07 \pm 0.11$ ,  $09.41 \pm 1.81$  and  $08.13 \pm 2.31$  ( $\mu\text{M}$ ) and Cell viability (%)  $2.04 \pm 0.09$ ,  $07.14 \pm 1.2$ ,  $05.41 \pm 07$  and  $04.71 \pm 0.62$  respectively. The compounds **5-6** showed minimum inhibitory effect of cancerous cell lines growth.

**Table 2.** Cytotoxicity of compounds 1-6, IC<sub>50</sub> values (μM) and cell survival (values ± standard Deviation)\*

Compounds	IC <sub>50</sub> ± SEM <sup>a</sup>			
	MCF-7		PC-3	
	IC <sub>50</sub> (μM)	Cell survival (%)	IC <sub>50</sub> (μM)	Cell survival (%)
<b>1</b>	5.37± 1.03	3.02± 2.11	4.12± 0.38	2.04± 0.09
<b>2</b>	9.02 ± 1.20	6.07 ± 0.30	12.07 ± 0.11	7.14 ± 1.2
<b>3</b>	8.04 ± 2.10	4.14± 0.81	9.41 ± 1.81	5.41 ± 07
<b>4</b>	10.04 ± 0.20	5.21± 0.01	8.13± 2.31	4.71 ± 0.62
<b>5</b>	24.18± 1.08	9.02 0.05	28.07± 0.02	7.02± 0.20
<b>6</b>	18.04± 0.10	10.27 ± 2.47	21.04 ± 0.20	8.04 ± 0.22
<b>Tamoxifen<sup>b</sup></b>	0.00053 ± 0.0004		0.0036 ± 0.0007	

\* Values shown are mean ± SEM, no. of experiments = 3

<sup>a</sup> Inhibitory effect of cancerous cell lines growth

<sup>b</sup> standard drug

### Acknowledgments

Thanks to Guangdong Provincial ministry of Education for financial support during post doctorate fellowship at Sun Yat-Sen University, China. Instrumental Analysis and Research Center of Sun Yat-Sen University is specially acknowledged for their assistance in acquisition of mass spectra regarding this work.

### Supplementary material

Supporting Information accompanies this paper on <http://www.acgpubs.org/RNP>

### Conflict of interest

The authors declare no conflict of interest.

### ORCID

Dilfaraz Khan: [0000-0003-0971-0124](https://orcid.org/0000-0003-0971-0124)

Shah Iram Niaz: [0000-0001-7083-9790](https://orcid.org/0000-0001-7083-9790)

Kamran Tahir: [0000-0001-9505-1414](https://orcid.org/0000-0001-9505-1414)

Hidayat Ullah Khan: [0000-0002-3099-6983](https://orcid.org/0000-0002-3099-6983)

Muhammad Haroon: [0000-0003-0763-448X](https://orcid.org/0000-0003-0763-448X)

Shahid Ullah Khan: [0000-0002-3082-6079](https://orcid.org/0000-0002-3082-6079)

Syed Badshah: [0000-0001-8888-7387](https://orcid.org/0000-0001-8888-7387)

Shafiullah Khan: [0000-0003-4306-4752](https://orcid.org/0000-0003-4306-4752)

Muafia Jabeen: [0000-0002-2610-0435](https://orcid.org/0000-0002-2610-0435)

Lui Lan: [0000-0003-2765-4015](https://orcid.org/0000-0003-2765-4015)

### References

- [1] D. Brayford (1993) *Cylindrocarpon*. In: Methods for research on soil-borne phytopathogenic fungi. (Singleton LL, Mihail JD, Rush M, eds). APS Press, Saint. Paul, U.S.A.: 103–106.
- [2] C. Booth (1966). The Genus *Cylindrocarpon*. Mycological Papers No. 104, Commonwealth Mycological Institute, England, pp. 35-37.
- [3] K. A. Seifert, C. R. McMullen, D. Yee, R. D. Reeleder and K. F. Dobinson (2003). Molecular differentiation and detection of ginseng-adapted isolates of the root rot fungus *Cylindrocarpon destructans*, *Phytopathology*, **93**, 1533–1542.
- [4] D. R. Maluta and P. Larignon (1991). Pied-noir: Mieuxvautprévenir, *Viticulture* **11**, 71–72.

- [5] S. Grasso and S. L. di Magnano (1975). Infections of *Cylindrocarpon obtusisporum* on grape vines in Sicily, *Vitis* **14**, 36–39.
- [6] H. J. Scheck, S. J. Vasquez, D. Fogle and W. D. Gubler (1998). Grape growers report losses to black-foot and grapevine decline, *Calif. Agric.* **52**, 19–23.
- [7] J. Kaliyamurthy, C. Jesudasan, D. A. Prasanth and P. A. Thomas (2006). Keratitis due to *Cylindrocarpon lichenicola*, *J. Postgrad. Med.* **52**, 155–157.
- [8] T. Gaujoux, E. Borsali, J. C. Gavrillov, O. Touzeau, P. Goldschmidt and M. O. Despiau (2012). Fungal keratitis caused by *Cylindrocarpon lichenicola*, *J. Fr. Ophthalmo.* **35** (5), 356.
- [9] D. E. Zoutman and L. Singler (1991). Mycetoma of the foot caused by *Cylindrocarpon destructans*, *J. Clin. Microbiol.* **29** (9), 1855–1859.
- [10] R. Sharma, C. K. Farmer, W. R. Gransden and C. S. Ogg (1988). Peritonitis in continuous ambulatory peritoneal dialysis due to *Cylindrocarpon lichenicola* infection, *Nephrol. Dial. Transplant.* **13**, 2662–2664.
- [11] B. Lancy, C. Blanc and J. Lapalu (1985). *Cylindrocarpon*: Anewathlete's foot agent, *Bull. Trimest. Soc. Mycol. Fr.* **14**, 73–76.
- [12] P. C. Iwen, S. R. Tarantolo, D. A. Sutton, M. G. Rinaldi and S. H. Hinrichs (2000). Cutaneous infections caused by *Cylindrocarpon lichenicolain* a patient with acute leukemia, *J. Clin. Microbiol.* **38** (9), 3375–3378.
- [13] E. A. James, D. K. Orchar, P. H. Mcwhinney, D. W. Warnock, E. M. Johnson, A. B. Mehta (1997). Disseminated infection due to *Cylindrocarpon lichenicola* in a patient with acute myeloid leukemia, *J. Infect.* **34**, 65–67.
- [14] C. Booth, Y. M. Clayto and M. Usherwood (1985). *Cylindrocarpon* species associated with mycotic keratitis, *Proceed. Ind. Acad. Sci.* **2** (3), 433–436
- [15] K. Mio, F. Takashi, U. Ryuji, N. Kenichi, M. Rokuro and T. Hiroshi (2013). A new ascochlorin derivative from *Cylindrocarpon* sp. FKI-4602, *J. Antibiot.* **66**, 23–29.
- [16] B. S. Sheo, G. B. Richard, F. B. Gerald, C. Carmen, B. G. Jackson, A. G. Michael, H. Karst, G. J. Rosalind, M. L. Jerrold, B. L. Russell, C. S. Keith and L. Z. Deborah (1996). Chemistry and biology of cylindrols: Novel inhibitors of ras farnesyl-protein transferase from *Cylindrocarpon lucidum*, *J. Org. Chem.* **61**, 7727-37.
- [17] W. Daniela, E. Gilda, S. Olov and A. Timm (2006). Cylindrocyclin A, a new cytotoxic cyclopeptide from *Cylindrocarpon* sp., *J. Antibiot.* **59** (8), 495–499.
- [18] G. R. Margarita, T. Cristina, R. Jaime, L. Mauricio and S. H. Guillermo (2005). Bioactive metabolites from the fungus *Nectriagalligena*, the main apple canker agent in Chile, *J. Agric. Food Chem.* **53**, 7701-7708.
- [19] G. Evans and N. H. White. (1966). Radicolin and radicol, two new antibiotics produced by *Cylindrocarpon radicola*, *Trans. Br. Mycol. Soc.* **49** (4), 563-576
- [20] T. Mosmann (1983). Rapid colorimetric assay for cellular growth and survival: Application to proliferation and cytotoxicity assays, *J. Immunol. Methods.* **16**, 55-63.
- [21] O. Jaturapat, M. Isaka, N. L. Hywel-Jones, Y. Lertwerawat, S. Kamchonwongpaisan, K. Klrtikara, M. Tanti-charoen and Y. Thebtaranonth (2001). Bioanthracenes from the insect pathogenic fungus *Cordycepspseudo militaris* BCC1 620, I, *J. Antibiot.* **54** (1), 29-35
- [22] M. Isaka, P. Kongsaree and Y. Thebtaranonth (2001). Bioanthracenes from the insect pathogen fungus *Cordycepspseudo militaris* BCC1 620, II, *J. Antibiot.* **54** (1), 36-43.
- [23] H. Lin, Y. W. Zhang, Y. Hua, Y. L. Bao, Y. Wu, S. L. Gun, C. L. Yu, Y. X. Huang, E. B. Wang, H. Y. Jiang and Y. X. Li (2014). Three new compounds from the stem bark of *Juglans mandshurica*, *J. Asian Nat. Prod. Res.* **16** (8), 819.
- [24] F. M. Talontsi, M. Lamshöft, C. Douanla-Meli, S. F. Kouam and M. Spiteller (2014). Antiplasmodial and cytotoxic dibenzofurans from *Preussiasp. Harbouredin enantiachlorantha oliv*, *Fitoterapia* **93**, 233–238.
- [25] A. Z. Huang, Y. Zhu, X. L. Guan, K. Tian, J. M. Guo, H. B. Wang and G. M. Fu (2012). A novel antioxidant isobenzofuranone derivative from fungus *Cephalosporium* sp. AL031, *Molecules* **17**, 4219-4224.