

A New Acylated Benzyl Alcohol Glucoside from *Syzygium austroyunnanense*

Feng Li[✉], Shaohua Xu[✉], Yan Zhao[✉], Mengjia Li[✉], Wen Xu[✉],
Yikao Hu[✉], Dingli Zhang[✉] and Yong Zhao^{✉*}

College of Chemistry and Chemical Engineering, Yunnan Normal University, Kunming 650500, China

(Received April 16, 2018; Revised June 08, 2018; June 21, 2018)

Abstract: A new acylated benzyl alcohol glucoside, benzyl alcohol 3',6'-di-*O*-galloyl- β -glucopyranoside (**1**), together with a known analogue, 6'-*O*-galloyl- β -glucopyranoside (**2**) and a cyanogenic glucoside, 6'-*O*-galloylprunasin (**3**) were isolated from the leaves of *Syzygium austroyunnanense*. Their structures were characterized based on the spectroscopic methods and comparison with literature. This is the first phytochemical study on *Syzygium austroyunnanense*.

Keywords: *Syzygium austroyunnanense*; Myrtaceae; acylated glycoside. © 2018 ACG Publications. All rights reserved.

1. Introduction

The genus *Syzygium* (Myrtaceae) comprises about 500 species, mostly grown in the tropic regions of the world [1]. Many of them have been used as edible and medicinal plants in Southeast Asian [2]. Previous research has demonstrated that hydrolysable tannins, flavonoids, chromone derivatives, phenylpropanoids, triterpenes, and phloroglucinols derivatives are the main bioactive constituents in this genus [3–9]. *Syzygium austroyunnanense* Chang et Miao, called as “Bajiamiao” in the Dai nationality, is a fruit tree which is native to Xishuang Banna Prefecture Yunnan province and Guangxi province in China [10]. To the best of our knowledge, no phytochemical research of this species have been reported so far. In our efforts to search for bioactive constituents, the leaves of *S. austroyunnanense* were investigated and one new acylated glycoside, benzyl 3',6'-di-*O*-galloyl- β -glucopyranoside (**1**), together with two known analogues, benzyl 6'-*O*-galloyl- β -glucopyranoside (**2**) and 6'-*O*-galloylprunasin (**3**) were obtained. Here, we present the isolation and characterization of the new compound.

* Corresponding author: E- Mail: zhaoy@126.com

2. Materials and Methods

UV spectra were acquired in MeOH with a Shimadzu UV-2401PC UV-vis spectrophotometer. IR spectra were measured on a Bruker Tensor 27 FTIR Spectrometer with KBr disks. NMR spectra were recorded on a Bruker Avance III-600 and a Bruker AM-400 instruments with TMS as internal standard. ESI-MS spectra were recorded on a Waters Xevo TQ-S UPLC Triple Quadrupole Mass Spectrometer. Column chromatography was performed using silica gel (Qingdao Marine Chemical Factory, China, 200–300 mesh), Sephadex LH-20 (Pharmacia Biotech Ltd., Sweden). Thin-layer chromatography (TLC) was performed using precoated silica gel GF₂₅₄ plates (Qingdao Marine Chemical Factory). Semipreparative HPLC was performed on a Hitachi Chromaster system (Hitachi, Ltd., Japan) equipped with an YMC-Triart C₁₈ column (250 mm × 10 mm i.d., 5 μm, YMC Corporation, Japan), using a flow rate of 3.0 mL/min at a column temperature of 25 °C, and detection was performed with a DAD detector.

2.2 Plant Material

The leaves of *Syzygium austroyunnanense* Chang et Miao were collected in October 2014 from Xishuang Banna Tropical Botanical Garden, Yunnan Province, People's Republic of China, and were authenticated by Mr. Yu Chen at the State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Science. Voucher specimens (KIB 20141009) were deposited at Kunming Institute of Botany, Chinese Academy of Science.

2.3 Extraction and Isolation

The air-dried and powdered leaves of *S. austroyunnanense* (2.5 kg) were extracted with CH₃OH (6 L × 3) at room temperature. The extracts were concentrated by rotary evaporator under reduced pressure to remove organic solvent. The extract (530 g) was suspended in H₂O (0.5 L) and then successively partitioned with petroleum ether (4 × 1 L), EtOAc (4 × 1 L), and n-BuOH (4 × 1 L), sequentially. The EtOAc extract (42.0 g) was subjected to silica gel column chromatography (CC) using a gradient system of petroleum ether (PE)-Me₂CO (1:0–0:1) to afford eight fractions (Fr A–H).

Fraction F (3.5 g) was decolorized on a MCI gel (CHP 20P) CC eluted by 92%MeOH-H₂O, and then divided into three subfractions (Fr. F-1–3) by silica gel (200–300 mesh) CC eluting with CHCl₃-MeOH (8:1). Subfraction Fr. F-1 was further divided into three subfractions (Fr. F-1-1–3) by silica gel CC eluting with CHCl₃-MeOH (14:1). Fr. F-1-1 was separated by Sephadex LH-20 column (MeOH-CHCl₃, 1:1), followed by semipreparative HPLC (HITACHI HPLC system; YMC-Triart C₁₈ column, 250 × 10 mm; DAD detector, MeOH-H₂O 35:65, 210 nm, 3.0 mL/min) to give compounds **2** (3.5 mg, t_R = 17.5 min) and **3** (21.0 mg, t_R = 12.5 min). Fr. F-2 was further divided into four subfractions (Fr. F-2-1–4) by silica gel CC eluting with CHCl₃-MeOH (10:1). Compound **1** (5.0 mg, t_R = 21.0 min) was purified from Fr. F-2-4 by Sephadex LH-20 column (MeOH-CHCl₃, 1:1), and semipreparative HPLC (HITACHI HPLC system; YMC-Triart C₁₈ column, 250 × 10 mm; DAD detector, MeOH-H₂O 35:65, 210 nm, 3.0 mL/min), sequentially.

Benzyl 3',6'-di-O-galloyl-β-glucopyranoside (1): Amorphous powder; ¹H-NMR (600 MHz, CD₃OD) and ¹³C-NMR (150 MHz, CD₃OD) spectral data see Table 1; HR-ESI-MS at *m/z* 597.1216 [M + Na]⁺ (calcd for C₂₇H₂₆O₁₄, 597.1215).

Benzyl 6'-O-galloyl-β-glucopyranoside (2): Amorphous powder; ¹H-NMR (600 MHz, CD₃OD) and ¹³C-NMR (150 MHz, CD₃OD) spectral data see Table 1.

6'-O-galloylprunasin (3): Amorphous powder; ¹H-NMR (500 MHz, CD₃OD) δH: 6.70 (1H, s, H-2), 6.87 (1H, d, *J* = 8.5 Hz, H-5), 6.71 (1H, d, *J* = 8.5 Hz, H-6), 3.35 (2H, d, *J* = 8.5 Hz, H-7), 5.97 (1H, m, H-8), 5.10 (1H, m, Ha-9), 5.08 (1H, m, Hb-9), 3.90 (3H, s); ¹³C-NMR (125MHz, CD₃OD) δC: 131.9 (C-1), 111.1 (C-2), 146.4 (C-3), 144.2 (C-4), 114.2 (C-5), 121.2 (C-6), 39.9 (C-7), 137.8 (C-8), 115.5 (C-9), -OCH₃ (55.9); ESI-MS at *m/z* 421 [M - H]⁻.

Table 1. ^1H NMR and ^{13}C NMR Data for **1–2** in MeOD at 600 MHz and 150 MHz, respectively

1		1		2		2				
NO.	δ_{H}	δ_{C}	NO.	δ_{H}	δ_{C}	NO.	δ_{H}	δ_{C}		
1		140.1	1''		121.6	1		140.1	1''	121.6
2	7.35 br d (7.3)	129.5	2''	7.12 s	110.4	2	7.26 br d (7.2)	129.5	2''	7.02 s 110.3
3	7.29 t (7.3)	129.5	3''		146.6	3	7.18 t (7.2)	129.6	3''	146.7
4	7.24 t (7.3)	129.0	4''		138.8	4	7.14 t (7.2)	128.9	4''	138.9
5	7.29 t (7.3)	129.5	5''		146.6	5	7.18 t (7.2)	129.6	5''	146.7
6	7.35 br d (7.3)	129.5	6''	7.12 s	110.4	6	7.26 br d (7.2)	129.5	6''	7.02 s 110.3
7a	4.86 d (11.8)		7''		168.3	7a	4.74 d (11.8)		7''	168.5
		72.1						71.8		
7b	4.66 d (11.8)		1'''		121.9	7b	4.53 d (11.8)			
1'	4.50 d (7.9)	103.3	2'''	7.13 s	110.5	1'	4.24 d (7.7)	103.2		
2'	3.53 dd (9.5,7.9)	73.8	3'''		146.7	2'	3.27 t (9.0)	75.3		
3'	5.14 t (9.2)	79.1	4'''		139.9	3'	3.32 t (9.2)	78.1		
4'	3.68 m	70.4	5'''		146.7	4'	3.42 m	72.0		
5'	3.68 m	75.7	6'''	7.13 s	110.5	5'	3.54 m	75.7		
6'a	4.58 d (11.6)		7'''		168.4	6'a	4.45 dd (11.9,1.8)			
		64.7						64.9		
6'b	4.48 m					6'b	4.33 dd (11.9,6.0)			

3. Results and Discussion

Compound **1** was obtained as a lavender amorphous powder. Its molecular formula, $\text{C}_{27}\text{H}_{26}\text{O}_{14}$, was deduced from the HR-ESI-MS (m/z 597.1216 $[\text{M} + \text{Na}]^+$, calcd. 597.1215). Strong UV absorptions at 214 nm and 276 nm implied it had large conjugated systems. Its IR spectrum showed absorption bands for conjugated carbonyl groups at 1700 cm^{-1} and hydroxyl groups at 3397 cm^{-1} . The ^1H NMR spectrum




Figure 1. The chemical structures of compounds **1–3**.

revealed the presence of two galloyl groups (δ 7.12, 2H, s; 7.13, 2H, s), a monosubstituted benzene ring (δ 7.35, 2H, br d, $J = 7.3$ Hz; 7.29, 2H, t, $J = 7.3$ Hz; 7.24, 1H, t, $J = 7.3$ Hz), and signals diagnostic of a 4C_1 -glucopyranosyl residue. An anomeric proton signal of the glucose at δ 4.50 (d, $J = 7.9$ Hz) indicated a β -configuration for the glucosidic bond. These spectral features along with two carboxyl carbon signals at δ 168.3 and 168.4 in the ^{13}C NMR spectrum suggested that **1** is a di-*O*-galloyl- β -glucopyranoside. Additionally, a methylene group [δ_H 4.86, 4.66 (each d, $J = 11.8$ Hz); δ_C 72.1] was evident. The 1H and ^{13}C NMR spectra of **1** closely resembled those of a known compound **2**, benzyl 6'-*O*-galloyl- β -glucopyranoside [11], except that **1** had one more galloyl than **2**. The location of galloyl units at 3'-*O* and 6'-*O* of the glucopyranose moiety was established by the low-field shifts of the H-3' (δ_H 5.14) and H-6' signals (δ_H 4.58, 4.48), which was proved by the correlations of H-3'/C-7'' and H-6'/C-7'''. The location of benzyl was attached to C-1' through O atom, which was confirmed by the cross peaks from H₂-7 to C-1' and from H₂-7 to C-2 and C-6. Thus, compound **1** was assigned as benzyl 3', 6'-di-*O*-galloyl- β -glucopyranoside.




Figure 2. Key HMBC (H \rightarrow C) correlations of **1**.

The known compounds were determined to be benzyl 6'-*O*-galloyl- β -glucopyranoside (**2**) and 6'-*O*-galloylprunasin (**3**) [11] by comparing their spectroscopic data with those in the literature.

Acknowledgments

This work was financially supported by The National Natural Science Foundation of China (No. 31560098) Mid-aged and Young Academic and Technical Leader Raising Foundation of Yunnan Province (No. 2010CI040), China.

Supporting Information

Supporting information accompanies this paper on <http://www.acgpubs.org/RNP>

ORCID

Feng Li: [0000-0002-3943-5058](https://orcid.org/0000-0002-3943-5058)

Shaohua Xu: [0000-0003-3526-1705](https://orcid.org/0000-0003-3526-1705)

Yan Zhao: [0000-0002-1638-7751](https://orcid.org/0000-0002-1638-7751)

Mengjia Li: [0000-0002-7454-3767](https://orcid.org/0000-0002-7454-3767)

Wen Xu: [0000-0003-1588-8850](https://orcid.org/0000-0003-1588-8850)

Yikao Hu: [0000-0003-3913-9408](https://orcid.org/0000-0003-3913-9408)

Dingli Zhang: [0000-0002-9073-6761](https://orcid.org/0000-0002-9073-6761)

Yong Zhao : [0000-0002-3996-2480](https://orcid.org/0000-0002-3996-2480)

References

- [1] L.W. Tian, M. Xu, D. Wang, H.T. Zhu, C.R. Yang and Y.J. Zhang (2011). Phenolic constituents from the leaves of *Syzygium forrestii* Merr. and Perry, *Biochem. System. Ecol.* **39**, 156-158.
- [2] G.Q. Li, Y.B. Zhang, P. Wu, N.H. Chen, Z.N. Wu, L. Yang, R.X. Qiu, G.C. Wang and Y.L. Li (2015). New phloroglucinol derivatives from the fruit tree *Syzygium jambos* and their cytotoxic and antioxidant activities, *J. Agric. Food Chem.* **63**, 10257-10262.
- [3] T. Manaharan, D. Appleton, H.M. Cheng and U.D. Palanisamy (2012). Flavonoids isolated from *Syzygium aqueum* leaf extract as potential antihyperglycaemic agents, *Food Chem.* **132**, 1802-1807.
- [4] M.N. Samy, S. Sugimoto, K. Matsunami, H. Otsuka and M.S. Kamel (2014). One new flavonoid xyloside and one new natural triterpene rhamnoside from the leaves of *Syzygium grande*, *Phytochem. Lett.* **10**, 86-90.
- [5] M. Miyazawa and M. Hisama (2003). Antimutagenic activity of phenylpropanoids from clove (*Syzygium aromaticum*), *J. Agric. Food Chem.* **51**, 6413-6422.
- [6] F.F. Liu, T. Yuan, W. Liu, H. Ma, N.P. Seeram, Y.Y. Li, L. Xu, Y. Mu, X.S. Huang and Y.L. Li (2017). Phloroglucinol derivatives with protein tyrosine phosphatase 1B inhibitory activities from *Eugenia jambolana* seeds, *J. Nat. Prod.* **80**, 544-550.
- [7] T. Manaharan, H.M. Cheng, U.D. Palanisamy (2013). *Syzygium aqueum* leaf extract and its bioactive compounds enhances pre-adipocyte differentiation and 2-NBDG uptake in 3T3-L1 cells, *Food Chem.* **136**, 354-363.
- [8] R. Omar, L.Y. Li, T. Yuan and N.P. Seeram (2012). α -Glucosidase inhibitory hydrolysable tannins from *Eugenia jambolana* seeds, *J. Nat. Prod.* **75**, 1505-1509.
- [9] N. Yao, W. Yao, J.X. Lei, K.W. Wang (2013). Lignans from *Syzygium grijsii* and their chemotaxonomic significance, *Biochem. System. Ecol.* **46**, 79-82.
- [10] J. Chen (1984). Flora of China. Vol **53** (1). Beijing: Science Press, pp. 72.10
- [11] J.H. Isaza, H. Ito, T. Yoshida (2001). A flavonol glycoside-lignan ester and accompanying acylated glucosides from *Monochaetum multiflorum*, *Phytochemistry* **58**, 321-327.

A C G
publications

© 2018 ACG Publications