

Structural Elucidation of a Coumarin with New Skeleton from *Artemisia ordosica*

Wugedunqiqige Bao^{id}, Qinghu Wang^{id}* and Junsheng Hao^{id}

College of Traditional Mongolian Medicine, Inner Mongolia University for Nationalities,
Tongliao, Inner Mongolia 028000, P. R. China

(Received October 31, 2018; Revised January 30, 2019; Accepted February 01, 2019)

Abstract: A new coumarin, named as arteordocoumarin A (**1**), together with eight known compounds (**2-9**) were isolated from the CHCl₃ extract of *Artemisia ordosica* (*A. ordosica*). The structures of **1** was elucidated by spectroscopic methods, including UV, IR, HR-ESI-MS and extensive 1D and 2D NMR techniques.

Keywords: Arteordocoumarin A; *Artemisia ordosica*; NMR. © 2019 ACG Publications. All rights reserved.

1. Introduction

A. ordosica, Asteraceae family, is one of the main arido-active shrubs growing in the arid and semi-arid areas of the north China including Inner Mongolia, Ningxia, Gansu and Shanxi [1, 2]. The aerial parts of *A. ordosica* is utilized as a folk medicine for expelling rheumatism, clearing heat, and dispelling swelling [3]. Sterols [4], coumarins [5], terpenoids [6], flavonoids [7, 8] and acetylenes [3] were isolated previously from this plant. However, the secondary metabolites from *Artemisia ordosica* often differ when grown in different ecological environments. In order to continue our research on the bioactive secondary metabolites from *Artemisia ordosica* collected in Tongliao of Inner Mongolia, China, we now describe the isolation and structure elucidation of a new coumarin compound, together with eight known ones.

2. Materials and Methods

2.1. Instrumentation and Reagents

A Shimadzu UV-2201 spectrometer (Shimadzu, Japan) was used to record the he UV spectra. The IR spectra were recorded in KBr discs on a Thermo Nicolet 200 double beam spectrophotometer (Shimadzu, Japan). A Waters Xevo G2-S QT (Waters, USA) was used to measure the HR-ESI-MS spectra. NMR spectra were measured on a Bruker AV-500 spectrometer (Bruker, Germany) with tetramethylsilane (TMS) as the internal reference, and chemical shifts are expressed in δ (ppm). Column

* Corresponding author: E-Mail: wqh196812@163.com; Phone:086-0475-8314242 Fax:086-0475-8314242

chromatography was performed by using silica gel (200-300 mesh, Marine Chemical Factory, Qingdao, China).

2.2. Plant Materials

The aerial parts of *A. ordosica* were collected in Tongliao, Inner Mongolia of China, in June 2017, and identified by Prof. Buhebateer (Inner Mongolia University for Nationalities). A voucher (ref. no. 20170612) has been deposited at the Herbarium of college of Traditional Mongolian Medicine of Inner Mongolia University for Nationalities.

2.3. Extraction

Ground-dried aerial parts of *A. ordosica* (2.0 kg) were extracted with CHCl₃ (25 L) under reflux after extraction with 10 L petroleum ether. Evaporation of the solvent under reduced pressure delivered the CHCl₃ extract (150.0 g). The CHCl₃ extract was fractionated by column chromatography on silica gel and gradiently eluted with petroleum ether-CH₃COCH₃ (60:1 to 20:1) to give 3 fractions (Fr. 1-3). Fr. 1 (320 mg) was separated by TLC (cyclohexane-ethyl acetate, 10:1) yielding **1** (17 mg), **2** (11 mg) and **3** (15 mg); Fr. 2 (400.0 mg) was further eluted on a Sephadex LH-20 column with MeOH:CHCl₃ (v:v, 1:1) and then separated by TLC (cyclohexane-ethyl acetate, 7:1) yielding **4** (13 mg), **5** (9 mg), **6** (15 mg) and **7** (18 mg). Fr. 3 (330 mg) was separated by TLC (cyclohexane-ethyl acetate, 3:1) yielding **8** (16 mg) and **9** (21 mg).

Arteordocoumarin A (1): White needle; ¹H-NMR (500MHz, DMSO-d₆) and ¹³C-NMR (125MHz, DMSO-d₆) spectral data see Table 1; HR-ESI-MS at *m/z* 209.0467 [M-H]⁻ (calcd for C₁₀H₉O₅, 209.0450).

6,7-dimethoxycoumarin (2) [9]: White needle; ¹H-NMR (500MHz, CDCl₃) δ_H: 6.31 (1H, d, *J* = 9.5 Hz, H-3), 7.65 (1H, d, *J* = 9.5 Hz, H-4), 6.86 (1H, s, H-5), 6.88 (1H, s, H-8), 3.94 (3H, s, 6-OCH₃), 3.97 (3H, s, 7-OCH₃); ¹³C-NMR (125MHz, CDCl₃) δ_C: 161.5 (C-2), 113.6 (C-3), 143.3 (C-4), 107.9 (C-5), 146.3 (C-6), 152.8 (C-7), 100.0 (C-8), 150.0 (C-9), 111.4 (C-10), 56.4 (6-OCH₃), 56.3 (7-OCH₃).

6,7,8-trimethoxycoumarin (3) [10]: White needle; ¹H-NMR (500MHz, CDCl₃) δ_H: 6.06 (1H, d, *J* = 9.0 Hz, H-3), 7.52 (1H, d, *J* = 9.0 Hz, H-4), 7.30 (1H, s, H-5), 3.88 (3H, s, 6-OCH₃), 3.86 (3H, s, 8-OCH₃), 3.82 (3H, s, 7-OCH₃); ¹³C-NMR (125MHz, CDCl₃) δ_C: 165.4 (C-2), 115.5 (C-3), 140.3 (C-4), 114.9 (C-5), 145.0 (C-6), 125.1 (C-7), 133.5 (C-8), 134.4 (C-9), 120.7 (C-10), 56.4 (6-OCH₃), 56.1 (8-OCH₃), 55.8 (7-OCH₃).

6-hydroxy-7-methoxycoumarin (4) [9]: White needle; ¹H-NMR (500MHz, CDCl₃) δ_H: 6.31 (1H, d, *J* = 9.5 Hz, H-3), 7.64 (1H, d, *J* = 9.5 Hz, H-4), 6.93 (1H, s, H-5), 6.85 (1H, s, H-8), 3.91 (3H, s, 7-OCH₃); ¹³C-NMR (125MHz, CDCl₃) δ_C: 161.6 (C-2), 113.5 (C-3), 143.5 (C-4), 108.3 (C-5), 147.3 (C-6), 152.0 (C-7), 103.0 (C-8), 153.0 (C-9), 113.4 (C-10), 56.5 (7-OCH₃).

4-hydroxyacetophenone (5) [11]: White needle; ¹H-NMR (500MHz, CDCl₃) δ_H: 7.93 (2H, d, *J* = 8.5 Hz, H-2,6), 6.91 (2H, d, *J* = 8.5 Hz, H-3,5), 2.58 (3H, s, -CH₃); ¹³C-NMR (125MHz, CDCl₃) δ_C: 130.4 (C-1), 131.0 (C-2), 115.3 (C-3), 160.1 (C-4), 115.3 (C-5), 131.0 (C-6), 197.9 (C=O), 26.3 (-CH₃).

4-hydroxy-5-methoxyacetophenone (6) [11]: White needle; ¹H-NMR (500MHz, CDCl₃) δ_H: 7.59 (1H, d, *J* = 2.0 Hz, H-2), 6.98 (1H, d, *J* = 8.0 Hz, H-5), 7.62 (1H, d, *J* = 8.0, 2.0 Hz, H-6), 3.90 (3H, s, -OCH₃), 2.51 (3H, s, -CH₃); ¹³C-NMR (125MHz, CDCl₃) δ_C: 129.9 (C-1), 111.1 (C-2), 147.0 (C-3), 146.8 (C-4), 114.4 (C-5), 126.9 (C-6), 190.3 (C=O), 56.8 (-OCH₃), 26.3 (-CH₃).

4-hydroxybenzaldehyde (7) [12]: White needle; ¹H-NMR (500MHz, CDCl₃) δ_H: 7.82 (2H, d, *J* = 8.5 Hz, H-2,6), 7.02 (2H, d, *J* = 8.5 Hz, H-3,5), 9.85 (1H, s, -CHO); ¹³C-NMR (125MHz, CDCl₃) δ_C: 139.3 (C-1), 132.5 (C-2), 116.0 (C-3), 162.0 (C-4), 116.0 (C-5), 132.5 (C-6), 191.3 (-CHO).

4-hydroxy-5-methoxybenzaldehyde (**8**) [13]: White needle; $^1\text{H-NMR}$ (500MHz, CDCl_3) δ_{H} : 7.43 (1H, brs, H-2), 7.05 (1H, d, $J = 8.5$ Hz, H-5), 7.45 (1H, brd, $J = 8.5$ Hz, H-6), 9.83 (1H, s, -CHO), 3.99 (3H, s, $-\text{OCH}_3$); $^{13}\text{C-NMR}$ (125MHz, CDCl_3) δ_{C} : 129.9 (C-1), 108.8 (C-2), 151.7 (C-3), 147.2 (C-4), 114.4 (C-5), 127.6 (C-6), 191.0 (C=O), 56.1 ($-\text{OCH}_3$).

4,5-dihydroxybenzaldehyde (**9**) [13]: $^1\text{H-NMR}$ (500MHz, CDCl_3) δ_{H} : 7.44 (1H, d, $J = 2.0$ Hz, H-2), 6.96 (1H, d, $J = 8.0$ Hz, H-5), 7.46 (1H, brd, $J = 8.0, 2.0$ Hz, H-6), 9.85 (1H, s, -CHO); $^{13}\text{C-NMR}$ (125MHz, CDCl_3) δ_{C} : 129.8 (C-1), 108.9 (C-2), 146.8 (C-3), 146.2 (C-4), 114.3 (C-5), 127.7 (C-6), 191.0 (C=O).

3. Results and Discussion

From the CDCl_3 of *A. ordosica*, six compounds were obtained using chromatographic methods (CC and TLC). On the basis of ^1H , ^{13}C NMR, COSY, HSQC, HMBC and HR-ESI-MS spectra, and modified Mosher's method as well as by comparison with previous reports [9-13], compound **1** was identified as a coumarin with new skeleton while the remaining eight compounds were found to be the known compounds, 6,7-dimethoxycoumarin (**2**), 6,7,8-trimethoxycoumarin (**3**), 6-hydroxy-7-methoxycoumarin (**4**), 4-hydroxyacetophenone (**5**), 4-hydroxy-5-methoxyacetophenone (**6**), 4-hydroxybenzaldehyde (**7**), 4-hydroxy-5-methoxybenzaldehyde (**8**) and 4,5-dihydroxybenzaldehyde (**9**) (Figure 1).




Figure 1. Structures of compounds **1-9**

Compound **1** was obtained as a white needle, mp 137-139 °C; IR (KBr) ν_{max} (cm^{-1}): 3312, 1683, 1643 and 1320 cm^{-1} . The molecular formula was determined to be $\text{C}_{10}\text{H}_{10}\text{O}_5$ by HR-ESI-MS exhibiting a pseudomolecular ion peak at m/z 209.0467 [M-H^-] (calcd for $\text{C}_{10}\text{H}_9\text{O}_5$, 209.0450). In the ^1H NMR spectrum (Table 1), two characteristic resonances for H-3 and H-4 of a coumarin at δ_{H} 6.03 (1H, d, $J = 10.0$ Hz, H-3) and 8.06 (1H, d, $J = 10.0$ Hz, H-4). In addition, the signals at δ_{H} 3.64 (3H, s) and 3.82 (3H, s) indicated the presence of two methoxy groups. The remaining signal at δ_{H} 6.39 (1H, s) was assigned to H-9 compared with the data of H-25 (δ_{H} 6.21) in fasciospongides **A** [14], which was confirmed by the HMBC correlations (Figure 2) from δ_{H} 6.39 (1H, s, H-9) to C-5 (δ_{C} 104.8), C-6 (δ_{C} 152.3) and C-7 (δ_{C} 133.5).





Figure 2. Selected HMBC correlations for **1**

Table 1. ^1H (500 MHz) and ^{13}C -NMR (125MHz) data of compound **1** in $\text{DMSO-}d_6$

Position	δ_{H} (ppm), J (Hz)	δ_{C} (ppm)
2	—	161.5
3	6.03 d (10.0)	109.1
4	8.09 d (10.0)	141.1
5	—	104.8
6	—	152.3
7	—	133.5
8	—	157.0
9	6.39 s	89.7
7-OCH ₃	3.64 s	60.6
8-OCH ₃	3.82 s	56.5

The ^{13}C NMR spectrum of **1** showed 10 carbon signals, of which 8 were assigned to the coumarin skeleton part and 2 to the two methoxy groups. In coumarin skeleton part, there were only 8 carbon signals (δ_{C} 161.5, 109.1, 141.8, 104.8, 152.3, 133.5, 157.0, 89.7), which was different from the 9 carbon signals of an usual coumarin skeleton [9, 10]. The benzene in an usual coumarin skeleton was substituted by the 1,3-diene cyclopentane in the new coumarin skeleton, in which the HMBC correlations from H-4 to C-2 (δ_{C} 161.5), C-6 (δ_{C} 152.3) and C-9 (δ_{C} 89.7), and H-3 to C-2 (δ_{C} 161.5) and C-5 (δ_{C} 104.8), H-9 to C-5 (δ_{C} 104.8), C-6 (δ_{C} 152.3) and C-7 (δ_{C} 133.5) were show. In addition, the HMBC correlations δ_{H} 3.64 (-OCH₃) to δ_{C} 133.5 (C-7) and 3.82 (-OCH₃) to δ_{C} 157.0 (C-8) revealed that the two methoxy groups were linked to the C-7 and C-8, respectively. The modified Mosher's method was used to produce (*R*)- and (*S*)-MTPA esters (**1a**, **1b**), and signals corresponding to H-3, H-4 and 8-OCH₃ were relatively deshielded in **1a** compared to **1b**, indicating that the absolute configuration of C-9 is *S* (Figure 3). Thus, the structure of compound **1** was elucidated and named as arteordocoumarin A.

**Figure 3.** Results with the modified Mosher's method ($\Delta = \delta_{\text{S}} - \delta_{\text{R}}$) for compound **1**

Acknowledgments

This work was supported by the Science and Technology Major Project of Inner Mongolia Mongolian medicine in China (No. GCY20150712) and Graduate Research Innovation Projects of Inner Mongolia (No. B20161013601).

Supporting Information

Supporting information accompanies this paper on <http://www.acgpubs.org/journal/records-of-natural-products>

ORCID

Wugedunqiqige Bao: [0000-0003-0744-8361](https://orcid.org/0000-0003-0744-8361)

Qinghu Wang: [0000-0002-6272-6398](https://orcid.org/0000-0002-6272-6398)

Junsheng Hao: [0000-0002-7733-133X](https://orcid.org/0000-0002-7733-133X)

References

- [1] M.J. Abad, L.M. Bedoya, L. Apaza and P. Bermejo (2012). The *Artemisia* L. genus: A review of bioactive essential oils, *Molecules* **17**, 2542-2566.
- [2] H. Zhang, D. Zhou, Y. Luo, J. Wang and S. Zong (2013). Identification of volatile compounds emitted by *Artemisia ordosica* (Artemisia, Asteraceae) and changes due to mechanical damage and weevil infestation, *Z. Naturforsch. C.* **68**, 313-317.
- [3] Z. Zhang, S.S. Guo, W.J. Zhang, Z.F. Geng, J.Y. Liang, S.S. Du, C.F. Wang and Z.W. Deng (2017). Essential oil and polyacetylenes from *Artemisia ordosica* and their bioactivities against *Tribolium castaneum* Herbst (Coleoptera:Tenebrionidae), *Ind. Crop. Prod.* **100**, 132-137.
- [4] D.B. Zhao, Y.F. Jiang, C.J. Xi, X.H. Liu and H.Q. Wang (2005). Isolation and identification of capillarin from *Artemisia ordosica* Krasch, *Chin. J. Struct. Chem.* **24**, 637-640
- [5] R.X. Tan, W.F. Zheng and H.Q. Tang (1998). Biologically active substances from the genus *Artemisia*, *Planta. Med.* **64**, 295-302
- [6] Y. Zhong, X.S. Feng and Y.R. Liu (2016). Chemical constituents from *Artemisia ordosica* Krasch and their bioactivities, *Chem. Bioeng.* **33**, 36-38.
- [7] D.B. Zhao, Y.X. Yang, W. Zhang, X.H. Liu and H.Q. Wang (2005). Studies on flavonoid constituents from herbs of *Artemisia ordosica* II, *Chin. J. Chin. Mater. Med.* **31**, 1959-1961.
- [8] W. Zhang, D.B. Zhao, M.J. Li, X.H. Liu and H.Q. Wang (2006). Studies on flavonoid constituents from herbs of *Artemisia ordosica* II, *Chin. J. Chin. Mater. Med.* **30**, 1430-1432.
- [9] J.Y. Wu, Y.B. Wu, J. Yi, T. Wang, Z.H. Huang, J.G. Wu and J.Z. Wu (2013). Chemical constituents in leaves of *Magnolia officinalis* var. *biloba*, *Chin. Trad. Herb. Drag.* **44**, 2965-2968.
- [10] C. Kofinas, I. Chinou, A. Loukis, C. Harvala and C. Roussakis (1998). Cytotoxic coumarins from the aerial parts of *Tordylium apulum* and their effects on a non-small-cell bronchial carcinoma line, *Planta. Med.* **64**, 171-179.
- [11] L.L. Song, X.W. Li and F.F. Yan (2010). study on the chemical constituents of *Daphne genkwa*, *Chin. Trad. Herb. Drag.* **41**, 536-539.
- [12] Y.W. Wang, Z.F. Li, M.Z. Zhen, Y.L. Feng, Q. Wang, X. Li and S.L. Yang (2013). Chemical constituents of *Gastrodia elata*, *Chin. Trad. Herb. Drag.* **44**, 2794-2796.
- [13] X.H. Duan, Z.L. Li, D.S. Yang, F.L. Zhang, Q. Lin and R. Dai (2013). Study on the chemical constituents of *Gastrodia elata*, *J. Chin. Med. Mat.* **36**, 1608-1611.
- [14] A. Montagnac, M. Pas and C. Debitus (1994). Fasciospongides A, B and C, new manoalide derivatives from the sponge *fasciospongia* SP, *J. Nat. Prod.* **57**, 186-190.

A C G
publications

© 2019 ACG Publications