

Essential Oil Composition of *Tilia platyphyllos* Scop. Collected from Different Regions of Kosovo

Nita Kelmendi¹, Behxhet Mustafa², Fitore Zahiri²
Dashnor Nebija^{3*} and Avni Hajdari²

¹ Department of Pharmacy, College of Medical Sciences “Rezonanca” Glloku te Shelgjet 10000-Prishtina, Kosovo

² Department of Biology, Faculty of Mathematical and Natural Science, University of Prishtina “Hasan Prishtina”. Mother Theresa St. 10000 Prishtina, Kosovo

³ Department of Pharmacy, Faculty of Medicine, University of Prishtina. “Hasan Prishtina” Mother Theresa St. 10000 Prishtina, Kosovo

(Received February 13, 2020; Revised March 02, 2020; Accepted March 06, 2020)

Abstract: The aim of this study was to assess variation of the chemical composition of essential oils obtained from inflorescences and leaves of *Tilia platyphyllos* Scop. collected from five different localities in Kosovo. GC-MS/GC-FID analysis revealed the presence of 96 compounds in essential oils obtained from inflorescences of large-leaved lime with the most prominent classes including hydrocarbons, with the respective concentrations in flowers and leaves of 41.7% - 46.2% and 27.5% - 36.8%, respectively, oxygenated sesquiterpenes (12.2%-23.0% and 21.3% -31.2%, in flowers and leaves, respectively) and fatty acids and their derivatives (8.0% - 19.6% and 18.1% – 23.1% in flowers and leaves, respectively) whereas monoterpenes were present in smaller amount. Experimental data showed that large-leaved lime is reach in different groups of volatile compounds, and this species could be further recommended for food and medicinal application.

Keywords: Large-leaved lime; essential oil; principal component analysis, © 2020 ACG Publications. All rights reserved.

1. Plant Source

Plant material (inflorescences) of the large-leaved lime (*Tilia platyphyllos*, Scop. Malvaceae), were collected from five different localities in Kosovo (Pejë, Gjiilan, Lipjan, Klinë and Suharekë regions) during the period June-July 2016. The plant was identified by Professor asoc. Avni Hajdari, Department of Biology, University of Prishtina. A voucher specimen (Herbarium Number: LEB/2016/1) has been deposited at the Department of Biology University of Prishtina, Kosovo.

2. Previous Studies

Tilia platyphyllos Scop., Malvaceae, known as large-leaved lime or linden, is large-sized deciduous tree, native through Europe. This large-leaved tree prefers warmer climates and can reach 30-40m in height, with a trunk up to around 1m in diameter [1]. Asymmetrical, heart shaped leaves are about 9cm long and have pointed tips with upper surface of dark green colour and paler underside. The cream to yellow flowers, are fragrant, have five petals and hang in tiny clusters of 4 to 10 [2]. Large-leaved lime has industrial cosmetics and medicinal importance [3, 4]. In Kosovo large leaved lime was traditionally used for different purposes, including tea preparation (recreational use/ panacea),

* Corresponding author: E-Mail: dashnor.nebija@uni-pr.edu Phone: + 383-45-401585 Fax: +383(0)38-512-223

treatment of sore throat and lung inflammations, abdominal pain and respiratory problems (cough, fever) [5-11]. It has been shown that the principal bioactive compounds of *Tilia platyphyllos* Scop. are mucilaginous substances, flavonoids and essential oils. [4, 11-13]. In a comparative metabolomics study of *Tilia platyphyllos* Scop. bracts totally 504 compounds were detected, including flavonoid glycosides, catechins, procyanidins, quinic acid derivatives and coumarins [14]. Chemical composition of essential oils obtained from linden inflorescences was studied in samples obtained from different region [15-19].

GC-FID and GC-MS analyses: GC analyses were performed using of an Agilent 7890A gas chromatography system equipped with flame ionization (FID) detector (Agilent Technologies) using HP-5MS column 30 m x 0.25 mm with 0.25 μm film thickness. Helium was used as the carrier gas with an initial flow rate of 0.6 mL/min and subsequently at a constant pressure of 16.6 psi. GC/MS analyses were carried out using an Agilent 7800A GC System coupled with a 575C MSD. The ionisation energy was 70 eV with a mass range of 40-400 m/z. The separation was performed with the same column and temperature program the same as for the analytical GC. Identification of the essential oils components was completed by comparing their Kovats retention indices with those reported in literature [20]. The components were also identified by comparing the mass spectra of each constituent with those stored in the MS library search (NIST 08.L and WILEY MS 9th) and with mass spectra from the literature [20].

Principal component analysis (PCA) was used to evaluate whether the identified essential oil components can be useful for reflecting the chemotaxonomy of *Tilia platyphyllos*. Constituents with concentrations higher than 3% (bold in Table 1) of were subjected to statistical analysis.




Figure 1. Graph obtained from the Principal Component Analysis of essential oils obtained from inflorescences of *Tilia platyphyllos* from 5 different localities. Inf. - Inflorescences; Leav. - leaves

Table 1 The chemical composition of the essential oils obtained from the inflorescences (IF) and leaves (L) of *Tilia platyphyllos*

KI ^a	Compound ^b	Pejë		Gjilan		Lipjan		Klinë		Suharekë	
		IF	L	IF	L	IF	L	IF	L	IF	L
838	4-methylpentanol	-	-	1.42	1.00	1.63	2.50	1.02	0.65	-	0.20
846	<i>E</i> -2-Hexenal	1.15	0.64	0.83	0.47	0.98	0.92	0.74	0.75	0.58	0.37
862	<i>E</i> -2-Hexenol	-	0.05	0.44	0.29	-	0.07	0.03	0.20	0.85	0.46
900	<i>n</i> -Nonane	0.16	0.03	0.19	0.04	0.26	0.14	0.21	0.37	0.17	0.14
902	Heptanal	1.07	0.28	0.80	0.28	0.72	0.51	1.47	0.73	0.84	0.42
960	Benzaldehyde	0.19	0.20	0.06	0.24	0.13	0.21	0.14	0.30	-	0.25
973	Hexanoic acid	0.38	0.24	0.40	0.21	0.49	0.26	0.39	0.36	0.71	0.26
987	<i>E</i> -3-Hexenoic acid	0.18	0.65	0.25	0.96	0.32	1.75	0.41	1.01	0.38	1.41
989	1-Decene	0.41	0.52	0.54	0.64	0.51	1.43	0.66	1.51	0.72	2.36
988	2-Pentylfuran	0.62	0.39	0.50	0.25	0.51	0.29	0.43	0.57	0.48	0.58
1000	<i>n</i> -Decane	0.16	0.16	0.11	0.15	0.24	0.22	0.21	0.39	-	0.20
1009	Hexyl acetate	0.77	0.56	0.52	0.56	0.63	0.69	0.50	0.70	0.42	0.64
1035	<i>E</i> -3-Octen-2-one	-	-	0.46	0.43	0.36	0.52	0.35	0.70	0.20	0.35
1042	Benzeneacetaldehyde	1.80	0.25	1.23	0.37	1.22	0.58	1.68	0.80	1.28	0.56
1068	<i>n</i> -octanol	0.65	0.81	0.91	1.26	0.68	1.41	0.69	1.21	0.49	1.00
1097	2-Nonanol	0.34	0.87	0.40	0.66	0.50	1.90	0.41	0.72	-	0.33
1097	Linalool	1.09	1.35	0.50	0.84	1.40	0.96	1.11	1.71	0.95	2.54
1100	<i>n</i>-Nonanal	8.97	4.10	3.30	2.76	8.27	6.15	8.35	6.96	5.40	2.78
1161	2<i>E</i>-Nonen-1-ol	0.62	0.23	4.19	2.22	0.21	0.36	0.68	0.36	-	0.38
1166	Coahuilensol	0.32	0.15	0.30	0.21	0.18	0.27	0.11	0.24	-	0.29
1169	Borneol	0.32	0.11	0.42	0.14	0.55	0.23	0.16	0.05	0.30	0.21
1175	<i>cis</i> -Pinocamphone	0.22	0.05	0.11	0.02	0.16	0.09	0.07	0.12	0.19	0.24
1192	2-Decanone	0.68	0.26	0.71	0.31	0.52	0.24	0.40	0.41	0.42	0.33
1191	Methyl salicylate	0.72	0.23	0.62	0.24	0.54	0.22	0.33	0.34	0.27	0.20
1196	Safranal	-	0.28	-	0.48	-	0.45	0.08	0.44	-	0.74
1201	<i>n</i>-Decanal	0.81	0.48	0.44	0.53	0.68	0.65	0.97	0.90	3.00	-
1219	β -Cyclocitral	0.46	0.28	0.49	0.28	0.23	0.37	0.26	0.25	-	0.35
1235	3 <i>Z</i> -Hexenyl 3-methylbutanoate	0.20	0.13	0.72	0.23	0.63	0.23	0.26	0.27	-	0.32
1258	Carvenone	1.61	0.05	1.40	0.33	1.38	0.28	0.72	0.39	0.81	0.36
1270	Nonanoic acid	0.52	1.09	0.52	0.39	0.54	0.45	0.46	0.61	0.61	0.78
1305	Undecanal	0.66	0.38	0.73	0.89	0.78	0.94	0.50	0.57	-	0.25
1311	2-Adamantanone	-	0.17	0.37	0.06	0.07	0.13	-	-	-	0.22
1324	Methyl geranate	2.48	1.21	2.38	1.70	1.71	1.10	1.55	1.50	1.58	0.89
1329	<i>Z</i> -Hasmigone	0.27	0.84	0.15	1.08	0.22	1.36	0.42	0.91	0.32	1.19
1349	α -Terpinyl acetate	-	0.08	0.17	0.26	0.03	0.10	0.18	0.79	0.61	1.08
1359	Eugenol	0.23	0.60	0.17	0.09	0.18	-	0.18	0.15	-	0.31
1366	<i>n</i> -decanoic acid	0.24	-	0.19	-	0.16	-	0.14	0.11	0.42	0.26
1380	<i>Z</i> -3-Hexenyl hexanoate	0.39	0.17	0.21	0.09	0.24	0.11	0.27	0.54	0.43	0.79
1384	<i>E</i> - β -damascenone	-	0.18	0.15	0.21	-	0.22	0.05	0.18	-	0.33
1391	<i>E</i> -Jasmone	0.18	0.49	0.30	0.56	0.08	0.11	0.05	0.14	-	0.15
1392	<i>Z</i>-Jasmone	3.06	1.60	0.65	0.94	2.29	1.81	2.11	1.07	0.77	1.81
1402	α -Funebrene	0.42	0.07	0.57	0.09	0.12	0.16	0.27	1.27	-	0.26

Table 1 continued..

1403	Methyl eugenol	0.25	0.66	0.02	0.01	0.04	0.35	0.12	0.11	-	0.47
1419	<i>E</i> -Caryophyllene	0.18	0.78	0.14	0.74	-	1.11	0.09	0.72	1.07	4.20
1422	β -Duprezianene	0.51	0.34	1.08	0.35	0.43	0.35	0.42	0.54	0.42	0.25
1430	<i>E</i> - α -Ionone	0.32	0.09	0.15	0.15	0.38	0.06	0.25	0.32	-	0.21
1441	Aromadendrene	0.15	0.02	0.06	0.32	0.13	0.53	-	0.07	-	-
1446	Seychellene	-	0.41	-	0.35	-	1.47	0.09	0.50	-	1.04
1454	α -Humulene	-	0.98	0.34	0.94	0.71	0.08	0.44	0.11	0.35	0.22
1466	2 <i>E</i> -Dodecenal	0.26	0.15	0.08	0.03	0.13	0.10	0.27	0.38	-	-
1476	α -Neocallitropsene	1.11	0.66	0.34	0.51	0.63	0.20	0.71	0.32	0.88	0.91
1488	<i>E</i> - β -Ionone	-	0.15	0.07	0.35	0.04	0.24	0.03	0.10	-	0.49
1493	β -etispirene	0.18	1.30	-	1.37	-	1.66	0.20	0.78	0.39	0.86
1500	Isodaucene	-	0.25	0.19	0.23	-	0.32	0.05	0.20	-	0.42
1502	<i>trans</i>- β-Guaiene	1.30	0.25	1.71	0.70	2.72	0.48	2.64	0.78	-	1.27
1509	α-Bulnesene	-	-	4.64	0.38	-	-	-	-	-	-
1512	δ -Z-Macrocarpene	0.31	0.33	0.35	0.24	0.54	0.56	0.39	0.51	0.33	0.27
1514	α -Cadinene	-	0.39	-	0.42	-	0.45	0.05	0.17	-	0.88
1520	1-endo-Bourbonanol	-	0.37	0.01	0.31	-	0.35	0.07	0.26	0.31	0.52
1523	Artedouglasia oxide C	0.44	1.62	0.56	1.27	0.35	1.50	0.62	1.55	0.37	2.59
1536	Liguloxide	-	1.49	0.07	1.00	-	1.82	0.11	0.54	-	0.90
1545	α -Calacorene	0.16	0.77	0.19	2.16	-	1.63	0.24	0.80	0.26	0.43
1563	<i>epi</i>-Longipinanole	1.26	4.05	0.82	2.08	1.74	6.96	2.33	5.42	2.02	7.82
1567	3<i>Z</i> -<i>epi</i> Hexyl benzoate	2.49	0.70	2.84	0.55	3.59	0.48	1.15	0.81	2.28	0.46
1580	<i>n</i> -Hexyl benzoate	-	0.16	-	0.08	-	0.28	0.07	0.37	-	0.14
1587	2 <i>E</i> -Hexenyl benzoate	-	0.35	-	0.31	-	0.09	0.34	0.50	0.32	0.69
	5-oxy-Isobornyl										
1603	isobutanoate	-	0.33	-	0.10	-	0.08	0.05	0.26	0.78	0.48
1612	Tetradecanal	0.45	3.34	0.60	3.72	0.58	4.12	1.34	3.63	0.70	4.67
1646	α-Muurolol	0.18	2.58	0.12	1.87	-	2.15	0.52	1.14	0.23	0.92
1654	α -Cadinol	-	0.20	0.24	0.19	0.30	0.13	0.48	0.21	0.26	-
1668	6 <i>Z</i> -Coniferyl alcohol	0.36	0.45	0.29	0.58	0.13	0.41	0.19	0.58	0.35	0.59
1673	5 iso-Cedranol	-	0.56	0.14	0.82	-	0.37	-	-	-	-
1676	Cadalene	-	0.36	-	0.17	-	0.11	0.27	0.07	-	0.26
1700	<i>n</i> -Heptadecane	0.25	0.53	0.09	0.49	0.13	0.90	0.60	1.12	0.71	2.36
1700	Amorpha-4,9-dien-2-ol	0.76	2.76	0.66	2.95	0.88	3.94	0.95	2.23	-	3.49
1713	Cedroxyde	-	0.39	0.02	0.65	-	0.56	0.05	0.22	-	0.40
	<i>E</i> -Sesquilandulyl										
1740	acetate	0.08	0.39	-	0.19	-	0.21	0.04	0.10	0.15	0.27
1768	Cedryl acetate	-	1.01	0.18	1.10	0.48	0.81	0.38	0.53	-	0.95
1768	14-oxy-α Muurolene	1.76	1.09	1.34	2.15	0.86	2.50	1.26	1.11	3.35	1.01
1788	8-Cedren-13-ol, acetate	0.87	0.44	1.12	1.58	0.65	0.72	1.61	0.84	0.47	0.22
	Isopropyl										
1829	tetradecanoate	0.61	0.64	0.73	0.60	0.97	1.15	0.96	1.66	0.64	0.41
	2,7(14),10-										
1845	isabolatriene-1-ol-4-one	0.26	1.64	0.04	0.82	-	0.53	0.04	0.36	0.47	0.70
1847	Phenylethyloctanoate	0.21	0.39	0.21	0.27	0.39	0.27	0.35	0.87	0.67	0.32

Table 1 continued..

1875	<i>n</i> -Hexadecanol	1.32	2.18	1.37	0.81	0.49	0.62	0.90	1.70	1.11	0.66
1900	<i>n</i> -Nanodecane	2.79	1.76	1.22	1.45	2.28	1.01	1.41	0.56	0.26	0.23
1913	5E,9E-Farnesyl acetone	2.42	1.47	3.79	1.07	3.45	0.55	2.46	0.74	11.98	0.56
2100	<i>n</i> -Heneicosane	13.50	4.07	3.59	2.10	12.38	2.16	4.92	5.94	9.88	3.91
2300	<i>n</i> -Tricosane	7.25	3.16	5.48	1.73	5.92	0.46	21.76	2.89	5.20	0.93
2400	<i>n</i> -Tetracosane	0.85	1.94	1.10	2.37	1.23	2.75	1.16	2.83	2.84	5.16
2500	<i>n</i> -Pentacosane	13.90	10.88	19.61	13.58	14.65	8.91	10.75	9.85	1.06	0.49
2600	<i>n</i> -Hexacosane	0.42	3.71	1.65	2.64	0.41	3.56	0.74	3.62	2.17	6.84
2700	<i>n</i> -Heptacosane	1.14	2.40	1.48	4.21	1.70	0.96	1.00	1.10	2.72	1.81
2725	Unidentified 1	5.87	5.67	5.49	4.96	5.88	2.70	3.48	3.63	-	4.83
2745	Unidentified 2	0.24	1.51	0.35	2.00	0.28	0.92	0.18	0.32	1.33	0.49
2800	<i>n</i> -Octacosane	1.74	2.74	2.96	2.26	1.36	1.55	0.98	1.38	6.47	1.77
2900	<i>n</i> -Nanocosane	1.49	4.91	3.66	4.96	2.80	3.43	1.54	3.37	13.98	3.69
Total (%)		99.9	100	100.0	100	99.9	100	100	99.9	99.9	100

^aKovats indices calculated against a C9- C28n-alkanes mixture on the HP5 MS column. ^bCompounds are listed in order of elution from a HP-5MS column. The percentage for each population represents the mean values of n calculated samples (n=3-5 samples). Boldface marked compounds (with concentrations higher than 3%) were chosen for PCA statistical analyses. ” –“ < 0.1%. inflorescences (IF) and leaves (L)

In this study, ninety-six compounds were separated (Table 1) from the essential oils obtained from inflorescences of *Tilia platyphyllos*, from which ninety-four were identified, while two compounds were not identified. (Supporting information, Figures S1 and S2). The most abundant classes of compounds were hydrocarbons with the respective concentrations in inflorescences and leaves of: 41.68 - 46.18% and 27.48 - 36.81%; followed by oxygenated sesquiterpenes (12.19 - 23.05% and 21.29 - 31.19% in inflorescences and leaves, respectively), fatty acids and their derivatives (8.03 - 19.61% and 18.05 - 23.1% in inflorescences and leaves, respectively); sesquiterpenes (3.44 - 9.42% and 6.04 - 10.84% in inflorescences and leaves, respectively) and oxygenated monoterpenes (4.23% - 6.18 % and 3.88 - 7.22% in inflorescences and leaves, respectively) (Table 1). The results of our study were in accordance with studies performed elsewhere [16, 17, 19]. In this work the total phenolic and flavonoid contents and antioxidant activity of methanol extract of *Tilia platyphyllos* was studied as well (Supporting information, Table 1 and 2).

The two-dimensional axis system of the PCA (Figure 1) shows the presence of two principal groups of compounds clustered based on the plant organs from which volatile compounds have been extracted, while the origin of plant material played a smaller role in the variability of the chemical composition of essential oils.

Supporting Information

Supporting Information accompanies this paper on <http://www.acgpubs.org/journal/records-of-natural-products>

ORCID 

Nita Kelmendi: [0000-0002-0346-3356](https://orcid.org/0000-0002-0346-3356)

Behxhet Mustafa: [0000-0002-1052-5252](https://orcid.org/0000-0002-1052-5252)

Fitore Zahiri: [0000-0003-4867-1906](https://orcid.org/0000-0003-4867-1906)

Avni Hajdari: [0000-0001-5688-9679](https://orcid.org/0000-0001-5688-9679)

Dashnor Nebija: [0000-0002-8853-794X](https://orcid.org/0000-0002-8853-794X)

References

- [1] E. Eaton, G. Caudullo and D. De Rigo (2016). *Tilia cordata*, *Tilia platyphyllos* and other limes in Europe: distribution, habitat, usage and threats. In: European Atlas of forest tree species. eds: J. S.M.Ayanz, D. de Rigo, G. Caudullo, T. H. Durrant, A. Mauri Publication Office of the European Union, Luxembourg. pp: 184-185.
- [2] S. Jensen (2003) EUFORGEN. Technical Guidelines for genetic conservation and use for lime (*Tilia spp.*). International Plant Genetic Resources Institute, Rome, Italy. 6 pages. European Forest Genetic Resources Programme. Available from <http://www.euforgen.org/> Last accessed: 17.11.2019
- [3] University of Connecticut Plant Database. M. H. Brand, Department of Plant Science and Landscape Architecture, Storrs, CT 06269-4. Available form: <http://hort.uconn.edu/plants>. Last accessed:16.11.2019
- [4] M. Blumenthal (1998). American Botanical Council, Bundesinstitut für Arzneimittel und Medizinprodukte, Therapeutic Guide to Herbal Medicines, The Complete German Commission E monographs. American Botanical Council
- [5] R. L. Söukand, C. L. Quave, A. Pieroni, M. Pardo-de-Santayana, J. Tardío, R. Kalle, Ł. Łuczaj, I. Svanberg, V. Kolosova, L. Aceituno-Mata, G. Menendez-Baceta, I. Kołodziejska-Degórska, E. Pirożnikow, R. Petkevičius, A. Hajdari and B. Mustafa. (2013) Plants used for making recreational tea in Europe: a review based on specific research sites, *J Ethnobiol Ethnomed.* **9(1)**, 58 (12 pages).
- [6] B. Mustafa, A. Hajdari, A. Pieroni, B. Pulaj, X. Koro and C.L. Quave (2015). A cross-cultural comparison of folk plant uses among Albanians, Bosniaks, Gorani and Turks living in south Kosovo, *J Ethnobiol Ethnomed.* **12**;11:39 (26 pages).
- [7] A. Pieroni, R. Söukand, C.L. Quave, A. Hajdari and B. Mustafa (2017). Traditional food uses of wild plants among the Gorani of South Kosovo, *Appetite* **108**, 83-92.
- [8] A. Hajdari, A. Pieroni, M. Jhaveri, B. Mustafa and C.L. Quave (2018). Ethnopharmacological practices of slavic speaking people in South Kosovo, *Ethnobiol. Conserv.* **7(6)**, (42 pages).
- [9] B. Mustafa, A. Hajdari, F. Krasniqi, E. Hoxha, H. Ademi, C.L. Quave, and A. Pieroni (2012). Medical ethnobotany of the Albanian Alps in Kosovo, *J. Ethnobiol. Ethnomed.* **8**, 6 (14 pages).
- [10] B. Mustafa, A. Hajdari, Q. Pajaziti, B. Sylja, C.L. Quave and A. Pieroni (2012). An ethnobotanical survey of the Gollak region, Kosovo. *Genet. Resour. Crop Ev.* **69**,739-754.
- [11] European Medicines Agency. Available from <https://www.ema.europa.eu/en> Last accessed: 17.11.2019
- [12] G. Toker, M. Aslan, E. Yeşilada, M. Memişoğlu and S. Ito (2001). Comparative evaluation of the flavonoid content in officinal *Tiliaeflos* and Turkish lime species for quality assessment, *J. Pharmaceut. Biomed. Anal.* **26(1)**,111-1121.
- [13] A. Karioti, L. Chiarabini, A. Alachkar, M. Fawaz Chehna, F.F. Vincieri and A. R. Bilia (2014). HPLC-DAD and HPLC-ESI-MS analyses of *Tiliaeflos* and its preparations, *J. Pharmaceut. Biomed. Anal.* **100**, 205-214.
- [14] Z. Szűcsa, Z. Cziáky, A. Kiss-Szikszai, L. Sinka, G. Vasas and S. Gonda (2019).Comparative metabolomics of *Tilia platyphyllos* Scop. bracts during phenological development, *Phytochemistry* **167**,112084.
- [15] L. Ahmadi and M. Mirza (1999). Volatile constituents of *Tiliaapl typhyllos* Scop. flowers, *J. Essent. Oil Res.***11(6)**,765-766.
- [16] G. Toker , K. H. C. Baser, M. Kürkçüoğlu and T. Özek (1999). The composition of essential oils from *Tilia L.* species growing in Turkey, *J. Essent. Oil Res.* **11(3)**, 369-374.
- [17] A. Prączko and J. Góra (2000). The chemical composition of essential oil distilled from Linden blossoms, *Żywność* **4(25)**, 90-94.
- [18] V. Rădulescu and E. Oprea (2008). Analysis of volatile compounds of officinal *Tiliaeflos* by gas – chromatography coupled with mass spectrometry, *Farmacía* **56(2)**, 129-138.
- [19] R. Kowalski, T.Baj, K.Kalwa, G. Kowalska and M. Sujka (2017) Essential oil composition of *Tilia cordata* flowers, *J. Essent. Oil Bear. Pl.* **20(4)**, 1137-1142.
- [20] R. Adams (2007). Identification of essential oil components by gas chromatography/mass spectrometry, Allured, Carol Stream, Ill, USA, 4th edition.

A C G
publications

© 2020 ACG Publications