

Aromatic Compounds from the Marine-Derived Fungus

*Aspergillus versicolor*Rong Yao ^{1,2} *, Shen Yao ¹ and Ailian Hua ¹

¹ Department of Pharmacy, The First People's Hospital of Yuhang District, Hangzhou, Zhejiang
311100, China

² School of Pharmacy, Zhejiang University of Technology, Hangzhou 310014, China

(Received May 21, 2020; Revised June 06, 2020; Accepted June 12, 2020)

Abstract: Chemical investigation of the EtOAc extract of a marine-derived fungus *Aspergillus versicolor* SZW-5 led to the isolation of one new compound, named aspeverether A (**1**), and seven known compounds. The structures were determined by extensive analyses of spectroscopic data (1D and 2D NMR, and HRESIMS). Compounds **1** and **3** possessed an unusual 1,3-dimethoxy-2-vinylbenzene moiety, which is rarely found in nature. Compound **6** showed strong inhibitory effect toward α -glucosidase with an IC₅₀ value of 71 μ M, being more active than the positive acarbose (210 μ M).

Keywords: *Aspergillus versicolor*; aspeverether A; structure elucidation; marine. © 2020 ACG Publications. All rights reserved.

1. Plant Source

Fungus SZW-5 was isolated from the sediments that were collected at a depth of -3.5 m from the Shenzhen River Estuary. The strain was identified as *Aspergillus versicolor* by analysis of the internal transcribed spacer (ITS) region of the rDNA sequence. The ITS sequence has been submitted to the GenBank data base (<http://www.ncbi.nlm.nih.gov>) with the accession number MG845247. The strain SZW-5 was deposited at the Marine Culture Collection of China.

2. Previous Studies

In recent years, marine-derived fungi have attracted much attention from natural medicinal chemists, and many secondary metabolites possessing intriguing structures and various activities have been reported. The fungus *A. versicolor* has been proved to be prolific. Previous chemical study of marine-derived *A. versicolor* led to the identification of many secondary metabolites, including alkaloids [1-3], cyclopeptides [4-6], polyketides [7, 8], and depsides [9, 10], some exhibited

* Corresponding author: E- Mail: yaorong1205@163.com (R. Yao).

pronounced biological activities, such as cytotoxic [2], antimicrobial [1], lipid-lowering [6], and DPPH radical scavenging effects [10]. In our study of bioactive secondary metabolites of a marine-derived fungus *A. versicolor* SZW-5, eight aromatic compounds including a new one were obtained (Figure 1). Herein, the isolation, structural identification, and the inhibitory effects toward α -glucosidase of the metabolites were described.

3. Present Study

The solid-state fermentation was carried out in 15 erlenmeyer flasks (500 mL) with 70 g of rice and artificial sea-water (90 mL), and the contents were autoclaved at 15 psi for 30 min. Each flask was inoculated with 3.0 mL of the spore inoculum and incubated at room temperature for 35 days. The fermented materials were extracted with EtOAc (3 \times 2000 mL) and to give an EtOAc extract (3.1g), which was subjected to ODS silica gel column chromatography (CC) eluted with MeOH/H₂O (20:80 to 100:0) to afford five fractions (F1–F5). F3 was chromatographed over ODS silica gel CC (MeOH/H₂O: 20:80 to 100:0) to give six fractions (F3a–F3f). F3a was purified by HPLC using CH₃CN/H₂O (28:72, 3 mL/min) as eluent to afford **4** (3.2 mg, *t_R* 28.1 min) and **5** (1.9 mg, *t_R* 33.2 min). F3c was separated by ODS silica gel CC eluting with MeOH/H₂O (20:80 to 100:0) to give six fractions (F3d1–F3d6). F3d2 was separated by HPLC (MeOH/H₂O = 50:50, 3 mL/min) to yield five fractions (F3d2a–F3d2e). F3d3c was purified by HPLC (CH₃CN/H₂O = 41:59, 3 mL/min) to afford **6** (6.5 mg, *t_R* 19.5 min). F5 was further chromatographed over ODS silica gel CC eluted with MeOH/H₂O (20:80→100:0) to afford four subfractions (F5a–F5d). Purification of F5a by HPLC (CH₃CN/H₂O = 54:46, 3 mL/min) gave **7** (5.3 mg, *t_R* 9.7 min) and **8** (3.7 mg, *t_R* 11.9 min). F5d was purified by HPLC (CH₃CN/H₂O=68:32, 3 mL/min) to give **2** (14 mg, *t_R* 24.5 min), **3** (8.8 mg, *t_R* 22.2 min), and **1** (24 mg, *t_R* 24.2 min).

Aspeverether A (1): light yellow oil; UV (MeOH) λ_{\max} (log ϵ) 299 (3.81), 258 (4.22), 220 (4.39); ¹H and ¹³C NMR data, see Table 1; positive ESIMS *m/z* 363.1561 [M + Na]⁺ (calculated for C₂₁H₂₄O₄Na, 363.1567).

α -Glucosidase Assay: The α -glucosidase inhibitory effect was assessed as described in the literature [11].




Figure 1. Structures of compounds **1–8** from *A. versicolor* SZW-5.

Compound **1**, a light yellow oil, had the molecular formula C₂₁H₂₄O₄ as established by the HRESIMS *m/z* 363.1561 [M + Na]⁺ (calculated 363.1567) and NMR data (Table 1), requiring ten double bond equivalents. The ¹H NMR data exhibited the presences of three aromatic methoxys (δ_{H} 3.84, 9H), four olefinic protons [δ_{H} 6.96 (1H, d, *J* = 16.2 Hz), 6.76 (1H, dt, *J* = 16.2, 6.0 Hz), 6.38 (1H, dt, *J* = 16.1, 6.0 Hz), 6.98 (d, *J* = 16.1 Hz)], a symmetrically 1,2,3-trisubstituted benzene [δ_{H} 6.65 (2H, d, *J* = 8.4 Hz), 7.18 (1H, d, *J* = 8.4 Hz)], and a 1,2-disubstituted benzene [6.98 (1H, d, *J* = 7.8 Hz), 6.99 (1H, t, *J* = 7.8, 7.6 Hz); 7.24 (1H, td, *J* = 7.8, 1.5 Hz); 7.51 (1H, dd, *J* = 7.6, 1.5 Hz)] (Table

1). The ^{13}C NMR spectrum indicated the presences of twenty-one carbon resonances, including sixteen aromatic carbons for two benzenes and two double bonds, together with three methoxys (δ 56.1, 56.1, 55.8), and two oxymethylenes (δ_{C} 73.0, 71.4) (Table 1).

Table 1. ^1H (400 Hz) and ^{13}C NMR (100 Hz) Data of **1** in Acetone- d_6 (δ in ppm)

1					
No.	δ_{H} (mult., J in Hz)	δ_{C}	No.	δ_{H} (mult., J in Hz)	δ_{C}
1		114.7, C	1'		126.7, C
2		159.5, C	2'		157.7, C
3	6.65, d (8.4)	104.9, CH	3'	6.98, d (7.8)	111.9, CH
4	7.18, d (8.4)	129.1, CH	4'	7.24, td (7.8, 1.5)	129.5, CH
5	6.65, d (8.4)	104.9, CH	5'	6.99, t (7.8, 7.6)	127.3, CH
6		159.5, C	6'	7.51, dd (7.6, 1.5)	127.5, CH
7	6.96, d (16.2)	123.4, CH	7'	6.99, d (16.1)	123.4, CH
8	6.76, dt (16.2, 6.0)	131.0, CH	8'	6.38, dt (16.1, 6.0)	128.1, CH
9/9'	4.19, dt (6.0, 1.4)	73.0, CH_2	9'	4.19, dt (6.0, 1.4)	71.4, CH_2
2-OCH ₃	3.84, s	56.1, CH ₃	2''-OCH ₃	3.84, s	55.8, CH ₃
6-OCH ₃	3.84, s	56.1, CH ₃			

The HMBC correlations from the three aromatic methoxys (δ_{H} 3.84) to the oxygenated aromatic carbons (δ_{C} 159.5 \times 2, 157.7) indicated the presences of a 1,3-dimethoxy-1,2,3-tri substituted and a 2-methoxy-1,2-disubstituted benzene rings (Figure 2). The COSY spectrum indicated the ^1H - ^1H spin systems (H-7/H-8/H₂-9 and H-7'/H-8'/H₂-9') for two propenyl moiety. Additional HMBC correlations from H-7 (δ_{H} 6.96) and H-7' (δ_{H} 6.99) to C-1 and C-1' assigned an 1,3-dimethoxy cinnamyl and a 2-methoxy cinnamyl moieties. And these two cinnamyl moieties were connected via a ether linkage by the HMBC correlations from H-9 to C-9' and H-9' to C-9. Thus, the structure of **1** was established as depicted and was named aspeverether A. Compound **1** possessed an unusual 1,3-dimethoxy-2-vinylbenzene moiety, which is rarely found in nature.




Figure 2. ^1H - ^1H COSY (—), HMBC (\rightarrow), and NOE (\curvearrowright) correlations of **1**.

Besides, the known compounds were established as O-methoxy cinnamaldehyde (**2**) [12], 2,6-dimethoxy cinnamaldehyde (**3**) [13], altechromone A (**4**) [14], saccharonol A (**5**) [15], sulochrin (**6**) [16], questinol (**7**) [17], 7-hydroxyemodin (**8**) [18] by comparing their ^1H and ^{13}C NMR data with reported data in the literature.

Compounds **1**–**8** were tested for their inhibitory activities against α -glucosidase. As results, compound **6** exhibited an IC_{50} value of 71 μM , which is more active than the positive control acarbose (210 μM). And other compounds showed inhibitions less than 30% at the concentration of 200 μM .

Supporting Information

Supporting Information accompanies this paper on <http://www.acgpubs.org/journal/records-of-natural-products>

ORCID Rong Yao: [0000-0002-0015-7396](https://orcid.org/0000-0002-0015-7396)Shen Yao: [0000-0001-6731-0255](https://orcid.org/0000-0001-6731-0255)Ailian Hua: [0000-0003-1453-1112](https://orcid.org/0000-0003-1453-1112)

References

- [1] J. Wang, W. He, X. Huang, X. Tian, S. Liao, B. Yang, F. Wang, X. Zhou and Y. Liu (2016). Antifungal new oxepine-containing alkaloids and xanthenes from the deep-sea-derived fungus *Aspergillus versicolor* SCSIO 05879, *J. Agric. Food Chem.* **64**, 2910-2916.
- [2] K. Nakanishi, M. Doi, Y. Usami, T. Amagata, K. Minoura, R. Tanaka, A. Numata and T. Yamada (2013). Anthcolorins A-F, novel cytotoxic metabolites from a sea urchin-derived *Aspergillus versicolor*, *Tetrahedron* **69**, 4617-4623.
- [3] Z. Cheng, D. Liu, W. Cheng, P. Proksch and W. Lin (2018). Versiquinazolines L–Q, new polycyclic alkaloids from the marine-derived fungus *Aspergillus versicolor*, *RSC Adv.*, **8**, 31427-31439.
- [4] Y. Zhuang, X. Teng, Y. Wang, P. Liu, H. Wang, J. Li, G. Li and W. Zhu (2011). Cyclopeptides and polyketides from coral-associated fungus, *Aspergillus versicolor* LCJ-5-4, *Tetrahedron* **67**, 7085-7089.
- [5] J. Peng, H. Gao, X. Zhang, S. Wang, C. Wu, Q. Gu, P. Guo, T. Zhu and D. Li (2014). Psychophilins E-H and versicotide C, cyclic peptides from the marine-derived fungus *Aspergillus versicolor* ZLN-60, *J. Nat. Prod.* **77**, 2218-2223.
- [6] R. Chen, Z. Cheng, J. Huang, D. Liu, C. Wu, P. Guo and W. Lin (2017). Versicotides D-F, new cyclopeptides with lipid-lowering activities, *RSC Adv.*, **7**, 49235-49243.
- [7] Y. M. Lee, H. Li, J. Hong, H. Y. Cho, K. S. Bae, M. A. Kim, D.-K. Kim and J. H. Jung (2010). Bioactive metabolites from the sponge-derived fungus *Aspergillus versicolor*, *Arch. Pharmacol. Res.* **33**, 231-235.
- [8] W. Lin, G. Brauers, R. Ebel, V. Wray, A. Berg, Sudarsono and P. Proksch (2003). Novel chromone derivatives from the fungus *Aspergillus versicolor* isolated from the marine sponge *Xestospongia exigua*, *J. Nat. Prod.* **66**, 57-61.
- [9] H. Gao, L. Zhou, S. Cai, G. Zhang, T. Zhu, Q. Gu and D. Li (2013). Diorcinols B-E, new prenylated diphenyl ethers from the marine-derived fungus *Aspergillus versicolor* ZLN-60, *J. Antibiot.*, **66**, 539-542.
- [10] Z. Wu, Y. Wang, D. Liu, P. Proksch, S. Yu and W. Lin (2016). Antioxidative phenolic compounds from a marine-derived fungus *Aspergillus versicolor*, *Tetrahedron* **72**, 50-57.
- [11] Z. Cheng, Y. Li, W. Liu, L. Liu, J. Liu, W. Yuan, Z. Luo, W. Xu and Q. Li (2019). Butenolide derivatives with α -glucosidase inhibitions from the deep-sea-derived fungus *Aspergillus terreus* YPGA10, *Marin. Drugs* **17**, 332.
- [12] D. T. L. Huong, Y. S. Jo, M. K. Lee, K. H. Bae and Y. H. Kim (2000). Monoamine oxidase inhibitors from *Cinnamomi Cortex*, *Nat. Prod. Sci.* **6**, 16-19.
- [13] K. V. Rao and J. H. Johnson (1998). Occurrence of 2,6-dimethoxy cinnamaldehyde in *taxus floridana* and structural revision of taxiflorine to taxchinin M, *Phytochemistry* **49**, 1361-1364.
- [14] L. Liu, W. Xu, S. Li, M. Chen, Y. Cheng, W. Yuan, Z. Cheng and Q. Li (2019). Penicindopene A, a new indole diterpene from the deep-sea fungus *Penicillium* sp. YPCMAC1, *Nat. Prod. Res.* **33**, 2988-2994.
- [15] G. Du, Z.C. Wang, Y.K. Yang, H.M. Yang, H.Y. Yang, M. Zhou, Y.Q. Ye, X.M. Li and Q.F. Hu (2015). New isocoumarins from the fermentation products of entophytic fungus *Aspergillus versicolor* and their anti-tobacco mosaic virus activities, *Heterocycles* **91**, 1996-2002.
- [16] D. Liu, L. Yan, L. Ma, Y. Huang, X. Pan, W. Liu and Z. Lv (2015). Diphenyl derivatives from coastal saline soil fungus *Aspergillus iizukae*, *Arch. Pharm. Res.* **38**, 1038-1043.
- [17] D. Zhao, F. Cao, X.-J. Guo, Y.-R. Zhang, Z. Kang and H.-J. Zhu (2018). Antibacterial indole alkaloids and anthraquinones from a sewage-derived fungus *Eurotium* sp., *Chem. Nat. Compd.* **54**, 399-401.
- [18] G. Bringmann, G. Lang, S. Steffens, E. Günther and K. Schaumann (2003). Evariquinone, isoemicellin, and stromemycin from a sponge derived strain of the fungus *Emericella varicolor*, *Phytochemistry* **63**, 437-443.

A C G
publications

© 2020 ACG Publications

