

Two-steps synthesis of hexasubstituted porphyrins at the β -pyrrolic positions

Sylvia Ostrysz[†], Agnieszka Mikus^{ID*} and Stanisław Ostrowski^{ID*}

Faculty of Chemistry, Warsaw University of Technology, ul. Noakowskiego 3,
00-664 Warszawa, Poland

(Received March 30, 2025; Revised June 15, 2025; Accepted June 21, 2025)

Abstract: *meso*-Tetraarylporphyrin chelates are easily available from parent porphyrins. Their reactions with nitric acid (yellow fuming HNO₃, $d = 1.52$), under optimized conditions in CHCl₃ at room temperature, can be carried out selectively, thus giving mainly β,β,β -trinitro-substituted moieties (usually two or three isomers of the above complexes). These intermediates (Cu²⁺-derivatives; obtained in overall yields of up to 38%), upon the reaction with carbanions bearing a leaving group X at the reactive centre afforded the target products, the ones of *vicarious nucleophilic substitution of hydrogen* (VNS). This approach allows the synthesis of porphyrin derivatives exhaustively β -substituted in three pyrrole rings (in yields of up to 69%), starting from a simple *meso*-tetraarylporphyrin complexes. The above two-steps procedure including very attractive VNS reaction gives hydrophilic or amphiphilic moieties that seem to be potential agents in anticancer photodynamic therapy (PDT), practically unavailable by alternative methods.

Keywords: Porphyrin complexes; β -derivatization; carbanions; vicarious nucleophilic substitution; nitro group; photodynamic therapy. © 2025 ACG Publications. All right reserved.

1. Introduction

Porphyrins are intensively studied in the past decades as they can be used as valuable materials in many areas.¹ In this context, it needs to be highlighted that β -nitro-porphyrins (and derivatives), due to their interesting properties, are in the spotlight. For example, electron-withdrawing β -NO₂ groups affect the electronic structure of porphyrin chelates.² A blue shift of the Soret band and red shift of the peaks around 630 nm are observed with an increase in the electron-withdrawing effect of these groups. The electronic absorption bands shift to the near infrared region cause that these systems may find use in photodynamic therapy³⁻⁵ or dye-sensitized solar cells.⁶⁻¹⁰ They could be also considered as the most suitable models for mimicking the key roles of naturally occurring dyes in vital processes. Thus, synthesis of β -substituted porphyrins is described in many papers. Some of them regarding nucleophilic β -derivatization come from our laboratory.¹¹⁻¹⁶ This type of approach has also been investigated by other groups.¹⁷⁻²¹


In our ongoing research we are trying to functionalize the β,β,β -trinitroporphyrin derivatives *via* nucleophilic substitution reactions to prepare highly decorated products: penta- and hexa-substituted ones at the β -positions. It was planned to introduce carbon substituents into unoccupied sites,

[†] PhD student at the University of Podlasie, Siedlce

* Corresponding authors: E-mail: a.mikus@ch.pw.edu.pl; phone: +48 22-234-5063 (A. Mikus); stan@ch.pw.edu.pl; phone: +48 22-234-5063 (S. Ostrowski).

neighbouring the nitro groups. Thus, the previously synthesized trinitro isomers have been reacted with the respective carbanions.

Excellent tool to realize this task is the *vicarious nucleophilic substitution of hydrogen* process (VNS).^{22,23} Its mechanism is presented below using an example reaction of (2-nitro-5,10,15,20-tetraphenylporphyrinato)nickel(II) with chloromethyl phenyl sulphone, recently published in the literature (Scheme 1).¹⁴ VNS consists of addition of carbanion (also *O*-anions and *N*-anions) bearing a leaving group *X* at the reactive centre to nitroarene (and other electrophilic arenes and heteroarenes) at positions occupied by hydrogen to form σ^H -adduct(s) that are subsequently followed by base induced β -elimination of HX to give product(s) of nucleophilic substitution of hydrogen (see below).



Scheme 1. General scheme of the vicarious nucleophilic substitution of hydrogen in porphyrin system.

VNS is a very powerful reaction for derivatization of electrophilic aromatic and heteroaromatic systems. It is a process of general character and its application for porphyrin functionalization is known for more than two decades.²³ This approach was used earlier in our laboratory for preparation of di-, tri-, and tetrasubstituted porphyrins at the β -positions, affording the respective products with good or very good yield.¹¹⁻¹⁶ Substrates for this type of nucleophilic functionalization were β -nitro- and β,β -dinitro-porphyrins described in several publications.²⁴⁻³³ In the last year we also published a paper dealing with the synthesis of β,β,β -trinitro-isomers³⁴ that can be used (by applying VNS reaction) for transformation to obtain hexasubstituted porphyrinyl macrocycles. Such preliminary attempts are described in this communication.

2. Experimental

2.1. Chemical Materials and Apparatus

¹H NMR spectra were recorded with a Varian MR-400 spectrometer, operating at 400 MHz. Coupling constants *J* are expressed in hertz [Hz]. Mass spectra were measured with a GCT Premier (Waters, FD-TOF) spectrometer (FD method) and MARINER (PerSeptive Biosystems, ESI-TOF) spectrometer (ESI method); *m/z* intensity values for peaks are given as a % of relative intensity. UV-vis spectra were measured with a Metertech SP-8001 spectrophotometer. TLC analysis was performed on aluminium foil plates pre-coated with silica gel (60 F-254, Merck AG). All the products were isolated by column chromatography (silica gel, 230-400 mesh; Merck AG). Some compounds were rechromatographed on preparative TLC plates (silica gel, 60 F-254, 2 mm and 0.5 mm; Merck AG). Molecular formulas of new compounds were confirmed by HR-MS (ESI method), and by comparing the isotope molecular patterns (theoretical and experimental).

All the reactions were carried out in light-shielded flasks equipped with a septum. Preparation of starting β,β,β -trinitro-5,10,15,20-tetraphenylporphyrinate **1** was described in the earlier literature.³⁴ Carbanion precursors were also obtained according to known procedures: α -chloromethyl *para*-tolyl sulphone (**2a**),³⁵ α -bromomethyl *para*-tolyl sulphone (**2b**),³⁶ and α -chloromethyl phenyl sulphone (**2c**).³⁶

2.2. Chemistry

2.2.1. Reaction of 3,7,12-trinitro-5,10,15,20-tetraphenylporphyrin-copper(II) (**1**) with chloromethyl *para*-tolyl sulphone (**2a**) and bromomethyl *para*-tolyl sulphone (**2b**)

In a round-bottomed, light-shielded flask 3,7,12-trinitro-5,10,15,20-tetraphenylporphyrin-copper(II) (**1**; 65 mg, 0.080 mmol) was dissolved in anhydrous DMSO (30 mL). The mixture was stirred for ca 30 min due to solubility problem of porphyrinate. To this mixture carbanion precursor (**2a** – 225

Synthesis of hexasubstituted porphyrins at the β -pyrrolic positions

mg, 1.10 mmol; or **2b** – 317 mg, 1.27 mmol) was added in one portion and the solution obtained was dropped *via* syringe (under argon) to a stirred *t*-BuOK (230 mg, 2.05 mmol) in anhydrous DMSO (10 mL) over a period of *ca* 5 min. After the next 5 min of stirring the second portion of CH-acid (**2a**, 97 mg, 0.47 mmol; or **2b**, 140 mg, 0.56 mmol) was added followed by addition of the next portion of *t*-BuOK (101 mg, 0.90 mmol).

The reaction was continued at room temperature for the next 20 min. Then, the mixture was poured into 3% NH₄Cl containing ice (400 mL). The precipitate was filtered, washed with water (200 mL), and then dissolved in CHCl₃ (45 mL). After drying with anhydrous MgSO₄ and evaporation of the solvent, the products were isolated: (a) for **2a**: column chromatography (eluent: CHCl₃/MeOH, 100:1), to give a mixture of **4-6** and [3,7,12-trinitro-2,8,13-tris{(toluene-4-sulphonyl)methyl}-5,10,15,20-tetraphenylporphyrinato]copper(II) (**3**; 34.0 mg, 32%); the above mixture **4-6** was separated on preparative TLC (CHCl₃/MeOH, 100:1) to give three pentasubstituted isomers respectively: product $R_f=0.40$ (18.4 mg, 20%), product $R_f=0.35$ (6.4 mg, 7%), and product $R_f=0.28$ (20.2 mg, 22%); (b) for **2b**: the products were isolated on preparative TLC (eluent: CHCl₃/MeOH, 100:1) to give respectively: **3** (7.4 mg, 7%) and a mixture of β,β,β -trinitro- β,β -bis[(toluene-4-sulphonyl)methyl]-5,10,15,20-tetraphenylporphyrins–copper(II) **4-6** (47.7 mg, 52%).

[3,7,12-Trinitro-2,8,13-tris{(toluene-4-sulphonyl)methyl}-5,10,15,20-tetraphenylporphyrinato]copper(II) (**3**): $R_f=0.24$ [CHCl₃/MeOH (50:1)]. mp>300°C. UV-vis (CHCl₃): $\lambda_{\max}$, nm (log ϵ), 667 (3.88), 610 (3.83), 482.5 (4.80, Soret), 346 (4.25). MS (ESI): m/z (% rel. int.) 2659 (3), 2658 (5), 2657 (7), 2656 (11), 2655 (13), 2654 (13), 2653 (13), 2652 (8), 2651 (5) (isotope [2M+Na]⁺); 1342 (16), 1341 (34), 1340 (61), 1339 (100), 1338 (81), 1337 (96) (isotope [M+Na]⁺). HR-MS (ESI): m/z 1337.1812 (calcd. for C₆₈H₄₉N₇O₁₂S₃CuNa [(M+Na)⁺] 1337.1795).

Product $R_f=0.40$: $R_f=0.40$ [CHCl₃/MeOH (50:1)]. mp>300°C. UV-vis (CHCl₃): $\lambda_{\max}$, nm (log ϵ), 652.5 (3.97), 598 (3.97), 471.5 (4.96, Soret), 345 (4.33). MS (ESI): m/z (% rel. int.) 2323 (5), 2322 (8), 2321 (13), 2320 (18), 2319 (26), 2318 (28), 2317 (30), 2316 (19), 2315 (14) (isotope [2M+Na]⁺); 1174 (12), 1173 (26), 1172 (56), 1171 (90), 1170 (79), 1169 (100) (isotope [M+Na]⁺). HR-MS (ESI): m/z 1169.1534 (calcd. for C₆₀H₄₁N₇O₁₀S₂CuNa [(M+Na)⁺] 1169.1550). The molecular formula was also confirmed by comparing the theoretical and experimental isotope pattern for the [M+Na]⁺ ion (C₆₀H₄₁N₇O₁₀S₂CuNa; MS (ESI)) – it was found to be identical within the experimental error limits.

Product $R_f=0.35$: $R_f=0.35$ [CHCl₃/MeOH (50:1)]. mp>300°C. UV-vis (CHCl₃): $\lambda_{\max}$, nm (log ϵ), 651 (3.95), 600.5 (4.00), 477 (4.91, Soret), 345 (4.34). MS (ESI): m/z (% rel. int.): 2323 (6), 2322 (9), 2321 (14), 2320 (22), 2319 (31), 2318 (33), 2317 (33), 2316 (22), 2315 (16) (isotope [2M+Na]⁺); 1174 (14), 1173 (28), 1172 (58), 1171 (91), 1170 (80), 1169 (100) (isotope [M+Na]⁺). HR-MS (ESI): m/z 1169.1534 (calcd. for C₆₀H₄₁N₇O₁₀S₂CuNa [(M+Na)⁺] 1169.1550). The molecular formula was also confirmed by comparing the theoretical and experimental isotope pattern for the [M+Na]⁺ ion (C₆₀H₄₁N₇O₁₀S₂CuNa; MS (ESI)) – it was found to be identical within the experimental error limits.

Product $R_f=0.28$: $R_f=0.28$ [CHCl₃/MeOH (50:1)]. mp>300°C. UV-vis (CHCl₃): $\lambda_{\max}$, nm (log ϵ), 657 (3.63), 593 (3.48), 463 (4.57, Soret), 349.5 (3.96), 316 (4.06). MS (ESI): m/z (% rel. int.) 2323 (4), 2322 (6), 2321 (9), 2320 (13), 2319 (15), 2318 (19), 2317 (19), 2316 (13), 2315 (10) (isotope [2M+Na]⁺); 1173 (28), 1172 (57), 1171 (92), 1170 (79), 1169 (100) (isotope [M+Na]⁺). HR-MS (ESI): m/z 1169.1534 (calcd. for C₆₀H₄₁N₇O₁₀S₂CuNa [(M+Na)⁺] 1169.1550). The molecular formula was also confirmed by comparing the theoretical and experimental isotope pattern for the [M+Na]⁺ ion (C₆₀H₄₁N₇O₁₀S₂CuNa; MS (ESI)) – it was found to be identical within the experimental error limits.

2.2.2. Reaction of 3,7,12-trinitro-5,10,15,20-tetraphenylporphyrin-copper(II) (**1**) with chloromethyl phenyl sulphone (**2c**)

In a round-bottomed, light-shielded flask 3,7,12-trinitro-5,10,15,20-tetraphenylporphyrin–copper(II) (**1**; 60 mg, 0.074 mmol) in anhydrous DMSO (30 mL) was dissolved. The mixture was stirred for *ca* 30 min due to solubility problem of porphyrinate. To this mixture carbanion precursor **2c** (210

mg, 1.10 mmol) was added in one portion and the solution obtained was dropped *via* syringe (under argon) to a stirred *t*-BuOK (225 mg, 2.00 mmol) in anhydrous DMSO (10 mL) over a period of *ca* 5 min. After 5 min of stirring the second portion of CH-acid (**2c**, 88 mg, 0.46 mmol) was added followed by dropwise addition of the next portion of *t*-BuOK (120 mg, 1.07 mmol). The reaction was continued at room temperature for *ca* 20 min. Then, the mixture was poured into 3% NH₄Cl containing ice (400 mL). The precipitate was filtered off, washed with water (200 mL), and then dissolved in CHCl₃ (45 mL). After drying with anhydrous MgSO₄ and evaporation of the solvent, the product was isolated by column chromatography (eluent: CHCl₃, then CHCl₃/MeOH, 100:1). It was rechromatographed on preparative TLC (eluent: CHCl₃/MeOH, 100:1) to give [3,7,12-trinitro-2,8,13-tris{(phenylsulphonyl)methyl}-5,10,15,20-tetraphenylporphyrinato]copper(II) (**7**; 64.6 mg, 69%).

Another experiment:

Modifications: *t*-BuOK (540 mg, 4.81 mmol) in anhydrous DMSO (20 mL) was added to the solution of porphyrinate **1** and CH-acid **2c** – in reverse order to the previous reaction. The following products were obtained: (a) [3,7,12-trinitro-2,8,13-tris{(phenylsulphonyl)methyl}-5,10,15,20-tetraphenylpor-phyrinato]copper(II) (**7**) – 52.8 mg, 56%; (b) [3,7,12-trinitro-β,β-bis{(phenylsulphonyl)methyl}-5,10,15,20-tetraphenylporphyrinato]copper(II) (**8**)* – 5.2 mg, 6%; (c) [3,7,12-trinitro-β-{(phenylsul-phonyl)methyl}-5,10,15,20-tetraphenylporphyrinato]copper(II) (**9****) – 22.4 mg, 31%.

*) one of the three possible pentasubstituted isomers (the structure was not assigned)

**) one of the three possible tetrasubstituted isomers (the structure was not assigned)

[3,7,12-Trinitro-2,8,13-tris{(phenylsulphonyl)methyl}-5,10,15,20-tetraphenylporphyrinato]copper(II) (**7**): mp>300°C. UV-vis (CHCl₃): λ_{max}, nm (log ε), 672.5 (3.98), 611 (3.91), 485.5 (4.83, Soret), 349.5 (4.31). MS (FD): *m/z* (% rel. int.) 1279 (8), 1278 (12), 1277 (9), 1276 (26), 1275 (62), 1274 (100), 1273 (68), 1272 (96) (isotope M⁺). HR-MS (ESI): *m/z* 1272.1415 (calcd. for C₆₅H₄₃N₇O₁₂S₃Cu [M⁺] 1272.1428). The molecular formula was also confirmed by comparing the theoretical and experimental isotope pattern for the M⁺ ion (C₆₅H₄₃N₇O₁₂S₃Cu; MS (ESI)) – it was found to be identical within the experimental error limits.

[3,7,12-Trinitro-β,β-bis{(phenylsulphonyl)methyl}-5,10,15,20-tetraphenylporphyrinato]copper(II) (**8**): mp>300°C. UV-vis (CHCl₃): λ_{max}, nm, 646, 589, 466 (Soret). MS (ESI(-)): *m/z* (% rel. int.) 1158 (15), 1157 (29), 1156 (36), 1155 (64), 1154 (42), 1153 (60) (isotope [M+Cl]⁻); 1121 (21), 1120 (54), 1119 (85), 1118 (78), 1117 (100) (isotope [M-H]⁻). HR-MS (ESI): *m/z* 1117.1256 (calcd. for C₅₈H₃₆N₇O₁₀S₂Cu [(M-H)⁻] 1117.1261).

[3,7,12-Trinitro-β-{(phenylsulphonyl)methyl}-5,10,15,20-tetraphenylporphyrinato]copper(II) (**9**): mp>300°C. UV-vis (CHCl₃): λ_{max}, nm (log ε), 641 (3.45), 589 (3.36), 457 (4.43, Soret), 329 (3.75). MS (ESI(-)): *m/z* (% rel. int.) 1004 (13), 1003 (32), 1002 (53), 1001 (90), 1000 (59), 999 (100) (isotope [M+Cl]⁻); 968 (8), 967 (20), 966 (32), 965 (48), 964 (49), 963 (52) (isotope [M-H]⁻). MS (ESI(+)): *m/z* (% rel. int.) 991 (13), 990 (32), 989 (67), 988 (62), 987 (100) (isotope [M+Na]⁺). HR-MS (ESI): *m/z* 999.0917 (calcd. for C₅₁H₃₁N₇O₈SClCu [(M+Cl)⁻] 999.0939).

2.2.3. Demetallation of Copper(II)–Complexes of Hexasubstituted Products

In a round-bottomed flask (20 mL), to the corresponding copper(II)–porphyrinate (**3**: 33 mg or **7**: 32 mg; 0.025 mmol) concentrated H₂SO₄ (95%; 0.6 mL) was added followed by dropwise addition of TFA (3 mL). The mixture was stirred under argon at room temperature during *ca* 20 min. Then, it was poured into water (30 mL) and extracted with CHCl₃ (3×30 mL). The combined organic layers were washed with water (3×100 mL) and dried over anhydrous MgSO₄/Na₂CO₃. The solvent was evaporated to dryness. The residue was subjected to column chromatography using a mixture of CHCl₃/CH₃OH (100:1) as eluent, and then it was rechromatographed on preparative TLC (eluent: CHCl₃/MeOH, 100:1) to give free-base porphyrin **10** (19.4 mg, 62%) or **11** (21.8 mg, 72%).

3,7,12-Trinitro-2,8,13-tris[(toluene-4-sulphonyl)methyl]-5,10,15,20-tetraphenylporphyrin (**10**): mp>300°C. UV-vis (CHCl₃): λ_{max}, nm (log ε), 650 (3.52), 584 (3.45), 480 (4.70, Soret), 362 (4.08). ¹H


Synthesis of hexasubstituted porphyrins at the β -pyrrolic positions

NMR (CDCl_3 , 400 MHz): δ , ppm, 8.69 and 8.68 (AB, $J = 5.3$ Hz, 2H, H^β -pyrrole), 8.34–8.02 (m, 8H, H-Ph), 7.98–7.62 (m, 12H, H-Ph), 6.91 (apparent d, $J = 8.4$ Hz, 2H, H-Tol), 6.83 (apparent d, $J = 8.0$ Hz, 2H, H-Tol), 6.79–6.68 (m, 8H, H-Tol), 5.12–4.90 (m, 6H, $3 \times \text{CH}_2$), 2.21 (s, 3H, CH_3), 2.20 (s, 3H, CH_3), 2.16 (s, 3H, CH_3), NH – undetected. MS (ESI(+)): m/z (% rel. int.) 1281 (2), 1280 (6), 1279 (15), 1278 (37), 1277 (69), 1276 (84) (isotope $[\text{M}+\text{Na}]^+$); 1259 (2), 1258 (6), 1257 (17), 1256 (43), 1255 (77), 1254 (100) (isotope $[\text{M}+\text{H}]^+$). MS (ESI(-)): m/z (% rel. int.) 1257 (1), 1256 (5), 1255 (15), 1254 (39), 1253 (76), 1252 (100) (isotope $[\text{M}-\text{H}]^-$). The molecular formula was confirmed by comparing the theoretical and experimental isotope pattern for the $[\text{M}+\text{H}]^+$ ion ($\text{C}_{68}\text{H}_{52}\text{N}_7\text{O}_{12}\text{S}_3$; MS (ESI)) – it was found to be identical within the experimental error limits.

3,7,12-Trinitro-2,8,13-tris[(phenylsulphonyl)methyl]-5,10,15,20-tetraphenylporphyrin (11): mp > 300°C. UV-vis (CHCl_3): λ_{max} , nm (log ϵ), 658 (3.68), 592 (3.49), 483 (4.81, Soret), 358 (4.19). ^1H NMR (CDCl_3 , 400 MHz): δ , ppm, 8.75 and 8.73 (AB, $J = 5.2$ Hz, 2H, H^β -pyrrole), 8.38–8.12 (m, 8H, H-Ph), 7.96–7.54 (m, 12H, H-Ph), 7.24–6.52 (m, 15H, H-Ph), 5.18–4.90 (m, 6H, $3 \times \text{CH}_2$), NH – undetected. MS (ESI(+)): m/z (% rel. int.) 1238 (1), 1237 (5), 1236 (10), 1235 (18), 1234 (23) (isotope $[\text{M}+\text{Na}]^+$); 1217 (2), 1216 (6), 1215 (16), 1214 (39), 1213 (73), 1212 (100) (isotope $[\text{M}+\text{H}]^+$). MS (ESI(-)): m/z (% rel. int.) 1215 (1), 1214 (5), 1213 (13), 1212 (34), 1211 (72), 1210 (100) (isotope $[\text{M}-\text{H}]^-$). The molecular formula was confirmed by comparing the theoretical and experimental isotope pattern for the $[\text{M}+\text{H}]^+$ ion ($\text{C}_{65}\text{H}_{46}\text{N}_7\text{O}_{12}\text{S}_3$; MS (ESI)) – it was found to be identical within the experimental error limits.

3. Results and Discussion

(3,7,12-Trinitro-5,10,15,20-tetraphenylporphyrinato)copper(II) (**1**) was selected to illustrate the synthesis of highly decorated porphyrins at the β -positions. This substrate is now easily available.³⁴ In the first experiment, chloromethyl *para*-tolyl sulphone (**2a**) was reacted in DMSO with Cu(II)-complex **1** in the presence of *t*-BuOK. Carbanion generated from the above CH-acid, bearing good leaving group at the reactive centre, is a species of moderate nucleophilicity. However, it can effectively attack any position in the porphyrinate ring neighbouring the NO_2 group. Elimination of HCl with *tert*-butoxide anion that plays a role of the base led to the anionic form of the product (see VNS mechanism presented in Scheme 1). The double repetition of this reaction scheme followed by protonation resulted in the formation of hexasubstituted moieties at the rim of the porphyrin core ring (Scheme 2). It is worth noting that we are dealing with only a two-step procedure (electrophilic nitration/nucleophilic substitution of hydrogen).



Scheme 2. Reactions of nitroporphyrinate **1** with carbanions of halomethyl *para*-tolyl sulphones.


In this case, within 30 min from the beginning of the reaction, all the porphyrin substrate disappeared (TLC control). We isolated from the post-reaction mixture (by column chromatography) two fractions. The first one consists of three possible pentasubstituted isomers **4-6** (three spots on TLC, with a total yield of 49%), while the next fraction was the desired pure hexasubstituted product **3** (32%; $R_f = 0.24$, $\text{CHCl}_3/\text{MeOH}$, 50:1) (Scheme 2). Thus, the combined yield of all four products reached 81%.

The R_f values of the isomers **4-6** are close to each other, however preparative TLC chromatography allowed us to separate them and partially characterize ($R_f = 0.40$, 18%; $R_f = 0.35$, 13%; and $R_f = 0.28$, 18%; respectively). The spectral data are in agreement with the postulated structures, *e.g.*, in all MS spectra the intensive ion $m/z = 1169$ ($[\text{M}+\text{Na}]^+$) was observed. We did not investigate further the problem concerning assignment of the structure to the particular isomers. It is rather a side issue here. First of all, this is not a trivial task. On the other hand, they are not the aims of this study.

Minor optimization of the reaction conditions, namely prolongation of the reaction time (to 2 hours) did not give better results. Instead of increasing conversion to exhaustively substituted moiety **3**, a spontaneous degradation of the products occurred. The total yield dropped from 81% to 69%; in particular, the yield of main product has decreased, from 32% to 20%.

Reaction of the above porphyrinate **1** with carbanion of bromomethyl *para*-tolyl sulphone ($^-\text{CH}(\text{Br})\text{SO}_2\text{Tol}$; **2b**) which is a species of comparable nucleophilicity to $^-\text{CH}(\text{Cl})\text{SO}_2\text{Tol}$ gave a mixture of the same target compounds, however the main product was formed with 7% yield only (accompanied with 52% of pentasubstituted derivatives **4-6**; see Scheme 2). A reasonable explanation of this result is that after addition of carbanion containing better leaving group ($\text{X} = \text{Br}$) to porphyrin moiety the σ^{H} -adduct(s) of the latter will undergo β -elimination faster to give anionic form of VNS product(s). Unfortunately, this triple anion is also more unstable what was observed in the first experiment (for $\text{X} = \text{Cl}$; reaction time: 2 h).


The best yield of the exhaustively hexasubstituted product was observed in the reaction of copper(II)-porphyrinate **1** with carbanion of chloromethyl phenyl sulphone (**2c**). Additionally, penta- and tetrasubstituted derivatives **8** and **9** were formed (Scheme 3). The combined yield was very high (93%; 56% of product **7**). Optimization of this reaction (in particular, the reverse order of addition of the reagents and variable amount of base) allowed us to obtain the desired compound **7** as the only product and achieve the yield of 69%.



Scheme 3. Reactions of nitroporphyrinate **1** with carbanion of chloromethyl phenyl sulphone.

For more detailed and accurate characterization of paramagnetic compounds **3** and **7** they were decomplexed in $\text{H}_2\text{SO}_4/\text{CF}_3\text{CO}_2\text{H}$ mixture to give free-base porphyrins **10** and **11**, which were analyzed including ^1H NMR (Scheme 4). In ^1H NMR spectrum of **10** one can find AB system originating from the remaining two β -pyrrole protons (8.68 and 8.69 ppm, coupled with $J = 5.3$ Hz), signals from CH_2 groups (4.90-5.12 ppm), and three intensive singlets at 2.16, 2.20, and 2.21 of $\text{CH}_3\text{-Tol}$ groups. They confirm the structure.

Synthesis of hexasubstituted porphyrins at the β -pyrrolic positions



Scheme 4. Demetallation of copper(II)-complexes of hexasubstituted products.

We also characterized free-base of multisubstituted porphyrinyl derivative **11**, and observed in ¹H NMR spectra a diagnostic AB system originating from the protons of unsubstituted pyrrole ring (8.73 and 8.75 ppm; *J* = 5.2 Hz). Signals from CH₂ groups appeared in the region 4.90-5.18 ppm.

4. Conclusions

In summary, the studies described in this communication focused on the synthesis of multisubstituted at the β -positions porphyrins. We demonstrated two-step procedure (electrophilic nitration / VNS) for preparation of highly derivatized moieties from parent *meso*-tetraphenylporphyrin Cu(II)-chelates. This type of functionalization gave rise to amphiphilic compounds with six polar groups on the rim of the core ring (*e.g.* NO₂, SO₂R; three pyrrole rings are exhaustively substituted). The products obtained would find interesting applications and can be considered as valuable intermediates for further transformations. Among others, such derivatives are sought in anticancer photodynamic therapy (PDT).

ORCID

Agnieszka Mikus: [0000-0002-0191-6024](https://orcid.org/0000-0002-0191-6024)

Stanisław Ostrowski: [0000-0003-3903-3769](https://orcid.org/0000-0003-3903-3769)

References

- [1] Kadish, K. M.; Smith, K. M.; Guillard, R. Eds; Handbook of Porphyrin Science, World Scientific Publishing Co.: New Jersey, London, Singapore, Beijing, Shanghai, Hong Kong, Taipei, Chennai, 2010–2022, vol. 1–46.
- [2] Fujii, H. Electronic structure and reactivity of high-valent oxo iron porphyrins. *Coord. Chem. Rev.* **2002**, 226, 51–60.
- [3] Denis, T. G. St.; Huang, Y.-Y.; Hamblin, M. R. Cyclic tetrapyrroles in photodynamic therapy: the chemistry of porphyrins and related compounds in medicine. In ref. [1]: Vol. 27, p. 255.
- [4] Singh, S.; Aggarwal, A.; Bhupathiraju, N. V. S. D. K.; Arianna, G.; Tiwari, K.; Drain, Ch. M. Glycosylated porphyrins, phthalocyanines, and other porphyrinoids for diagnostics and therapeutics. *Chem. Rev.* **2015**, 115, 10261–10306, and references therein.
- [5] Abrahamse, H.; Hamblin, M. R. New photosensitizers for photodynamic therapy. *Biophys. J.* **2016**, 473, 347–364.
- [6] Campbell, W. M.; Jolley, K. W.; Wagner, P.; Wagner, K.; Walsh, P. J.; Gordon, K. C.; Schmidt-Mende, L.; Nazeeruddin, M. K.; Wang, Q.; Grätzel, M.; Officer, D. L. Highly efficient porphyrin sensitizers for dye-sensitized solar cells. *J. Phys. Chem. C* **2007**, 111, 11760–11762.
- [7] Ishida, M.; Park, S. W.; Hwang, D.; Koo, Y. B.; Sessler, J. L.; Kim, D. Y.; Kim, D. Donor-substituted β -functionalized porphyrin dyes on hierarchically structured mesoporous TiO₂ spheres. Highly efficient dye-sensitized solar cells. *J. Phys. Chem. C* **2011**, 115, 19343–19354.
- [8] Li, L.-L.; Diau, E. W.-G. Porphyrin-sensitized solar cells. *Chem. Soc. Rev.* **2013**, 42, 291–304.
- [9] Higashino, T.; Imahori, H. Porphyrins as excellent dyes for dye-sensitized solar cells: recent developments and insights. *Dalton Trans.* **2015**, 44, 448–463.
- [10] Puttaiah, B. Recent advances in mixed β -pyrrole substituted *meso*-tetraphenylporphyrins. *Tetrahedron Lett.* **2016**, 57, 5150–5167.

- [11] Ostrowski, S.; Raczko, A. M. Selective double functionalization of *meso*-tetraphenylporphyrin complexes on the same pyrrole unit by tandem electrophilic/nucleophilic aromatic substitution. *Helv. Chim. Acta* **2005**, *88*, 974-978.
- [12] Ostrowski, S.; Grzyb, S. Direct β -amination reaction in porphyrin systems – a simple route to compounds containing two nitrogen substituents at both β -positions of the same pyrrole unit. *Tetrahedron Lett.* **2012**, *53*, 6355-6357.
- [13] Rosa, M.; Ostrowski, S. Studies on synthesis of porphyrins – β -substituted with multiple nitrogen-containing groups. *ChemistrySelect*, **2022**, *7*: e202200290.
- [14] Mikus, A.; Ostrowski, S. Vicarious nucleophilic substitution of hydrogen and synthesis of nickel-, copper-, and cobalt-porphyrinates – double functionalized at the β -positions of the same pyrrole unit. *J. Porphyrins Phthalocyanines* **2024**, *28*, 647-653.
- [15] Mikus, A. In two steps from simple *meso*-tetraphenylporphyrin to its derivatives exhaustively β -substituted at the 'Eastern half'. *Mendeleev Commun.* **2024**, *34*, 70-73.
- [16] Mikus, A.; Ostrysz, S.; Ostrowski, S. Fast nucleophilic approach to exhaustively substituted porphyrin derivatives at the 'Eastern half'. *Eur. J. Org. Chem.*; DOI: 10.1002/ejoc.202401054 (accepted).
- [17] Malinovskii, V. L.; Vodinskii, S. V.; Zhilina, Z. I.; Andronati, S. A.; Mazepa, A. V. Porphyrins and their derivatives. Part 18. Vicarious Nucleophilic Substitution of Hydrogen in 2-Nitro-5,10,15,20-tetraphenylporphyrin. *Zh. Org. Khim.* **1996**, *32*, 119-123.
- [18] Richeter, S.; Jeandon, Ch.; Ruppert, R.; Callot, H. J. Reactivity of oxonaphthoporphyrins. Efficient β -functionalization of the porphyrin ring on reaction with nitrogen or carbon nucleophiles. *Tetrahedron Lett.* **2001**, *42*, 2103-2106.
- [19] Eddahmi, M.; Moura, N. M. M.; Ramos, C. I. V.; Bouissane, L.; Faustino, M. A. F.; Cavaleiro, J. A. S.; Rakib, E. M.; Neves, M. G. P. M. S. An insight into the vicarious nucleophilic substitution reaction of 2-nitro-5,10,15,20-tetraphenylporphyrin with *p*-chlorophenoxyacetonitrile: Synthesis and gas-phase fragmentation studies. *Arabian J. Chem.* **2020**, *13*, 5849-5863.
- [20] Richeter, S.; Hadj-Aïssa, A.; Taffin, C.; van der Lee, A.; Leclercq, D. Synthesis and structural characterisation of the first *N*-heterocyclic carbene ligand fused to a porphyrin. *Chem. Commun.* **2007**, 2148-2150.
- [21] Lefebvre, J.-F.; Leclercq, D.; Gisselbrecht, J.-P.; Richeter, S. Synthesis, characterization, and electronic properties of metalloporphyrins annulated to exocyclic imidazole and imidazolium rings. *Eur. J. Org. Chem.* **2010**, 1912-1920.
- [22] Małosza, M. Nucleophilic substitution of hydrogen in electron-deficient arenes, a general process of practical value. *Chem. Soc. Rev.* **2010**, *39*, 2855-2868 (review).
- [23] Ostrowski, S.; Mikus, A. Vicarious nucleophilic substitution of hydrogen – excellent tool for porphyrins functionalization. *SynOpen*; DOI: 10.1055/s-0040-1720167 (review, accepted).
- [24] Evans, B.; Smith, K. M.; Cavaleiro, J. A. S. Bile pigment studies. Part 4. Some novel reactions of metalloporphyrins with thallium(III) and cerium(IV) salts. Ring cleavage of *meso*-tetraphenylporphyrin. *J. Chem. Soc., Perkin Trans.1* **1978**, 768-773.
- [25] Shine, H. J.; Padilla, A. G.; Wu, S.-M. Ion radicals. 45. Reactions of zinc tetraphenylporphyrin cation radical perchlorate with nucleophiles. *J. Org. Chem.* **1979**, *44*, 4069-4075.
- [26] Baldwin, J. E.; Crossley, M. J.; DeBernardis, J. Efficient peripheral functionalization of porphyrins. *Tetrahedron* **1982**, *38*, 685-692.
- [27] Catalano, M. M.; Crossley, M. J.; Harding, M. M.; King, L. G. Control of reactivity at the porphyrin periphery by metal ion co-ordination: a general method for specific nitration at the β -pyrrolic position of 5,10,15,20-tetraarylporphyrins. *J. Chem. Soc., Chem. Commun.* **1984**, 1535-1536.
- [28] Crossley, M. J.; King, L. G. A new method for regiospecific deuteration and reduction of 5,10,15,20-tetraphenylporphyrins: nucleophilic reaction of borohydride ion with 2-nitro-5,10,15,20-tetraphenylporphyrins. *J. Org. Chem.* **1993**, *58*, 4370-4375.
- [29] Hombrecher, H. K.; Gherdan, V. M.; Ohm, S.; Cavaleiro, J. A. S.; Neves, M. G. P. M. S.; Condesso, F. M. Synthesis and electrochemical investigation of β -alkyloxy substituted *meso*-tetraphenylporphyrins. *Tetrahedron* **1993**, *49*, 8569-8578.
- [30] Ostrowski, S.; Szerszeń, D.; Ryszczuk, M. Electrophilic nitration of *meso*-tetraarylporphyrin complexes at the β -pyrrolic position. *Synthesis* **2005**, 819-823.
- [31] Ostrowski, S. Preparation of 2-nitro-5,10,15,20-tetraphenylporphyrin zinc(II) complex – *post scriptum* to electrophilic nitration of porphyrins at the β -pyrrolic position. *Polish J. Chem.* **2005**, *79*, 1169-1172.
- [32] Wyřbek, P.; Ostrowski, S. Synthesis of some β -nitro-*meso*-tetraphenylporphyrin derivatives. *J. Porphyrins Phthalocyanines* **2007**, *11*, 822-828.

Synthesis of hexasubstituted porphyrins at the β -pyrrolic positions

- [33] (a) Mikus, A.; Rosa, M.; Ostrowski, S. Isomers of β,β -dinitro-5,10,15,20-tetraphenylporphyrin derivatives: valuable starting materials for further transformations. *Molecules* **2019**, *24*, 838. (b) Mikus, A.; Ostrowski, S. Revised structure of 2,13-dinitro-5,10,15,20-tetraphenylporphyrin. Inside high-resolution NMR. *Struct. Chem.* **2022**, *33*, 1251-1255. (c) Mikus, A.; Ostrowski, S. Correction to: Revised structure of 2,13-dinitro-5,10,15,20-tetraphenylporphyrin. Inside high-resolution NMR. *Struct. Chem.* **2022**, *33*, 2261.
- [34] Ostysz, S.; Mikus, A.; Ostrowski, S. Synthesis and structure elucidation of β,β,β -trinitro-*meso*-tetraphenylporphyrin derivatives. *Macroheterocycles* **2024**, *17*, 9-15.
- [35] Mąkosza, M.; Danikiewicz, W.; Wojciechowski, K. Reactions of chloroalkyl sulfones with nitronaphthalene derivatives. *Liebigs Ann. Chem.* **1987**, 711-715.
- [36] Mąkosza, M.; Goliński, J.; Baran, J. Vicarious nucleophilic substitution of hydrogen in nitroarenes with carbanions of α -haloalkyl phenyl sulfones. *J. Org. Chem.* **1984**, *49*, 1488-1494.

A C G
publications

© 2025 ACG Publications