

Supporting Information

Org. Commun. 12:3 (2019) 143-148

A study of Minisci reaction by changing Fe²⁺ equivalency: Preparation of arylpyridinyl methanol

Bünyamin Özgeriş¹ and Ferhan Tümer^{2*}

¹*Erzurum Technical University, Faculty of Science, Department of Basic Sciences, Erzurum, Türkiye*

²*Faculty of Science and Arts, Department of Chemistry, Sutcu Imam University, Kahramanmaraş,
Türkiye*

Table of content	Page
Figure S1 : 400 MHz ¹ H-NMR and 100 MHz ¹³ C-NMR (DMSO-d6) spectrums of 9	2
Figure S2 : 400 MHz ¹ H-NMR and 100 MHz ¹³ C-NMR (DMSO-d6) spectrums of 12	3
Figure S3 :400 MHz ¹ H-NMR and 100 MHz ¹³ C-NMR (DMSO-d6) spectrums of 13	4
Figure S4 : 400 MHz ¹ H-NMR and 100 MHz ¹³ C-NMR (DMSO-d6) spectrums of 14	5
Figure S5 :400 MHz ¹ H-NMR and 100 MHz ¹³ C-NMR (DMSO-d6) spectrums of 15	6




Figure S1: 400 MHz ¹H-NMR and 100 MHz ¹³C-NMR Spectra of compound 9 (DMSO-d₆)




Figure S2: 400 MHz ^1H -NMR and 100 MHz ^{13}C -NMR Spectra of compound 12 (DMSO-d₆)




Figure S3: 400 MHz ¹H-NMR and 100 MHz ¹³C-NMR Spectra of compound 13 (DMSO-d₆)




Figure S4: 400 MHz ^1H -NMR and 100 MHz ^{13}C -NMR Spectra of compound 14 (DMSO-d₆)





Figure S5: 400 MHz ^1H -NMR and 100 MHz ^{13}C -NMR Spectra of compound 15 (DMSO-d₆)