

Supporting Information

Rec. Nat. Prod. X:X (202X) XX-XX

The Leaf and the Gall Volatiles of *Salvia fruticosa* Miller from Turkey: Chemical Composition and Biological Activities

Sevda Süzgeç-Selçuk^{1*}, Temel Özek^{2,3}, Gülmira Özek², Süleyman Yur^{2,3}, Fatih Göger^{2,3}, M. Bahar Gürdal⁴, Gizem Gülsoy Toplan¹, Ali H. Meriçli⁵ and K. Hüsnü Can Başer⁵

¹ Istanbul University, Faculty of Pharmacy, Department of Pharmacognosy, 34116 Istanbul, Türkiye

² Anadolu University, Faculty of Pharmacy, Department of Pharmacognosy, 26470, Eskişehir, Türkiye

³ Medicinal Plant, Drug and Scientific Research and Application Center (AUBIBAM), Anadolu University, 26470 Eskişehir, Türkiye

⁴ Istanbul University, Faculty of Pharmacy, Department of Pharmaceutical Botany, 34116, Istanbul, Türkiye

⁵ Near East University, Faculty of Pharmacy, Department of Pharmacognosy, Nicosia, Turkish Republic of Northern Cyprus

Table of Contents		Page
Figure S1. Distribution of the major compound groups in the leaf (L) and gall (G) oils of <i>S. fruticosa</i> obtained with hydrodistillation (HD).		2
Figure S2 : Chromatographic profile of the volatiles obtained with hydrodistillation (HD) and MSD-SPME techniques from the leaves (L) and the galls (G) of <i>S. fruticosa</i>		3




Figure S1. Distribution of the major compound groups in the leaf (L) and gall (G) oils of *S. fruticosa* obtained with hydrodistillation (HD).




Figure S2 : Chromatographic profile of the volatiles obtained with hydrodistillation (HD) and MSD-SPME techniques from the leaves (L) and the galls (G) of *S. fruticosa*