

Rec. Nat. Prod. X:X (2022) XX-XX

records of natural products

Terpenes from the Fresh Stems of *Commiphora gileadensis* with Antimicrobial Activity

Maged S. Abdel-Kader (1)1,2*, Elmutasim O. Ibnouf (1)3,4,

Mohammed H. Alqarni (1)1, Abdallah S. AlQutaym (1)1, Ayman A. Salkini (1)1

and Ahmed I. Foudah (1)1

¹Department of Pharmacognosy, College of Pharmacy, Prince Sattam Bin Abdulaziz University, P.O.

Box 173, Al-Kharj 11942, Saudi Arabia

²Department of Pharmacognosy, Faculty of Pharmacy, Alexandria University, Alexandria 21215, Egypt

Box 173, Al-Kharj 11942, Saudi Arabia

⁴Department of Medical Microbiology, Faculty of Medical Laboratory Sciences, Omdurman Islamic University, Sudan

(Received 17 February, 2022; Revised March 13, 2022; Accepted April 12, 2022)

Abstract: Phytochemical study directed by antimicrobial activity of the fresh *Commiphora gileadensis* stems CHCl₃ extract resulted in the isolation of compounds **1-9**. The structures of the isolated compounds were elucidated utilizing various spectroscopic methods including 1D, 2D NMR and HRESIMS. Compounds **1**, **3-5** were identified as the *ent*-verticillane-type diterpenes (*1S*, *3E*, *7E*, *11R*)-(+)-verticilla-3,7,12(18)-triene (**1**), (*13S*, *14S*)-*Ent*-13,14-epoxyverticillol (**3**), (*9S*, *10S*)-*Ent*-9,10-epoxyverticillol (**4**) and *Ent*-Verticillol (**5**) for the first time from *Commiphora* species. Compound **9** was identified as a diterpene with new skeleton and was given the name "gileadenol". In addition, four know triterpenes were also identified as friedelin (**2**), oleanonic aldehyde (**6**), canyophyllal (**7**) and urs-12-en-3-one-28-al (**8**). The three tested *ent*-verticillane-type diterpenes expressed promising antimicrobial activity especially against *K. pneumonia*. The new diterpene as well as canyophyllal were less active.

Keywords: *Commiphora gileadensis*; antimicrobial; *ent*-verticillane-type diterpenes; gileadenol; triterpenes. © 2022 ACG Publications. All rights reserved.

1. Introduction

Genus Commiphora is represented by about 7 species in Saudi Arabia particularly in the southwest region [1]. Commiphora gileadensis (L) C.Chr. is synonymous with Commiphora opobalsamum [2]. The plant bark is the source of the balm of Gilead. Different parts of the plant are used in the traditional medicine. The dried layer under the bark was used to treat infected wounds [3]. The resins of

³Department of Pharmaceutics, College of Pharmacy, Prince Sattam Bin Abdulaziz University, P.O.

^{*} Corresponding author: E-Mail: mpharm101@hotmail.com

Commiphora species are used for the treatment of microbial infections, wounds, inflammation, gastrointestinal diseases, tumors, obesity, pain, arthritis and fractures [4].

The methanol extract of *C. gileadensis* bark as well as the callus extracts of the plant showed a broad-spectrum antimicrobial effect against Gram-positive, Gram-negative bacteria and fungi [5,6]. *C. gileadensis* can interfere with bacterial lectin-dependent adhesion important for the function and survival of *P. aeruginosa*. These properties support the old claims of using the balsam as anti-infective agent [7]. The methanol extract of *C. gileadensis* leaves demonstrated antiviral activity via target the enveloped viruses (HSV-2 and RSV B) [8]. *C. gileadensis* leaf and twig aqueous extracts controlled alloxan-induced diabetes in hypercholesterolemic male rats [2]. The plant also expressed potential to protect against acute gastric ulcers induced by different necrotizing agents in rats [9] and protective effect against carbon tetrachloride induced hepatotoxicity as well [10].

Previous phytochemical study of the genus revealed the presence of pimarane, abietane diterpenoids and dammarane triterpenoids [11,12]. Triterpenoids are the major constituents isolated from *Commiphora* species. More than twenty-one dammarane triterpenoids were identified in the resins of four *Commiphora* species namely; *C. kua*, *C. dalzieli* [13], *C. confuse* and *C. myrrha* [14]. The flavonoids of this genus occurs mainly in the flowers, stems and barks, but not in the resinous exudates. Phytochemical study of the stem bark of *C. opobalsamum* resulted in the identification of six prenylated flavonoids comophorin A-E with cytotoxic activity against MCF-7 and HepG-2 cell lines [15].

The current study designed to identify the antimicrobial constituents from the stems of *C. gileadensis* by conducting detailed biologically directed phytochemical study.

2. Materials and Methods

2.1. General

Melting points were measured using open capillary tubes Thermosystem FP800 Mettler FP80 central processor supplied with FP81 MBC cell apparatus and were uncorrected. Infrared (IR) spectra were recorded on FT-IR spectrophotometer (Perkin Elmer) as film. ¹H, ¹³C-NMR, and 2D-NMR data were collected on a Bruker UltraShield Plus 500 MHz spectrometer at the NMR Unite, College of Pharmacy, Prince Sattam Bin Abdulaziz operated at 500 MHz for protons and 125 MHz for carbons, respectively. Chemical shift values were reported in δ (ppm) relative to the residual solvents peaks. Coupling constants (J) were reported in Hertz (Hz). HRESIMS were determined by direct injection using Thermo Scientific UPLC RS Ultimate 3000 - Q Exactive hybrid quadrupole-Orbitrap mass spectrometer combined with high-performance quadrupole precursor selection with high resolution, accurate mass (HR/AM) OrbitrapTM detection. MPLC was performed using a Buchi medium pressure system composed of Buchi pump module C-605 controlled by Buchi control unit C-620 equipped with Buchi fraction collector C-660. The column eluate was detected by Buchi UV photometer C-640. Column 15/460-044032 was used and the system was operated by Sepacore control chromatography software. Sephadex LH20 (Amersham Biosciences, Uppsala, Sweden), Silica gel 60/230-400 mesh (EM Science), (Loba Chemie PVT. LTD. 100- 200 Mesh) and RP18 silica gel 40-63/230-400 mesh (Fluka) were used for column chromatography. The thin-layer chromatography (TLC) analysis was performed on Kiesel gel 60 F254 and RP18 F254 (Merck) plates. A UV lamp (entela Model UVGL-25) operated at 254 nm was used for spots detection on the TLC plates.

2.2. Plant materials

The plants of *Commiphora gileadensis* (L.) C.Chr (Syn. *Commiphora opobalsamum* (L.) Engl.) were collected from "Badr" area near Al-Madinah city during November, 2019. The plants were identified by Prof Saniya Kamal, Prof of Taxonomy, College of Science, Alexandria University. Voucher specimen (ALX 111119) was deposited at the Department of Botany, College of Science, Alexandria University.

2.3. Extraction and isolation

Fresh Stems (5.8 kg) were extracted with CHCl₃ at room temperature till exhaustion to yield 117.5 g of the CHCl₃ extract after evaporation of the solvent under reduces pressure. The stems were then extracted with MeOH at room temperature till exhaustion. Evaporation of the solvent under reduced pressure yielded 226.7 g of the MeOH soluble extract.

Column chromatography for 20 g of the $CHCl_3$ extract was preformed using silica gel (400 g, 5 cm i.d.) eluting with 5% EtOAc in hexane with increasing the percentage of EtOAc in a gradient system. Fractions of 250 mL each were collected, screened with TLC and similar fractions were pooled to yield ten fractions A- J.

Fraction A (4.69 g) eluted with 5% EtOAc was subjected to fractionation over Sephadex LH20 column (150 g, 3 cm i.d.) starting with petroleum ether followed by petroleum ether EtOAc mixtures with 25% increase in EtOAc contents each time. Fractions of 15 mL each were collected, screened with TLC and similar fractions were pooled. Fraction A-7 (136 mg) was subjected to Preparative TLC (PTLC) on silica gel plates using 9:1 petroleum ether / EtOAc mixture as developing system to yield 10 mg of compound 1. Fraction A-13 (64 mg) was subjected to PTLC on silica gel plates using 9:1 petroleum ether / EtOAc mixture developing system to yield 52 mg of compound 2.

Fraction B (4.99 g) eluted with 10% EtOAc was subjected to column chromatography on silica gel column (150 g, 3 cm i.d.) eluting with 5% EtOAc. Fractions of 100 mL each were collected, screened with TLC and similar fractions were pooled. Fraction B-a (1.2 g) was fractionated over MPLC silica gel column (92 cm, 1.5 cm i.d.) eluting with petroleum ether followed by petroleum ether EtOAc mixtures in a gradient system. Fractions of 15 mL each were collected, screened with TLC and similar fractions were pooled. Fraction B-a-12 (625 mg) was further purified on MPLC RP18 column eluted with 40% MeOH in water with gradual increase of MeOH in a gradient system. Fraction Fractions of 10 mL each were collected, screened with TLC and similar fractions were pooled. Fraction 64 eluted with 80% MeOH afforded 8 mg of 3. Fraction 67 eluted with 85% MeOH afforded 6 mg of 4. Fractions 71-75 eluted with 85% MeOH afforded 213 mg of 5. Fractions 80-83 eluted with 90% MeOH afforded 86 mg of 6. Fractions 87-92 eluted with 95% MeOH afforded 16 mg of 7. Fraction B-a-13 (161 mg) was repurified over silica gel column (20 g, 1.5 cm) eluting with petroleum ether followed by petroleum ether EtOAc mixtures in a gradient system. Fractions 19- 20 afforded compounds 6 (8 mg), 8 (16 mg) and 20 mg of 9 after PTLC using silica gel plates and 9:1 petroleum ether / EtOAc mixture as developing system. Plates were developed for four times. Fraction B-a-14 (132 mg) was subjected to silica gel column (20 g, 1.5 cm) eluting with petroleum ether followed by petroleum ether EtOAc mixtures in a gradient system to afford 26 mg of 5.

Gileadenol (9): $C_{20}H_{32}O$. Oily; $[\alpha]^{20}D = +112^{\circ}$ (c 0.63, CHCl₃). IR (film) v_{max} : 3611 (OH), 1651 (C=C), 1212 cm⁻¹ (C-O). ¹H and ¹³C NMR: Table 1. HRESIMS: m/z 287.2363 (*Calcd.* 287.2375) [M⁺-1], 271.2414 (Calcd. 271.2426) [M⁺-OH].

2.3. Antimicrobial Activity

2.3.1. Bacterial Strains

The antimicrobial assays were performed using reference strains of the American Type Culture Collection (ATCC) available in the Microbiology Laboratory of College of Pharmacy/Prince Sattam University (Al-Kharj- Arabia Saudi). The standard microbial strains used were as follows: gram positive bacteria (*Bacillus subtilis* ATCC10400, *Staphylococcus aureus* ATCC35501), gram-negative bacteria (*Escherichia coli* ATCC25992, *Klebsiella pneumonia* ATCC138222) and *Candida albicans* ATCC14053.

2.3.2. Antimicrobial Assay

The obtained bacteria were sub-cultured onto Mueller-Hinton agar for 24 h and the fungal strain *C. albicans* was cultured on Sabouraud-dextrose agar for 5 days at 37 °C. Colonies from sub-cultured

Terpenes from the fresh stems of Commiphora gileadensis

plates were then grown in M_H broth and S-D broth to match the 0.5 McFarland turbidity standard equivalent to 1.5 $_{-}$ 10 8 CFU/mL. M_H agar and S-D agar were used for antibacterial and antifungal assays, respectively. The agar-well diffusion method was used for inhibition zone determination [16] using ampicillin as positive control. MIC were measured following the reported broth dilution method and the Clinical and Laboratory Standards Institute guidelines [17]. Serial dilutions were prepared and tested in from $7.812\mu g/mL$ to $50\mu g/mL$. Inoculum of 10 μL from each tested microbes were added to each tested dilution. Mueller-Hinton broth (MHB, Scharlau) alone was tested as sterility control, and untreated microbes inoculated on MHB alone and with different concentration of the solvent DMSO as negative control. All tubes were incubated at $37^{\circ}C$ for 24 h and the lowest concentration of antimicrobial agent that inhibited visible growth MIC was determined.

2.3.3 Data analysis

Values of three replicates are presented as mean \pm standard deviation. The statistical parameters applied were the Student's *t*-test. Differences were considered statistically significant at P < 0.05.




Figure 1. Chemical structures of compounds 1-9

3. Results and Discussion

3.1. Structure Elucidation

Compound 1 showed 20 carbon resonances in the 13 C NMR spectrum (Table S2) sorted by DEPT135 experiment into 4 X CH₃, 8 X CH₂, 4 X CH and 4 quaternary carbons. The chemical shifts of the carbon atoms pointed out to a hydrocarbon skeleton free from oxygenation. The molecular formula supported by HRESIMS $C_{20}H_{32}$ indicated 5 degree of unsaturation. The data of 1 (Tables S1, S2) were very similar to those reported for (1S,3E,7E,11R)-(+)-verticilla-3,7,12(18)-triene [18, 19].

¹³C NMR of **5** (Table S1, Figure S12) showed twenty carbon resonances sorted by DEPT135 experiment (Figure S13) into 5 X CH₃, 7 X CH₂, 4 X CH and 4 quaternary carbons. Similar to **1** the ¹H NMR of **5** (Table S2, Figure S11) showed two doublets for olefinic protons at δ_H 5.55, 4.80 correlated in an HSQC experiment (Figure S15) to carbons at δ_C 127.67 and 129.88 ppm. The main differences between **5** and **1** were the disappearance of the signals assigned for the exocyclic methylene moiety; instead one extra methyl singlet at δ_H 1.16; δ_C 24.04 ppm and an oxygenated quaternary carbon signal at δ_C 75.79 ppm were observed. HRESIMS data (Figure S16) showed ion peak at 273.2594 m/z representing M⁺-OH. Comparison of the obtained data of **5** with those reported for verticillol and its different isomers revealed a close similarity with *ent*-Verticillol [20-22].

Compounds 3 and 4 ¹³C NMR (Table S2, Figures S2, S7) showed two more oxygenated carbon signals at δ_C 63.60, 64.32 and 62.24, 66.37 ppm, respectively when compared with 5. The two oxygenated carbons in DEPT135 experiment (Figures S3, S8) were sorted into one CH and one quaternary carbon in 3 and 4. The HRESIMS showed an M+OH at m/z 289.2538 (Figure S5) and 289.2541 (Figure S10) for 3 and 4, respectively, supporting an oxirane ring in each compound. The ¹H and 13 C NMR spectrum (Figures S1, S2) of 3 revealed the disappearance of the olefinic signals at δ_H 5.55, δ_C 127.67 and the quaternary carbon at δ_C 132.90 ppm in 5, instead a proton signal at δ_H 3.45 (d, J=9.5 Hz) correlated in an HSQC experiment (Figure S4) to the oxygenated carbon signal at $\delta_{\rm C}$ 64.32 ppm and quaternary oxygenated carbon signal at $\delta_{\rm C}$ 63.60 ppm assigned for C13-C14 oxirane ring structure. The data of 3 were identical with those reported for (13S,14S)-Ent-13,14-epoxyverticillol [22]. The ¹H and ¹³C NMR spectrum (Figures S6, S7) of **4** revealed the disappearance of the olefinic signals at δ_H 4.64, δ_C 128.37 and the quaternary carbon at δ_C 133.90 ppm in 5 and they were replaced by a proton signal at $\delta_{\rm H}$ 2.95 (d, J= 9.9 Hz) correlated in an HSQC experiment (Figure S9) to the oxygenated carbon signal at δ_C 66.37 and quaternary oxygenated carbon signal at δ_C 62.24 ppm assigned for C9-C10 oxirane ring structure. The data of 4 were closely related with those reported for (9S,10S)-Ent-9,10epoxyverticillol [22]. This is the first report of the ent-verticillane-type diterpenes from the genus Commiphora.

Table 1. ¹H and ¹³C NMR data for compounds **9** (CDCl₃, δ in ppm, J in Hz).

Pos.	¹ H	¹³ C	Pos.	¹H	¹³ C
1	-	24.31	11	1.49 m ^a , 1.81 m ^a	26.34
2	0.63 m ^a	26.73	12	2.12 bq ^d (6.2)	53.01
3	0.40 dd ^b (9.8, 10.8)	29.22	13	-	153.43
4	1.26 m ^a	53.68	14	1.95 m ^a 2.33 dd ^b (6.0, 8.2)	38.87
5	-	131.08	15	0.92 m $^{\rm a}$, 1.87 m $^{\rm a}$	24.78
6	5.01 bt ^c (7.0)	124.86	16	0.95 s ^e	13.66
7	1.95 m ^a	25.26	17	1.59 s ^e	25.72
8	1.05 m a, 1.26 m a	43.29	18	1.52 s ^e	17.52
9	-	81.07	19	1.21 s ^e	25.90
10	1.48 m ^a , 1.68 m ^a	41.58	20	4.58 bs ^f , 4.61 bs ^f	106.30

^a multiplet, ^b doublet of doublet, ^c broad triplet, ^d broad quartet, ^e singlet, ^f broad singlet.

¹³C NMR of **9** (Table 1, Figure S18, S19) showed 20 carbon resonances sorted by DEPT135 experiment (Figure S20) into 4 X CH₃, 7 X CH₂, 5 X CH and 4 quaternary carbons. The HRESIMS data (Figure S33) showed M⁺-1 at m/z 287.2363 and M⁺-OH at m/z 271.2414 for the molecular formula C₂₀H₃₂O with 5 degrees of unsaturation. One of the degree of unsaturation was assigned from the NMR data (Figures S17-S20) of **9** as exo-methylene moiety at δ_H 4.59, 4.61 (each as broad singlet) correlated in HSQC experiment (Figure S23) with the carbon signal at δ_C 106.30 and the quaternary carbon at δ_C 153.43 ppm. Another degree of unsaturation represented by the olefinic signals at δ_H 5.01 (bt) correlated in an HSQC experiment with the CH at δ_C 124.86 and the quaternary carbon at δ_C 131.08 ppm. With the absence of other Sp^2 carbon signals the remaining unsaturation were accounted for three ring structures.

Terpenes from the fresh stems of Commiphora gileadensis




Figure 2. ¹H-¹H COSY (A), H2BC (B) and selected HMBC (C) correlations of compound 9.

The group of signals representing two CH, one quaternary carbon and two CH₃ groups at δ_H 0.63, δ_C 26.73 (C2); δ_H 0.40, δ_C 29.22 (C3); δ_C 24.31 (C1); δ_H 0.95, δ_C 13.66 (C16); δ_H 1.59, δ_C 25.72ppm (C17) were typical for 1,1-dimethyl cyclopropane moiety [23]. Combination of COSY and H2BC connectivity experiments (Figures 2, S21, S22, S25-S28) enable the identification of a CH-CH-CH spin system at $\delta_{\rm H}$ 0.40, $\delta_{\rm C}$ 29.22 (C3); $\delta_{\rm H}$ 1.26, $\delta_{\rm C}$ 53.68 (C4) and $\delta_{\rm H}$ 2.12, $\delta_{\rm C}$ 53.01ppm (C12). Another spin system was detected between CH at 0.63, δ_C 26.73 (C2), CH₂ at δ_H 0.92, 1.87, δ_C 24.78 (C15) and CH₂ at $\delta_{\rm H}$ 1.95, 2.33, $\delta_{\rm C}$ 38.87 ppm (C14). The terminal CH (C12) and CH₂ (C14) in these two spin systems showed two ways 3 bonds HMBC correlations (Figures 2, S29-S32) with the exo-methylene moiety at $\delta_{\rm H}$ 4.59, 4.61, $\delta_{\rm C}$ 106.30 (C20). H14 protons showed 2 bonds HMBC correlations with the quaternary carbon at δ_C 153.43 ppm (C13). This arrangement enables the assignment of a 7 membered ring fused with the 1,1-dimethyl cyclopropane ring structure. With the remaining two methyl groups signals at δ_H 1.52, δ_C 17.52 and δ_H 1.21, δ_C 25.90 ppm the rest of carbons must form a 9 membered ring coupled with the 7 membered ring via the two methines at δ_H 1.26, δ_C 53.68 (C-4) and δ_H 2.12, δ_C 53.01ppm (C12). The 9 membered ring contains the olefinic array and oxygenated quaternary carbon at δ_C 81.07 ppm. The positions of the two substituents was undoubtfully assigned based on COSY, H2BC and HMBC experiments results. The methyl signal at $\delta_{\rm H}$ 1.52 (C18) showed HMBC correlations with carbon signals at δ_C 131.08, 124.86 and 43.29 ppm assigned for C-5, C-6 and C-8, respectively. Clear H2BC were observed between the H-8 protons at δ_H 1.05, 1.26 and the C7 carbon at δ_C 25.26 ppm (Figure S26). The methyl group protons signal at δ_H 1.21 showed 2 bonds HMBC correlations with the carbons at δ_C 81.07 and 3 bonds correlations with the carbon at δ_C 41.58 ppm assigned for C-9 and C-10, respectively (Figure S31). COSY and H2BC correlations were observed between C10 (δ_H 1.48, 1.68; δ_C 41.58 ppm), C11 (δ_H 1.48, 1.81; δ_C 26.34 ppm) and C12 (δ_H 2.12; δ_C 53.01ppm) (Figures 2, S21, S22, S25-S28) confirming assignments of their position. In the HMBC experiment long range correlations were observed through 5 and 6 bonds away. This correlation can be explained by the "W" coupling [24] that could happen with trans fused 7 and 9 membered rings with α-oriented H-4. However, from the available data the complete stereochemistry cannot be determined.




Figure 3. Proposed stereo models of **9** to explain the long range HMBC correlations. **A**: Trans ring junction with α-oriented H-4 in favour of "W" coupling. **B**: Trans ring junction with β-oriented H-4 not in favour of "W" coupling.




Figure 4. Proposed biosynthetic origin of compound 9

The structure of **9** represents a rearranged verticillene skeleton identified for the first time from natural source. The name "gileadenol" was given to the new compound **9** with "gileadene" novel diterpene skeleton. The novel skeleton of **9** proposed to be biosynthesized from four isoprene unites in similar fashion as verticillene diterpenes. However, different cyclization pattern takes place to produce such newly discovered skeleton (Figure 4).

The four isolated triterpenes were identified as friedelin (2) [25, 26], oleanonic aldehyde (6) [27, 28], canyophyllal (7) [29] and urs-12-en-3-one-28-al (8) [30]. Compounds 6-8 are reported for the first time from *Commiphora* species.

Table 2. MIC (µg/mL) of different extracts of *C. gileadensis*.

	Staph. aureus	B. subtilis	E. coli	K. pneumonia	C. albicans
Fresh stem CHCl ₃ Ext.	750	750	750	250	125
Fresh stem MeOH Ext.	17000	35000	175	50000	16000
Fresh Leaves CHCl3 Ext.	750	750	750	250	125
Fresh Leaves MeOH Ext.	17000	25000	175000	25	7500
3	NT ^a	NT ^a	NT ^a	46.8	NT ^a
4	NT ^a	NT ^a	NT ^a	25	NT ^a
5	NT a	10	NT a	93.8	NT a

a Not tested.

3.2. Antimicrobial Activity

MIC were determined for the different extracts obtained from the fresh, dried leaves and stems (Table 2) using Gram-positive bacteria (Bacillus subtilis ATCC10400, Staphylococcus aureus ATCC35501), Gram-negative bacteria (Escherichia coli ATCC25992, Klebsiella pneumonia ATCC138222) and Candida albicans ATCC14053. All the chromatographic purification steps were monitored by the antimicrobial testing using E. coli and C. albicans and activity were determined based on the inhibition zone measurement. The CHCl₃ extract of the fresh stem was the most active and further purified to afford the nine isolates from the active fractions. The antimicrobial activity of compounds 2 and 6 was previously reported [26, 27]. Compounds 3-5, 7 and 9 were tested against the five organisms and the corresponding inhibition zones are presented in Table 3. Both 7 and 9 were less active against the tested microorganisms. MIC were determined for 3-5 against K. pneumonia and for 5 against B. subtilis as well (Table 2). Compound 4 was the most active with MIC= 0.025 mg/mL followed by 3 with MIC= 0.0468 mg/mL, while 5 was the least active at MIC= 0.0938 mg/mL against K. pneumonia. The MIC of 5 against B. subtilis was 0.01 mg/mL. The obtained results indicated that the compounds with verticillene skeleton were the most active. The presence of the epoxide moiety increased the activity against the Gram-negative bacteria K. pneumonia probably due to better penetration of the cell walls. Compound 5 lacking the epoxide function was more active against the Gram-positive bacteria B. subtilis. The antimicrobial activity of the isolated compounds supports the traditional uses of the plant to treat infected wounds [3, 4].

Terpenes from the fresh stems of Commiphora gileadensis

Table 3. Inhibition zones (mm) of compounds **3-5, 7** and **9** at 1 mg/mL against four bacterial strains

	Staph. aureus	B. subtilis	E. coli	K. pneumonia
Ampicillina	17±0.2	20±0.1	16±0.3	18±0.1
3	11 ± 0.3	16 ± 0.2	13 ± 0.3	18 ± 0.2
4	12 ± 0.2	16 ± 0.2	13 ± 0.3	20 ± 0.2
5	13±0.2	18 ± 0.3	12 ± 0.2	15 ± 0.1
7	11±0.3	15 ± 0.2	13 ± 0.3	15 ± 0.2
9	12 ± 0.4	13 ± 0.4	13 ± 0.2	16 ± 0.4

^a Ampicillin tested at 10 μg/mL

Acknowledgments

This Publication was supported by the Deanship of Scientific Research at Prince Sattam bin Abdulaziz University.

Supporting Information

Supporting information accompanies this paper on http://www.acgpubs.org/journal/records-of-natural-products

ORCID 🗓

Abdel-Kader M.S.: 0000-0002-5128-2656 Ibnouf E.O.: 0000-0001-9018-1391 Alqarni M.H.: 0000-0002-7061-9111 AlQutaym A.S.: 0000-0002-7128-1415 Salkini M. A.: 0000-0002-5244-2308

Ahmed I. Foudah: 0000-0002-5244-2308

References

- [1] S.A. Chaudhary (2013). Vegetation of the Kingdom of Saudi Arabia. Riyadh (Saudi Arabia) Ministry of Agriculture and Water.
- [2] H.A. El Rabey, A.I. Al-sieni, M.N. Al-seeni, M.A. Alsieni, A.I. Alalawy and F.M. Almutairi (2020). The antioxidant and antidiabetic activity of the Arabian balsam tree "Commiphora gileadensis" in hyperlipidaemic male rats, J. Taibah Univ. Sci. 14(1), 831-841.
- [3] A.G. Miller, M. Morris and S. Stuart-Smith (1988). Plants of Dhofar, the southern region of Oman: traditional, economic, and medicinal uses. Office of the Adviser for Conservation of the Environment, Diwan of Royal Court, Sultanate of Oman.
- [4] M.C. Marcotullio, O. Rosati and D. Lanari (2018). Phytochemistry of *Commiphora erythraea*: A review, *Nat. Prod. Comm.* **13(9)**, 1209–1212.
- [5] H.M. Al-mahbashi, A. El-shaibany and F.A. Saad (2015). Evaluation of acute toxicity and antimicrobial effects of the bark extract of Bisham (*Commiphora gileadensis* L.), *J. Chem. Pharm. Res.* **7** (6), 810-814.
- [6] O.M. Al Zoubi (2019). Evaluation of anti-microbial activity of ex vitro and callus extracts from *Commiphora gileadensis*, *Pak. J. Biol. Sci.* **22(2)**, 73–82.
- [7] D. Iluz, M. Hoffman, N. Gilboa-Garber and Z. Amar (2010). Medicinal properties of *Commiphora gileadensis*, *Afr. J. Pharmacy Pharmacol.* **4(8)**, 516–520.
- [8] L. Bouslama, B. Kouidhi, Y.M. Alqurashi and K. Chaieb A. Papetti (2019). Virucidal Effect of Guggulsterone Isolated from *Commiphora gileadensis*, *Planta Med.* **85**(16), 1225–1232.
- [9] T.A. Al-Howiriny, M. Al-Sohaibani, M. Al-Said, M. Al-Yahya, K. El-Tahir and S. Rafatullah (2005). Effect of *Commiphora opobalsamum* (L.) Engl. (Balessan) on experimental gastric ulcers and secretion in rats, *J. Ethnopharmacol.* **98**(3), 287–294.
- [10] A. Al-Howiriny, M. Al-Sohaibani, M. Al-Said, M. Al-Yahya, K. El-Tahir and S. Rafatullah (2004). Hepatoprotective properties of *Commiphora opobalsamum* ("Balessan"), a traditional medicinal plant of

- Saudi Arabia, *Drugs Exp. Clin. Res.* **30**(**5–6**), 213–220.
- [11] S.-L. Su, J. A. Duan, Y.-P. Tang, X. Zhang, L. Yu, F.-R. Jiang, W. Zhou, D. Luo and A.-W. Ding (2008). Isolation and Biological Activities of Neomyrrhaol and Other Terpenes from the Resin of *Commiphora myrrha*, *Planta Med.* **75(4)**, 351–355.
- [12] M.M. Rahman, M. Garvey, L.J.V. Piddock and S. Gibbons (2008). Antibacterial terpenes from the oleoresin of *Commiphora molmol* (Engl.), *Phytother. Res.* **22(10)**, 1356–1360.
- [13] P.G. Waterman and S. Ampofo (1985). Dammarane Triterpenes from the stem bark of *Commiphora Dalzielii*, *Phytochemistry* **24(12)**, 2925–2928.
- [14] T. Shen, W.-Z. Wan, X.-N. Wang, H.-Q. Yuan, M. Ji and H.-X.Lou (2009). A Triterpenoid and Sesquiterpenoids from the Resinous Exudates of *Commiphora myrrha*, *Helv. Chim. Acta.* **92**(4), 645–652.
- [15] A.A. El-Gamal, S.M. Al-Massarani, W.M. Abdel-Mageed, A. El-Shaibany, H.M. Al-Mahbashi, O.A. Basudan, F.A. Badria, M.S. Al-Said and M.S. Abdel-Kader (2017). Prenylated flavonoids from *Commiphora opobalsamum* stem bark, *Phytochemistry* **141**, 80–85.
- [16] J. Alkahtani, M. Soliman Elshikh, K.S. Almaary, S. Ali, Z. Imtiyaz and S. Bilal Ahmad (2020). Antibacterial, anti-scavenging and cytotoxic activity of garden cress polysaccharides, *Saudi J. Biol. Sci.* **27**(11), 2929–2935.
- [17] CLSI (2014). Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Fourth Informational Supplement. CLSI document M100-S24. Wayne, PA: Clinical and Laboratory Standards Institute.
- [18] J.D. Hernández- Hernández, L.U. Román Marín, M. Carlos, R. Cerda-García and P. Joseph-Nathan (2005). Verticillane Derivatives from *Bursera suntui* and *Bursera kerberi*, *J. Nat. Prod.* **68(11)**, 1598–1602
- [19] H.L. Greve, M. Kaiser, R. Brun and T.J. Schmidt (2017). Terpenoids from the Oleo-Gum-Resin of *Boswellia serrata* and Their Antiplasmodial Effects In Vitro. *Planta Med.* **83**(**14-15**), 1214–1226.
- [20] S. Bai and M. Jain (2008). ¹H and ¹³C assignments of five cembrenes from guggul, *Magn. Reson. Chem.* **46(8)**, 791–793.
- [21] F. Nagashima, A. Tamada, N. Fujii and Y. Asakawa (1997). Terpenoids from the Japanese liverworts *Jackiella javanica* and *Jungermannia infusca*, *Phytochemistry* **46**(7), 1203–1208.
- [22] F. Nagashima, K. Kishi, Y. Hamada, S. Takaoka and Y. Asakawa (2005). *ent*-Verticillane-type diterpenoids from the Japanese liverwort *Jackiella javanica*, *Phytochemistry* **66(14)**,1662-1670.
- [23] U. Warmers, K. Wihstutz, N. Bülow, C. Fricke and W.A. König (1998). Sesquiterpene constituents of the liverwort Calypogeia muelleriana, *Phytochemistry* **49**(6),1723-1731.
- [24] M.G. Constantino, V. Lacerda, G.V.J da Silva, L. Tasic and R. Rittner (2001). Principal component analysis of long-range 'W' coupling constants of some cyclic compounds, *J. Mol. Struct.* **597**, 129-136.
- [25] F.A. Abbas, S.M. Al-Massarany, S. Khan, T.A. Al-Howiriny, J.S. Mossa and E.A. Abourashed (2007). Phytochemical and biological studies on Saudi *Commiphora opobalsamum* L., *Nat. Prod. Res.* **21**(5), 383–391.
- [26] N.S.S. Ambarwati, B. Elya, A. Malik, M. Hanafi and H. Omar (2019). Isolation, characterization, and antibacterial assay of friedelin from *Garcinia latissima* Miq. leaves, *J. Phys. Conf. Ser.* **1402(5)** 055078.
- [27] L. Karygianni, M. Cecere, A. Argyropoulou, E. Hellwig, A.L. Skaltsounis, A. Wittmer, J.P. Tchorz and A. Al-Ahmad (2019). Compounds from *Olea europaea* and *Pistacia lentiscus* inhibit oral microbial growth, *BMC Complement. Altern. Med.* **19**(1), 51.
- [28] M. Ruhul Kuddus, F. Rumi, M. Abul Kaisar and C. Mahmood Hasan (2011). Secondary Metabolites from *Melocanna baccifera* (Roxb.), *Asian J. Chem.* **23**(1), 85–88.
- [29] Y.Z. Li, Z.L. Li, S.L. Yin, G. Shi, M.S. Liu, Y.K. Jing and H.M. Hua (2010). Triterpenoids from *Calophyllum inophyllum* and their growth inhibitory effects on human leukemia HL-60 cells, *Fitoterapia* **81(6)**, 586-589.
- [30] I. Wahlberg, K. Karlsson, M.B. Hjelte and C.R. Enzell. (1971). Constituents of commercial tolu balsam, *Acta Chem. Scand.* **25**, 3285–3295.

