








A New Lignan from Leaves of *Ormosia xylocarpa*

Wenjuan Zhou ¹, Yingxuan Quan ¹, Yuanan Chen ¹,
Qin Wang ², Xiaoxing Zou ², Fangyou Chen ³ and Lin Ni ^{1,2*}

¹Key Laboratory of Biopesticide and Chemical Biology, Ministry of Education, Fujian Agriculture and Forestry University, Fuzhou, Fujian 350002, China

²Engineering Research Center of Natural Biological Resources Conservation & Utilization of Fujian Province, Fujian Agriculture and Forestry University, Fuzhou, Fujian 350002, China

³College of Pharmacy, Jiangxi University of Traditional Chinese Medicine, Nanchang, Jiangxi 330004, China

(Received March 14, 2022; Revised June 17, 2022; Accepted June 18, 2022)

Abstract: A new lignan, 4,4''-dihydroxy-3,3',5',3'',5'',7''-pentamethoxy-7,9';7',9'-diepoxy-4',8''-oxy-8,8'-sesqueneo-lignan-propanol (**1**), along with six known lignans (**2-7**) was isolated from the leaves of *Ormosia xylocarpa* (Chun ex L. Chen). The structure of compound **1** was elucidated through comprehensive 1D and 2D NMR, UV, IR, and HRMS analyses. Compounds **2-3** performed strong antioxidant activity, the median clearance concentration of DPPH, ABST⁺, and ·OH were lower than 40 μM.

Keywords: *Ormosia xylocarpa*; lignans; chemical constituents; antioxidant activity. © 2022 ACG Publications. All rights reserved.

1. Plant Source

The leaves of *Ormosia xylocarpa* (Chun ex L. Chen) were collected from the 15-year-old tree in September 2019 in Shaxian County, Fujian Province, China (117°78' N latitude, 26°40' E longitude), and were identified by one of the authors (Xiaoxing Zou). The voucher specimen (accession number: 20190812) was preserved in the Engineering Research Center of Natural Biological Resources Conservation & Utilization of Fujian Province, FAFU, Fuzhou, China.

2. Previous Studies

Ormosia xylocarpa is a valuable timber tree species belonging to the genus *Ormosia*, widely distributed in southern China [1]. Its huge canopy produces a lot of fallen leaves, which possess rich

*Corresponding author: E-mail: nilin_fjau@126.com

A new lignan from *Ormosia xylocarpa*

medicinal value and can be used for treating eye diseases in the folk [2]. The ancient method of decoction with its branches and leaves to prevent cardiovascular diseases indicated its potential antioxidant ability [3]. Some lignans have been isolated from *O. xylocarpa*, including xylocarpalignan, syringaresino-4'-O- β -D-glucoside, and zhebeir esinol [4-5]. In our research, seven lignans were isolated and identified (Figure 1), one of them was a new compound. They all showed certain antioxidant activity in the DPPH, ABST⁺, and \cdot OH radical scavenging activity assays. All compounds were first found in the *Ormosia* genus.

3. Present Study

The leaves of *O. xylocarpa* (10.0 kg) were powdered and then extracted with 70% ethanol (2 \times 20 L, 2 \times 2 h) to give an extract (1.86 kg). The extract was extracted with EtOAc (2 \times 45 L) to obtain the EtOAc fraction. The EtOAc fraction (407 g) was chromatographed over polyamide macroporous (60-100 mesh) with EtOH-H₂O gradient (H₂O, 3:7, 1:1, 6:4, 7:3, EtOH) as eluent to obtain 18 fractions (Fr.1~Fr.18). Fr.1 (87.3 g) was chromatographed over silica gel with CH₂Cl₂-MeOH as eluent to obtain 10 fractions (Fr.1.1~Fr.1.10). Compound **1** (3.2 mg) was isolated from Fr.1.1 (3.1 g) by HPLC (CH₃CN/H₂O (28: 72), 8.0 mL/min) after separating by Sephadex LH-20 with MeOH. Fr.1.3 (8.1g) was separated by PRP-512A and eluted with EtOH-water (1:9, 3:7, 1:1, 7:3, 9:1, EtOH) to give 6 subfractions (Fr.1.3.1~Fr.1.3.6). Fr.1.3.2 (640 mg) was further separated by Sephadex LH-20 with MeOH to obtain 5 subfractions (Fr.1.3.1.1~Fr.1.3.6.5). Compound **2** (5.3 mg) and **4** (3.4 mg) were separated from Fr.1.3.1.2 (168 mg) by semi-preparative HPLC (MeOH/H₂O (47: 53), 8.0 mL/min). Fr.1.3.1.3 (180 mg) was subjected to semi-preparative HPLC (CH₃CN/H₂O (28: 72), 8.0 mL/min) to obtain compound **3** (7.2 mg). Fr.1.3.3 (700 mg) was chromatographed over Sephadex LH-20 eluted with MeOH and further purified by HPLC (MeOH/H₂O (30: 70), 8.0 mL/min) to yield compound **5** (2.8 mg), **6** (3.7 mg) and **7** (3.2 mg).





Figure 1. The structure of compounds **1-7**. * indicates new compound

Compound **1**: Yellow oil; UV (MeOH) λ_{max} : 280, 250, 230 nm, IR (KBr) λ_{max} 3413.0, 1611.0, 1461.4, 1224.8, 1026.4 cm⁻¹. CD (MeOH) λ_{max} ($\Delta\epsilon$) 247 (+6.73), 278 (+9.48); HR-ESI-MS m/z 651.2409 [M + Na]⁺ (calcd for C₃₃H₄₀O₁₂Na, 651.2412). ¹H-NMR (DMSO-*d*₆, 400 MHz) and ¹³C-NMR (DMSO-*d*₆, 100 MHz) data see Table 1.

Table 1. ^1H -NMR (400MHz, $\text{DMSO-}d_6$) and ^{13}C -NMR (100 MHz, $\text{DMSO-}d_6$) data of compound **1**

Position	δ_{H}	δ_{C}
1		132.2
2	6.88 (1H, d, $J = 2.0$ Hz)	115.2
3		147.7
4		146.0
5	6.72 (1H, d, $J = 8.0$ Hz)	110.4
6	6.75 (1H, dd, $J = 8.0, 2.0$ Hz)	118.7
7	4.61 (1H, d, $J = 3.6$ Hz)	85.2
8	3.05 (1H, m)	53.5
9	4.14 (2H, m)	71.3
1'		136.8
2', 6'	6.61 (2H, d, $J = 2.0$ Hz)	103.2
3', 5'		152.6
4'		134.7
7'	4.64 (1H, d, $J = 3.6$ Hz)	85.2
8'	3.02 (1H, m)	53.8
9'	3.77 (overlapped)	71.0
1''		126.7
2'', 6''	6.54 (2H, m)	104.9
3'', 5''		147.6
4''		134.7
7''	4.39 (1H, d, $J = 6.8$ Hz)	82.6
8''	4.19 (1H, m)	84.9
9''	3.65 (1H, overlapped)	59.8
	3.47 (1H, overlapped)	
3-OCH ₃	3.76 (3H, s)	55.6
3', 5'-OCH ₃	3.72 (6H, s)	56.0
3'', 5''-OCH ₃	3.70 (6H, s)	55.9
7''-OCH ₃	3.71 (3H, s)	56.6

**Figure 2.** Key HMBC and ^1H - ^1H COSY Correlations of Compound **1**

Compound **1** was isolated as a yellow oil, and its UV spectrum showed three absorption peaks at 280, 250 and 230 nm. Its molecular formula was defined as $\text{C}_{33}\text{H}_{40}\text{O}_{12}$, based on analysis of the

A new lignan from *Ormosia xylocarpa*

HRESIMS (m/z 651.2409 $[M + Na]^+$, calcd. 651.2412). In the IR spectrum, absorption bands at 3413 cm^{-1} (hydroxy), 1611.0 and 1461.4 cm^{-1} (aromatic ring) were observed.

The ^1H NMR spectrum of **1** showed hydrogen proton signals of a benzene ring ABX system [δ_{H} 6.88 (1H, d, $J = 2.0$ Hz), 6.75 (1H, dd, $J = 8.0, 2.0$ Hz), and 6.72 (1H, d, $J = 8.0$ Hz)], two sets of symmetrical aromatic hydrocarbon proton signals [δ_{H} 6.61 (2H, m) and 6.54 (2H, m)], four oxygenated methines [δ_{H} 4.64 (1H, d, $J = 3.6$ Hz), 4.61 (1H, d, $J = 3.6$ Hz), 4.39 (1H, d, $J = 6.8$ Hz), and 4.19 (1H, m)], and six methoxy groups [δ_{H} 3.76 (3H, s), 3.76 (6H, s), and 3.72 (9H, s)]. The ^{13}C NMR associated with the DEPT spectra of **1** classified 33 carbon resonances indicated the existence of a 3,4-distributed phenyl and two symmetric 3',4',5'-trisubstituted phenyl, three methylene groups, and two methines. These data manifested compound **1** as a sesquieolignan [6-7].

In the ^1H - ^1H COSY spectrum, the correlations of H-7/H-8/H-9 [δ_{H} 4.61 (1H, d, $J = 3.6$ Hz), 3.05 (1H, m), 4.14 (2H, m)] and H-7'/H-8'/H-9' [δ_{H} 4.64 (1H, d, $J = 3.6$ Hz), 3.02 (1H, m), and 3.77 (overlapped)] suggested the existence of a 7,9';7',9-diepoxy moiety [8]. The HMBC correlations from δ_{H} 3.72 (6H, s, 3', 5'-OCH₃) to C-3' (δ_{C} 152.6), C-2' (δ_{C} 103.2), C-5' (δ_{C} 152.6), and C-6' (δ_{C} 103.2), from δ_{H} 3.76 (3H, s, 3-OCH₃) to C-2 (δ_{C} 115.2) and C-3 (δ_{C} 147.7) indicated the location of MeO-3, 3', 5'. The ^1H - ^1H COSY correlations of H-5/H-6 [δ_{H} 6.72 (1H, d, $J = 8.0$ Hz), 6.75 (1H, dd, $J = 8.0, 1.6$ Hz)], and the HMBC correlations from H-6 [δ_{H} 6.75 (1H, dd, $J = 8.0, 1.6$ Hz)] to C-2 (δ_{C} 115.2), from H-2 [δ_{H} 6.88 (1H, d, $J = 2.0$ Hz)] to C-7 (δ_{C} 85.2), from H-7 [δ_{H} 4.61 (1H, d, $J = 3.6$ Hz)] to C-1 (δ_{C} 132.2) and C-6 (δ_{C} 118.7), from H-6' [δ_{H} 6.61 (2H, dd, $J = 2.0$ Hz)] to C-1' (δ_{C} 136.8), C-5' (δ_{C} 152.6), and C-4' (δ_{C} 134.7), from H-7' [δ_{H} 4.64 (1H, d, $J = 3.6$ Hz)] to C-1' (δ_{C} 136.8) indicated that the compound **1** has a 4-hydroxy-3,4',5',-trimethoxy-7,9';7',9-diepoxy lignan structural unit. The ^1H - ^1H COSY spectrum indicated a glycerol-type moiety at H-7''/H-8''/H-9'' [δ_{H} 4.39 (1H, d, $J = 6.8$ Hz), 4.19 (1H, m), 3.65 (1H, overlapped) and 3.47 (1H, overlapped)] [9]. In addition, the HMBC correlations from H-2'' [δ_{H} 6.54 (2H, m)] to C-1'' (δ_{C} 126.7), C-3'' (δ_{C} 147.6), and C-4'' (δ_{C} 134.7), from H-7'' [δ_{H} 4.39 (1H, d, $J = 6.8$ Hz)] to C-1'' (δ_{C} 126.7), from δ_{H} 3.70 (6H, s, 3'', 5''-OCH₃) to C-2'' (δ_{C} 104.9), C-3'' (δ_{C} 147.6), C-5'' (δ_{C} 147.6), and C-6'' (δ_{C} 104.9) indicated that the compound **1** has a 3'',5''-dimethyl-4-hydroxy-phenylpropanol structural unit. The HMBC correlations from H-7'' [δ_{H} 4.39 (1H, d, $J = 6.8$ Hz)] to MeO-7'' (δ_{C} 56.6) indicated that MeO-7'' was located at C-7''. Moreover, the HMBC spectrum confirmed the 4'-8''-oxy linkage between the two units at H-8'' [δ_{H} 4.19 (1H, m)] to C-4' (δ_{C} 134.7) [10-11].

According to H-7' [δ_{H} 4.64 (1H, d, $J = 3.6$ Hz)] and H-7 [4.61 (1H, d, $J = 3.6$ Hz)], the coupling constant of 3.6 Hz between H-7/H-8 and H-7'/H-8' confirmed an *erythro* relative configuration [12]. The coupling of 6.8 Hz between H-7'' and H-8'' defined a *threo* relative configuration [13]. The spectrum showed a positive cotton effect at 247 nm and 278 nm agreeing with the configuration of **1** to be 7*S*, 7'*S*, 7''*S*, 8*R*, 8'*R*, and 8''*S* [14-15]. Compound **1** was given a trivial name xylocarpalignan B, and its structure was determined and illustrated in Figure 2.

Six known lignans (**2-7**) isolated from *O. xylocarpa* were identified as hedyotol C (**2**) [16], buddlenol C (**3**) [17], (+)-medioresinol (**4**) [18], (+)-isolariciresinol (**5**) [19], 5-methoxy-(+)-isolariciresinol (**6**) [20], and (+)-lyoniresinol (**7**) [21], by comparison with the published NMR data.

Table 2. Antioxidant activity values of Compounds **1-7**

Compounds	DPPH (EC ₅₀ , μM)	ABTS ⁺ (EC ₅₀ , μM)	$\cdot\text{OH}$ (EC ₅₀ , μM)
VC	48.79	5.03	39.10
1*	26.61	18.01	28.44
2	15.43	7.63	17.43
3	29.38	13.30	32.54
4	38.33	18.29	37.26
5	121.87	11.45	28.36
6	173.71	12.23	22.31
7	101.76	5.07	12.78

The DPPH, ABTS⁺ and ·OH radical scavenging activity assays [22-24] were conducted to determine the antioxidant ability of compounds **1-7**. The results showed that compounds **2-3** and **5-7** exerted strong antioxidant activity (Table 2). Compounds **2-3** are 8.O.4'-neolignans and their antioxidant properties were related to the conformation of C-7'' and C-8'' [25], and 8.O.4'-neolignans with the *threo* series possessed stronger activity. The lignans with a phenolic hydroxyl (compounds **5-7**) showed significant antioxidant activity like phenolic compounds.

Acknowledgments

This work was supported by the Outstanding-Young Scientific Research Talents Program of Fujian Agriculture and Forestry University (xjq202103), Open Project of the Key Laboratory of Biological Pesticide and Chemical Biology of the Ministry of Education (No. KeyLab2021-01), and National Agricultural Education Advisory Committee Research Project (No:2021-NYYB-12).

Supporting Information

Supporting Information accompanies this paper on <http://www.acgpubs.org/journal/records-of-natural-products>

ORCID

Wenjuan Zhou: [0000-0003-2498-940X](https://orcid.org/0000-0003-2498-940X)

Yingxuan Quan: [0000-0001-5683-8003](https://orcid.org/0000-0001-5683-8003)

Yuanan Chen: [0000-0002-5684-9340](https://orcid.org/0000-0002-5684-9340)

Qin Wang: [0000-0001-6528-3329](https://orcid.org/0000-0001-6528-3329)

Xiaoxing Zou: [0000-0002-3364-6719](https://orcid.org/0000-0002-3364-6719)

Fangyou Chen: [0000-0002-4650-3987](https://orcid.org/0000-0002-4650-3987)

Lin Ni: [0000-0001-6118-6724](https://orcid.org/0000-0001-6118-6724)

References

- [1] S. X. Su (2011). Growth and Biomass of a *Ormosia xylocarpa* plantation in Sanming, Fujian, *J. Fujian Nor. Univ.* 27, 148-152.
- [2] D. S. Di, H. Z. Liao, C. N. Zhang and Z. Z. Chen (1991). A study on ecosystem productivity and tree growing rules of *Ormosia xylocarpa* plantation stand, *J. Nanjing For. Univ.* 15, 60-65.
- [3] S. Zhu (2020). Research on seedling raising techniques of *Ormosia xylocarpa*, *Anhui Agri. Sci. Bull.* 26, 65-66.
- [4] Q. Wang, X. Chen, S. Q. Ruan, W. H. Yan, L. Ni, S. Q. Zou and X. X. Zou (2022). A new lignan from leaves of *Ormosia xylocarpa*, *Chin. Herb. Med.* 53, 653-658.
- [5] Q. Wang, X. Chen, X. X. Zou, S. Q. Zou, L. Ni and H. Y. Xu (2022). A new flavonoid from leaves of *Ormosia xylocarpa*, *Nat. Prod. Commun.* 17, 1-5.
- [6] F. Cutillo, M. Dellagrecia, M. Gionti, L. Previtera and A. Zarrelli (2006). Phenols and lignans from *Chenopodium album*, *Phytochem Anal.* 17, 344-349.
- [7] M. Miyazawa, H. Kasahara and H. Kameoka (1993). Biotransformation of lignans: a specific microbial oxidation of (+)-eudesmin and (+)-mahanolin by *Aspergillus niger*, *Phytochemistry* 34, 1501-1507.
- [8] S. Y. Shao, Y. N. Yang, Z. M. Feng, J. S. Jiang and P. C. Zhang (2018). An Efficient method for determining the relative configuration of furofuran lignans by ¹H NMR spectroscopy, *J. Nat. Prod.* 81, 1023-1028.
- [9] Y. Y. Lu, Y. B. Xue, J. J. Liu, G. M. Yao, D. Y. Li, B. Sun, J. W. Zhang, Y. F. Liu, C. X. Chang, M. Xiang and et al (2015). (+/-)-Acortatarinowins A-F, norlignan, neolignan, and lignan enantiomers from *Acorus tatarinowii*, *J. Nat. Prod.* 78, 2205-2214.
- [10] L. Xiong, C. G. Zhu, Y. R. Lu, Y. Tian, S. Lin, S. P. Yuan, J. F. Hu, Q. Hou, N. C. Chen, N. H. Chen, Y. C. Yang and et al (2011). Lignans and neolignans from *Sinocalamus affinis* and their absolute

A new lignan from *Ormosia xylocarpa*

configurations, *J. Nat. Prod.* 74, 1188-1200.

- [11] X. W. Yang, P. J. Zhao, Y. L. Ma, H. T. Xiao, Y. Q. Zuo, H. P. He, L. Li and X. J. Hao (2007). Mixed lignan-neolignans from *Tarenna attenuata*, *J. Nat. Prod.* 70, 521-525.
- [12] Z. Yuan and X. Li (2003). NMR Methods for determining the configuration of 8-*O*-4' neolignans, *J. Magn. Reson.* 20, 307-314.
- [13] N. Hao, L. R. Han, Y. T. Li, J. Li, X. L. Tian, D. Kong and X. Tian (2020). New 8-*O*-4' Neolignans and their antibacterial activity from the whole plants of *Clematis lasiandra*, *ACS Omega* 5, 19661-19666.
- [14] X. Chen, Q. Ren, Y. Liu, Z. X. Zou, K. P. Xu and G. S. Tan (2020). Circular dichroism in the determination of absolute configuration of lignans and neolignans, *Cent. South Pharm.* 18, 1-10.
- [15] P. Fang, X. Liang, Y. L. He, L. Fei and P. Cheng (2021). 8-*O*-4' Neolignans from the fruits of *Leonurus japonicus*, *Nat. Prod. Res.* 33, 1320-1325.
- [16] J. H. Liu (2014). Chemical constituents from the fruits of *Capparis masaikai*, *Nat. Prod. Res.* 26, 1780-1784.
- [17] X. Z. Liu, X. Lu, S. L. He, W. J. Tian, T. Lin and H. F. Chen (2020). Lignans from *Vernonia amygdalina* Del., *Chin. J. Med. Chem.* 12, 738-744.
- [18] L. Y. Y. J. Xu and D. H. Xu (2018). Lignanoids from *Solanum lyratum*, *J. Chin. Pharm. Sci.* 27, 289-294.
- [19] X. B. Yang, S. X. Sheng, M. Y. Li, J. Y. Pan, Y. N. Zheng and M. Y. Li (2010). Chemical constituents in seeds of Indian mangrove *Xylocarpus granatum*, *Chin. Tradit. Herb. Drugs.* 41, 846-851.
- [20] C. S. Zou, M. M. Yuan, X. Wu, S. Zeng, Y. L. Ou and G. P. Zhou (2020). Chemical constituents from n-butanol extract of *Cynachum paniculatum*, *J. Chin. Med. Mater.* 43, 606-611.
- [21] X. Cao, C. J. Li, J. Z. Yang, B. X. Wei, Y. M. Luo and D. M. Zhang (2011). Study on chemical constituents from leaves of *Tripterygium wilfordii*, *Chin. J. Chin. Mater. Med.* 36, 1028-1031.
- [22] W. Li, J. R. Fu, L. J. Zheng, L. Ni, J. Q. Liu, J. W. Zhai, Z. Zhou and S. S. Wu (2022). Two new bibenzyls from *Pleione grandiflora* (Rolfe) Rolfe and their antioxidant activity, *Nat. Prod. Res.* 15, 1-7.
- [23] R. Wang, Q. Luo and Y. Feng (2018). Determination of antioxidant effects of kaempferol by micro-model of DPPH, ABTS and FRAP assay, *Guangdong Chem. Ind.* 43, 58-63.
- [24] P. Shu, Y. Li, Y. Luo, S. Cai, Y. Fei, W. Yi, Y. Yang, X. Wei, Y. Zhang, T. Tu and L. Zhang (2021). Isolation, characterization and antioxidant, tyrosinase inhibitory activities of constituents from the flowers of *Cercis glabra* 'Spring-1', *Rec. Nat. Prod.* 25, 254-260.
- [25] K. Kónyal, Zs. Varga and S. Antus (2001). Antioxidant properties of 8-*O*.4'-neolignans, *Phytomedicine* 8, 454-459.

A C G
publications

© 2022 ACG Publications