

Sesquiterpenoids from the Leaves of *Dalbergia odorifera*

Chenxiao Shen ¹, Can Yue Ouyang ², Ying Zhang ², Qing Zhu ³,
Ronghua Liu ^{3*} and Lanying Chen ^{2*}

¹ Key Laboratory of Modern Preparation of TCM, Ministry of Education, Jiangxi University of Chinese Medicine, Nanchang, 330004, China

² National Pharmaceutical Engineering Center for Solid Preparation of Chinese Herbal Medicine, Jiangxi University of Chinese Medicine, Nanchang, 330006, China

³ School of Pharmacy, Jiangxi University of Chinese Medicine, Nanchang, 330006, China

(Received May 27, 2022; Revised July 19, 2022; Accepted July 20, 2022)

Abstract: Two new sesquiterpenoids (1*R*,5*R*,6*R*)-11-hydroxyl-drocostuslactone (**1**) and (1*R*,10*S*)- gibberodione A (**2**) were acquired from the leaves of *Dalbergia odorifera*. The structures of the new compounds were elucidated by 1D and 2D NMR techniques, and X-ray crystallography analyses. The results of the bioactivity screening tests of the two new compounds revealed the potential anti-inflammatory effects of these two compounds, their IC₅₀ values were 107.2 ± 4.02 and 54.64 ± 1.89 µg/mL, respectively. They could significantly decrease the production of NO (P<0.01) with 8~16 and 1~2 µg/mL and inhibited LDH (P<0.01) with 1~16 and 1~2 µg/mL in lipopolysaccharide (LPS)-stimulated RAW 264.7 macrophages, respectively.

Keywords: *Dalbergia odorifera* leaves; sesquiterpenoids; anti-inflammatory activity. © 2022 ACG Publications. All rights reserved.

1. Introduction

Dalbergia odorifera T. Chen, which belongs to the family Fabaceae, is a vital TCM. Its heartwood was utilized in China and Korea to treat blood stasis, ischemic symptoms, swelling, necrotic and rheumatic pains [1-2]. Phytochemical examination the heartwood of *D. odorifera* has identified substantial chemical substances, which involve neoflavonoids, flavonoids, furans, benzophenones and terpenoids [3-8]. The leaves of *D. odorifera* exhibited anti-inflammatory, antibacterial and antioxidant activities [9-10]. We studied the chemistry constituents of *D. odorifera*

* Corresponding author: E-Mail: rhliu@163.com (R. Liu); (C. Shen); cly2513@163.com (L.Chen)

leaves.

Two new sesquiterpenoids were acquired from the leaves of *D. odorifera* (Figure 1). In addition, the bioactivity assay showed that two compounds had potential anti-inflammatory activity.




Figure 1. Structural features of chemical substances **1** and **2**

2. Materials and Methods

2.1. General Experimental Procedures

The nuclear magnetic resonance (NMR) spectra of the isolated compounds were acquired on a 400 MHz NMR spectrometer (Bruker Corporation, Switzerland). Ultraviolet (UV) patterns of the compounds were determined on a 210A double beam spectrophotometer (Shimadzu, Japan). High resolution mass spectrometry (HRMS) data were acquired on an LCMS-IT-TOF mass spectrometer (Shimadzu, Kyoto, Japan). Circular dichroism (CD) spectra were recorded using a JASCOJ-1500 spectropolarimeter (CA, USA). Optical rotations were measured using a JASCO P-1020 polarimeter (JASCO Corporation, Tokyo, Japan).

2.2. Plant Material

The leaves of *Dalbergia odorifera* were harvested from Shanya, Hainan Province in China, in July 2020, which was identified by associate researcher Zhou Hong. A voucher specimen (No20200704.) was preserved in Key Laboratory of Modern Preparation of TCM, Ministry of Education, Jiangxi University of Chinese Medicine.

2.2. Extraction and Isolation

The leaves of *Dalbergia odorifera* (24.0 kg) were isolated under reflux by 70% ethyl alcohol thrice. Ethyl alcohol was evaporated using a rotary evaporator at reduced pressure and 4.50 kg crude ethyl alcohol extract was obtained. The crude extract was subjected to D101 resin column using H₂O, EtOH-H₂O (15:85-95:5) as eluents to yield 7 fractions (Fr.1-Fr.7). The fraction 4 (282.8 g) was furtherly separated on silica gel column chromatography (CC) (CH₂Cl₂-MeOH, 500:1-0:1) to yield twelve fractions (Frs.4.A-Frs.4.L), and the Frs.4.C (6.9 g) was subjected to silica gel CC (petrol

Sesquiterpenoids from the leaves of *Dalbergia odorifera*

ether-acetone, 200:1-1:1) to produce 3 fractions (Frs.4.C.1-Frs.4.C.3). The Frs.4.C.1 (301 mg) was subjected to Sephadex LH-20 CC with CH₂Cl₂-MeOH (1:1, v/v) solvent system and compound **1** (15 mg) was obtained. The Frs.4.D (2.8 g) was subjected to silica gel CC with (petrol ether-acetone, 200:1-1:1) solvent system and yielded to as 5 fractions as (Frs.4.D.1-Frs.4.D.5). By further separation of one of those fractions, Frs.4.D.3 (183.2 mg), on a silica gel column with CH₂Cl₂-MeOH (from 150:1-0:1, v/v) solvent system, compound **2** (8 mg) was isolated.

Table 1. The NMR data for **1** and **2** in CDCl₃ (δ in ppm, J in Hz)

No.	1		2	
	δ_{H} , mult. (J in Hz)	δ_{C}	δ_{H} , mult. (J in Hz)	δ_{C}
1	2.89 (1H, m)	48.9	2.31 (1H, m)	58.1
2	2.44 (2H, m)	30.7	1.94 (1H, m) 1.83 (1H, m)	24.0
3	2.56 (1H, m) 2.16 (1H, m)	31.1	2.49 (1H, ddd, $J = 17.4, 9.0, 5.2$ Hz) 2.28 (1H, m, H-11)	42.0
4	-	149.5	-	208.9
5	2.60 (1H, d, $J = 12.6$ Hz)	51.8	-	205.7
6	4.61 (1H, d, $J = 11.1$ Hz)	79.4	5.84 (1H, d, $J = 1.9$ Hz)	126.5
7	-	136.3	-	167.5
8	2.97 (1H, m) 2.35 (1H, m)	27.3	2.40 (1H, m) 2.25 (1H, m)	28.4
9	2.00 (1H, m) 1.81 (1H, m)	29.0	1.68 (1H, m) 1.64 (1H, m)	34.5
10	-	149.2	1.74 (1H, m)	34.8
11	-	137.1	2.35 (1H, m)	37.4
12	-	170.5	1.06 (3H, d, $J = 2.6$ Hz)	21.0
13	4.94 (1H, s) 4.91 (1H, s)	112.5	1.08 (3H, d, $J = 2.6$ Hz)	21.3
14	5.15 (1H, s) 5.10 (1H, s)	113.2	1.10 (3H, d, $J = 6.5$ Hz)	20.1
15	-	-	2.11 (3H, s)	30.1

(*1R,5R,6R*)-11-hydroxyl-drocostuslactone (**1**): Colorless crystal (MeOH); $[\alpha]_{\text{D}}^{20} = + 56.21$ ($c = 0.58$, MeOH). UV (MeOH) λ_{max} : 240 nm, IR (KBr) ν_{max} : 3417, 1944, 1901, 1832, 1637, 1031 cm⁻¹. HR-ESI-MS m/z 233.1162 ($[\text{M}+\text{H}]^+$ calcd. for C₁₄H₁₇O₃, 233.1172). ¹H-NMR (CDCl₃, 400 MHz) and ¹³C-NMR (CDCl₃, 100 MHz): see Table 1.

(*1R,10S*)-gibberodione A (**2**): Colorless oil (MeOH); $[\alpha]_{\text{D}}^{20} = - 3.64$ ($c = 0.07$, MeOH). UV (MeOH) λ_{max} : 240 nm, IR (KBr) ν_{max} : 2963, 2936, 1715, 1664 cm⁻¹. HR-ESI-MS m/z 237.1835 ($[\text{M}+\text{H}]^+$ calcd. for C₁₅H₂₅O₂, 237.1849). ¹H-NMR (CDCl₃, 400 MHz) and ¹³C-NMR (CDCl₃, 100 MHz): see Table 1.

3. Results and Discussion

3.1. Structure Elucidation

Compound **1** was isolated as achromatic crystal, with a molecular formula of $C_{14}H_{16}O_3$ was deduced from the HR-ESI-MS peak at m/z 233.1162, in correspondence to seven levels of under saturation. The 1H NMR spectral data of **1** displayed resonances for two olefinic protons [δ_H 5.15 (1H, s, H-14 α), 5.10 (1H, s, H-14 β), 4.94 (1H, s, H-13 α), 4.91 (1H, s, H-13 β)], three methines groups [δ_H 4.61 (1H, d, $J = 11.1$ Hz, H-6), 2.89 (1H, m, H-1), 2.60 (1H, d, $J = 12.6$ Hz, H-5)], one oxidative methine [δ_H 4.61 (1H, d, $J = 11.1$ Hz, H-6)], four methylenes [δ_H 2.97 (1H, m, H-8 α), 2.56 (1H, m, H-3 α), 2.44 (2H, m, H-2), 2.35 (1H, m, H-8 β), 2.16 (1H, m, H-3 β), 2.00 (1H, m, H-9 α), 1.81 (1H, m, H-9 β)]. The ^{13}C NMR spectra of **1** displayed 14 carbon resonances involving an ester carbonyl carbon [δ_C 170.5 (C-12)], six olefinic carbons [δ_C 149.5 (C-4), 149.2 (C-10), 137.1 (C-11), 136.3 (C-7), 113.2 (C-14), 112.5 (C-13)], one oxygenated methine [δ_C 79.4 (C-6)], two methines groups [δ_C 51.8 (C-5), 48.9 (C-1)], three methylenes [δ_C 31.1 (C-3), 30.7 (C-2), 29.0 (C-9), 27.3 (C-8)]. The data are closely comparable to those of dehydrocostuslactone [10]. The 1H - 1H COSY correlation of H-14/H₂-3, H₂-3/H₂-2, H₂-2/H-1, H-13/H₂-9, H₂-9/H₂-8 showed the sequence =CH (1) –CH₂(2) –CH₂(3)/CH₂(8) –CH₂(9). The HMBC association of H-1 with C-13/C-10 /C-2/C-5/C-6, H-13 with C-1/C-10/C-9, H-14 with C-4/C-3/C-5. The relative stereochemistry was acquired based on the NOESY spectrum, where H-1 displayed association with H-5 revealing that H-1 was on the same side with H-5. However, no correlation was observed between H-5 and H-6. It reflects the trans orientation of H-5 and H-6 (Figure 2). The absolute stereochemistry of chemical substance **1** was identified as (1*R*,5*R*,6*R*)-11-hydroxyl-drocostuslactone via X-ray crystallographic analysis, as well (Figure 3).




Figure 2. 2D NMR correlation of **1**


Sesquiterpenoids from the leaves of *Dalbergia odorifera*

Figure 3. ORTEP drawing of compound **1**

Compound **2** was separated as achromatic oil. The molecular formula $C_{15}H_{24}O_2$ was determined by HR-ESI-MS data (m/z 235.1835 $[M+H]^+$). The 1H NMR spectral data of **2** displayed resonances for an olefinic proton [δ_H 5.84(1H, d, $J=1.9$ Hz, H-6)], three methines groups [δ_H 2.35 (1H, m, H-11), 2.31 (1H, m, H-1), 1.74 (1H, m, H-10)], four methylenes [δ_H 2.49 (1H, ddd, $J=17.4, 9.0, 5.2$ Hz, H-3 α), 2.28 (1H, m, H-11, H-3 β), 2.40 (1H, m, H-8 α), 2.25 (1H, m, H-8 β), 1.94 (1H, m, H-2 α), 1.83 (1H, m, H-2 β), 1.68 (1H, m, H-9 α), 1.64 (1H, m, H-9 β)], and four methyl groups [δ_H 2.11 (3H, s, H-15), 1.10 (3H, d, $J=6.5$ Hz, H-14), 1.08 (3H, d, $J = 2.6$ Hz, H-13), 1.06 (3H, d, $J = 2.6$ Hz, H-12)]. Inspection of its ^{13}C -NMR spectra exhibited 15 carbon resonances assignable to two carbonyls carbon at [δ_C 208.9 (C-4), 205.7 (C-5)], one double bonds at [δ_C 167.5 (C-7), 126.5 (C-6)], three methines groups [δ_C 58.1 (C-1), 37.4 (C-11), 34.8 (C-10)], three methylenes [δ_C 24.0 (C-2), 42.0 (C-3), 28.4 (C-8), 34.5 (C-9)], and four methyl carbons [δ_C 30.1 (C-15), 21.3 (C-13), 21.0 (C-12), 20.1 (C-14)]. Systematic analyses of 2D NMR spectra, which involved HSQC, HMBC (Figure 4), allowed the total evaluation of every proton and carbon signal, and the establishment of the planar architecture of **2**. Those spectroscopy characteristics revealed that the architecture of **2** resembled that of gibberodione [11-12], except for the configuration. The relative configuration was assigned from the NOESY spectrum, in which H-1 [δ_H 2.31 (1H, m, H-1)] showed correlation with H₃-14 [δ_H 1.10 (3H, d, $J=6.5$ Hz, H-14)]. However, no correlation was observed between H-1 [δ_H 2.31 (1H, m, H-1)] and H-10 [δ_H 1.74 (1H, m, H-10)]. It reflects the trans orientation of H-10 and H-1 (Figure 4). To further reinforced the structure of **2**, the ^{13}C NMR chemical shifts of four possible isomers (1*R*, 10*S*)/(1*S*, 10*R*)-**2** and (1*S*, 10*S*)/(1*R*, 10*R*)-**2** were calculated using the GIAO method at the mPW1PW91/6-311G (d, p) level with the Gaussian 09 software. The predicted chemical shifts for each isomer were weighted according to the Boltzmann distributions. The results showed that (1*R*, 10*S*)/(1*S*, 10*R*)-**2** exhibited a better coefficient of determination ($R^2 = 0.9989$) of linear correlation between the experimental and calculated ^{13}C NMR chemical shifts than (1*S*, 10*S*)/(1*R*, 10*R*)-**2** ($R^2 = 0.9978$) (Figure 5). As shown in (Figure 6), the MAE (mean absolute error) and MD (maximum deviation) for (1*R*, 10*S*)/(1*S*, 10*R*)-**2** were 1.79 and 4.6 ppm, respectively, which are acceptable to meet MAE < 2.2 and MD < 5 [13]. The DP4 + probability analysis is a method that provides high probabilities to determine the relative stereochemistry of natural products [14]. The DP4 +

probability analysis showed that the best match of with the 100% with (1*R*, 10*S*)/(1*S*, 10*R*)-**2** isomer. The (1*R*,10*S*) of **2** was determined by ECD spectra (Figure 7). Accordingly, the architecture of **2** was revealed and named as (1*R*, 10*S*)-gibberodione A.




Figure 4. Selected 2D NMR correlation of **2**




Figure 5. Linear correlation plots of calculated-experimental ^{13}C NMR chemical shift values for (1*R*, 10*S*)/(1*S*, 10*R*)-**2** and (1*S*, 10*S*)/(1*R*, 10*R*) **2**




Figure 6. Relative errors between experimental and calculated ^{13}C NMR chemical shift of (1*R*, 10*S*)/(1*S*, 10*R*)-**2** and (1*S*, 10*S*)/(1*R*, 10*R*) **2**


Sesquiterpenoids from the leaves of *Dalbergia odorifera*

Figure 7. Experiment-derived and computed ECD spectra of compound

3.2. Anti-inflammatory Activity

Two new sesquiterpenoids were evaluated for their anti-inflammatory activity on the model of the LPS-stimulated RAW 264.7 macrophages. The cytotoxicity of compounds was tested by cell counting kit-8 (CCK-8) assay, using RAW 264.7 cells by using the same method in the literature [15-17]. The results of assays revealed that, in the range of 0~16 and 0~2 $\mu\text{g/mL}$, two compounds had no cytotoxicity to RAW 264.7, and the IC_{50} values with 107.2 ± 4.02 and 54.64 ± 1.89 $\mu\text{g/mL}$, respectively. We further investigated the effects of compounds in the LPS-activated release of the inflammation mediators nitric oxide (NO) and lactate dehydrogenase (LDH) activity by RAW 264.7 cells (Figure 8). Compound **1** had significantly decreased the production of NO with 2~4 $\mu\text{g/mL}$ ($P < 0.05$), and at 8~16 $\mu\text{g/mL}$, concentrations significantly decreased the production of NO ($P < 0.01$), in contrast to the LPS group. In contrast to the LPS group, compound **1** at 1~16 $\mu\text{g/mL}$ concentration levels significantly inhibited LDH activity ($P < 0.01$). In addition, compound **2** at 1 and 2 $\mu\text{g/mL}$ concentrations significantly decreased the production of NO and inhibited LDH activity ($P < 0.01$). Both compounds showed anti-inflammatory activity by decreasing nitric oxide (NO) production and inhibited lactate dehydrogenase (LDH) activity in LPS-stimulated RAW 264.7.




Figure 8. Effects of compounds on the production of NO and LDH in LPS-stimulated RAW264.7 cells. The values shown represent the mean (n=6). ^{##} $P < 0.01$ compared to the control group; ^{*} $P < 0.05$ and ^{**} $P < 0.01$ compared to the LPS group

Acknowledgments

The work was partially supported by the National Natural Science Foundation of China (82160732) and National Key R&D Program of China (2018YFC1706102).

Supporting Information

Supporting Information accompanies this paper on <http://www.acgpubs.org/journal/records-of-natural-products>

Sesquiterpenoids from the leaves of *Dalbergia odorifera***ORCID** Chenxiao Shen: [0000-0002-1331-5621](https://orcid.org/0000-0002-1331-5621)Can Yue Ouyang: [0000-0002-8108-0369](https://orcid.org/0000-0002-8108-0369)Ying Zhang: [0000-0001-7068-7395](https://orcid.org/0000-0001-7068-7395)Qing Zhu: [0000-0003-4036-0011](https://orcid.org/0000-0003-4036-0011)Ronghua Liu: [0000-0001-5623-9000](https://orcid.org/0000-0001-5623-9000)Lanying Chen: [0000-0001-8115-8114](https://orcid.org/0000-0001-8115-8114)**References**

- [1] Chinese Pharmacopoeia Committee (2020). Pharmacopoeia of People's Republic of China, Part 1. Bei Jing: The medicine science and technology press of China, p. 210.
- [2] T. H. Kang, T. H. Tian and Y. C. Kim (2005). Isoliquiritigenin: A competitive tyrosinase inhibitor from the heartwood of *Dalbergia odorifera*, *Biomol. Ther.* **13**, 32-34.
- [3] D. S. Lee, K. S. Kim, W. Ko, B. Li, S. Keo, G. S. Jeong and Y. C. Kim (2014). The neoflavonoid latifolin isolated from MeOH extract of *Dalbergia odorifera* attenuates inflammatory responses by inhibiting NF- κ B activation via Nrf2-mediated heme oxygenase-1 expression, *Phytother. Res.* **28**, 1216-1223.
- [4] C. Lee, J. W. Lee, Q. Jin, D. S. Jang, S. Lee, D. Lee, J. T. Hong, Y. Kim, M. K. Lee and B. Y. Hwang (2013). Inhibitory constituents of the heartwood of *Dalbergia odorifera* on nitric oxide production in RAW 264.7 macrophages, *Bioorg. Med. Chem. Lett.* **23**, 4263 - 4266.
- [5] L. L. Sun (2005). Studies on the chemical constituents of parts n-butanol of Lignum *Dalbergia odorifera*, *Guangdong Pharmaceut. Uni.* 11-33.
- [6] H. Wang, W. L. Mei, Z. K. Guo, Z. F. Xia, M. H. Zhong and H. F. Dai (2014). Chemical constituents of *Dalbergia odorifera*, *China J. Chin. Mater. Med.* **39**, 1625-1629
- [7] F. Y. Ma, M. Luo, C. J. Zhao, C. Y. Li, W. Wang, C. B. Gu, Z. F. Wei, Y. G. Zu and Y. J. Fu (2013). Simple and efficient preparation of biochanin A and genistein from *Dalbergia odorifera* T. Chen leaves using macroporous resin followed by flash chromatography, *Sep. Purif. Technol.* **120**, 310-318.
- [8] D. Y. Zhang, Y. G. Zu, Y. J. Fu, M. Luo, C. B. Gu, W. Wang and X. H. Yao (2011). Negative pressure cavitation extraction and antioxidant activity of biochanin A and genistein from the leaves of *Dalbergia odorifera* T. Chen, *Sep. Purif. Technol.* **83**, 91-99.
- [9] R. Ma, H. Liu, F. Shi, Y. Fu, P. Wei and Z. Liu, (2020). The chemical composition and anti-oxidant activity of essential oils and extracts of *Dalbergia odorifera* leaves, *Holzforchung* **74**, 755-763.
- [10] D. H. De Luengo, M. Miski, D. A. Gage and T. J. Mabry (1986). 7 α -Hydroxy-sesquiterpene lactones from *Decachaeta ovatifolia*, *Phytochemistry* **25**, 1917-1922.
- [11] A. F. Ahmed, Y. H. Kuo, C. F. Dai and J. H. Sheu (2005). Oxygenated terpenoids from a formosan soft coral *sinularia gibberosa*, *J. Nat. Prod.* **68**, 1208-1212.
- [12] K. Ng, M. Khoury and T. G. Minehan (2021). Total Syntheses and Absolute Configuration Assignment of (+)- Sootepdienone, (-)-Jambolanin C, (-)-Jambolanin I, and (-)-Gibberodione, *J. Org. Chem.* **86**, 3074-3080.

- [13] X. Zheng, A. Kadir, G. Zheng, P. Jin and G. Yao (2020). Antiproliferative abietane quinone diterpenoids from the roots of *salvia deserta*, *Bioorg. Chem.* **104**, 104261.
- [14] D. Xin, P. J. Jones and N. C. Gonnella (2018). DiCE: diastereomeric in silico chiral elucidation. expanded DP4 probability theory method for diastereomer and structural assignment, *J. Org. Chem.* **83**, 5035-5043.
- [15] Y. Lu and Y. F. Chen (2021). Eudesmane sesquiterpenoids from *Salvia plebeia*, *Rec. Nat. Prod.* **15**, 613-616.
- [16] D. Yuan, Z. Ying, F. He, X. Ying, L. Jia, G. Yang (2021). A new skeleton flavonoid and a new lignan from *Portulaca oleracea* L. and their activities, *Fitoterapia.* **153**, 104993.
- [17] Y. Gao, W. Jiang, C. Dong, C. Li, X. Fu, L. Min (2012). Anti-inflammatory effects of sophocarpine in lps-induced raw 264.7 cells via nf- κ b and maps signaling pathways, *Toxicology in Vitro.* **26**, 1-6.

A C G
publications

© 2022 ACG Publications