






Plagranline A, a Novel β -carboline and the First Alkaloid Isolated from *Platycodon grandiflorus*

Ju-Tao Wang ^{1,2,3}, Lan Ding ¹, Ren-Nai Yang ¹, Liang-Ping Zha ^{1,3}
Yang Yu ^{*1,2,3} and Shuang-Ying Gui ^{*1,4,5,6}

¹ School of Pharmacy, Anhui University of Chinese Medicine, Hefei, 230012, PR China

² Institute of Medicinal Chemistry, Anhui Academy of Chinese Medicine, Hefei 230012, PR China

³ Anhui Province Key Laboratory of Research & Development of Chinese Medicine, Hefei, 230012, PR China

⁴ Institute of Pharmaceutics, Anhui Academy of Chinese Medicine, Hefei 230012, PR China

⁵ Anhui Province Key Laboratory of Pharmaceutical Preparation Technology and Application, Hefei, Anhui, 230012, PR China

⁶ Engineering Technology Research Center of Modernized Pharmaceutics, Anhui Education Department (AUCM), Hefei, Anhui, 230012, PR China

(Received July 10, 2022; Revised August 11, 2022; Accepted August 14, 2022)

Abstract: A novel β -carboline, namely Plagranline A (**1**), and a known alkaloid (**2**) were isolated from the roots of *Platycodon grandiflorus*. Their structure was elucidated using a combination of spectroscopic analyses and computational chemistry, including electronic circular dichroism (ECD) and optical rotatory dispersion (ORD). Alkaloids **1** and **2** exhibited moderate inhibitory effects against NO production. Interestingly, **1** and **2** are the first alkaloid constituents reported from the plant *P. grandiflorus*.

Keywords: *Platycodon grandiflorus*; β -carboline; computational chemistry; alkaloid; anti-inflammatory activity.
© 2022 ACG Publications. All rights reserved.

1. Plant Source

P. grandiflorus roots were collected from Taihe, Anhui Province of China, in July 2021. The plant was identified by Qing-shan Yang of the Anhui University of Chinese Medicine. A voucher specimen (No. 202101) was deposited at Anhui University of Chinese Medicine.

* Corresponding author: E- Mail: ywhxyu@126.com (Yang Yu), guishy0520@126.com (Shuang-Ying Gui)

Plagranline A, a novel β -carboline from *Platycodon grandiflorus*

2. Previous Studies

Platycodon grandiflorus (Jacq.) A. DC. is a resascent herb belonging to the family Campanulaceae, which is widely distributed in Asian countries. As a well-known food material, it is frequently used as an ingredient in healthy foods and vegetable dishes. Meanwhile, it is used in folk medicine to treat cough, lung abscess, chest pain, sore throat, and diabetes [1]. Chemical investigations on *Platycodon grandiflorus* have mainly focused on and characterized triterpenoid saponins [2-4], flavonoids [5], phenolic acids (lignans) [6], polyacetylene [7], as well as sterols [1] and polysaccharides [8]. Moreover, there were a handful of records of the alkaloid content of the plant with UPLCQTOF-MS/MS techniques in the literature [9]. Among the above constituents, triterpenoid saponins, especially Platycodin D have been studied further, while polysaccharides have also been investigated. In summary, there are only limited chemical studies on *Platycodon grandiflora*, and their structural variation has not been further investigated in recent years.

3. Present Study

Two β -carboline-type alkaloids (**1-2**) (Figure 1) were isolated from the roots of *Platycodon grandifloras*, providing better insights into its constituents.

The dried roots of *P. grandiflorus* (20 kg) were reflux extracted with 80% ethanol (2 h \times 3). The solution was evaporated under reduced pressure to obtain an extract (1.2 kg). The crude extract was suspended in distilled water (1.5 L), and then partitioned with EtOAc and *n*-BuOH (each 1.5 L \times 3 times). The EtOAc-soluble extract (276 g) was separated by silica gel CC elution with a step gradient of CH₂Cl₂/MeOH (100:0 to 0:100, v/v) to obtain six fractions (Fraction A–Fraction F). Fraction E (39 g) was subjected to C₁₈ MPLC using MeOH–H₂O (20–80%, v/v) to yield five subfractions (E-1 to E-5). Subfraction E-2 (5 g) was chromatographed on Sephadex LH-20 (MeOH) twice and was further purified on the HPLC preparative column with CH₃CN–H₂O (20–35%, v/v, 40 min) to afford **2** (5.2 mg, 28 min). Subfraction E-5 (1 g) was chromatographed on Sephadex LH-20 (MeOH) and was further purified on the HPLC preparative column with CH₃CN–H₂O (35–50%, v/v, 40 min) to afford **1** (0.8 mg, 35 min).

Plagranline A (1): Yellow amorphous powder; C₂₀H₂₀N₂O₄, [α]_D²⁴ = -21.3 (*c* 0.6, CH₃OH); UV (CH₃OH) λ_{\max} (log ϵ): 202 (4.39), 278 (3.95), 362 (3.68) and 379 (3.67) nm, ¹H (800 MHz) and ¹³C (200 MHz) NMR data (CD₃)₂CO, acetone-*d*₆, Table 1; HRESIMS *m/z* 375.1314 [M+Na]⁺ (calcd. for C₂₀H₂₀N₂O₄Na 375.1315).




Figure 1. ¹H–¹H COSY correlations and the selected HMBC correlations of alkaloid **1** and the structures of alkaloids **1-2**

Alkaloid **1** was obtained as a yellow amorphous powder with maximum UV absorbance at 202, 278, 362, and 379 nm, which revealed the presence of a conjugated group. Its molecular formula was C₂₀H₂₀N₂O₄ based on HRESIMS with twelve degrees of unsaturation. The ¹H NMR data of **1** (Table 1)

and the HSQC spectrum showed the characteristic signals of an aromatic ring, including a group of ABX system spin coupling [7.94 (1H, d, $J = 2.5$ Hz, H-5), 7.36 (1H, dd, $J = 8.8, 2.5$ Hz, H-7) and 8.54 (1H, d, $J = 8.8$ Hz, H-8)], and a pair of olefinic protons at δ_{H} 8.83 (1H, d, $J = 4.9$ Hz, H-3) and 8.20 (1H, d, $J = 4.9$ Hz, H-4)]. The ^{13}C NMR and DEPT data (Table 1) revealed several conspicuous quaternary carbon signals at δ_{C} 125-140 ppm. Meanwhile, by combining the above hydrogen signals, in particular, an unusual pair of coupling constants (4.9 Hz) suggested that **1** was a β -carboline type. Furthermore, the key HMBC correlations from H-3 (δ_{H} 8.83) to C-1 (δ_{C} 137.4)/C-4a (δ_{C} 130.5), from H-4 (δ_{H} 8.20) to C-1a (δ_{C} 132.4)/C-5a (δ_{C} 127.3) and from H-5 (δ_{H} 7.94) to C-4a (δ_{C} 130.5)/C-8a (δ_{C} 134.5), as well as, the correlations of H-3 (δ_{H} 8.83) with H-4 (δ_{H} 8.20) and of H-7 (δ_{H} 7.36) with H-8 (δ_{H} 8.54) in the ^1H - ^1H COSY spectrum, supported the above deduction. Moreover, the remaining olefinic proton (δ_{H} 8.06) indicated that the correlations with C-1a (δ_{C} 132.4)/C-2' (δ_{C} 138.7)/C-7' (δ_{C} 162.3) in the HMBC spectrum, pointed to the presence of α, β -unsaturated lactam ring between N₉-C₁ and C₁'-C₇'. Finally, five high field signals in the 2D spectra were distinctly observed: the ^1H - ^1H COSY correlation with H-5' and H-6', together with the HMBC correlation from H-8' (δ_{H} 3.37/3.41) to C-3' (δ_{C} 38.4)/C-5' (δ_{C} 30.7), from H-5' (δ_{H} 1.64/1.69) to C-3' (δ_{C} 38.4) and from H-6' (δ_{H} 1.00) to C-4' (δ_{C} 75.5), suggested the presence of 2-methylbutane-1,2-diol group. The group was located at C-2' that confirmed by the crucial correlations from H-1' (δ_{H} 8.06) to C-3' (δ_{C} 38.4) and from H-3' (δ_{H} 2.97/3.13) to C-7' (δ_{C} 162.4). Therefore, the planar structure of **1** was assigned.

Because one chiral center is present in the structure of **1**, the ECD spectrum was applied to unambiguously confirm its absolute configuration, which was further supported with the aid of specific rotation. The calculated ECD spectra were determined using time-dependent density functional theory (TD-DFT) calculations at the M06-2X-D3/def2-TZVP level with MeOH as the solvent [10] using the polarizable continuum model (PCM), as well as, the calculated specific rotation was established by TDDFT at the APFD/6-311++g(2d,2p) (589 nm). As a result, the calculated ECD or ORD showed a good match with their experimental ones. Consequently, the absolute configuration of **1** was determined as 4'*R* (Figure 2 and S13-Figure 1).

Alkaloid **2** was identified as (3*S*,4*R*)-1-(β -carbolin-1-yl)-3,4,5-trihydroxy-1-pentanone by comparing their NMR spectroscopic data with those reported in the literature [11]. Interestingly, alkaloids **1** and **2** are the first alkaloids to be identified from *P. grandiflorus*, which have expanded the chemical diversity of this plant.

Table 1. ^1H (800 MHz) and ^{13}C (200 MHz) NMR spectroscopic data for **1** in $(\text{CD}_3)_2\text{CO}$ (δ in ppm and J in Hz)

No.	δ_{H}	δ_{C}	NO.	δ_{H}	δ_{C}
1		137.4, C	1'	8.06 (s)	139.8, C
1a		132.4, C	2'		138.7, C
3	8.83 (d, 4.9)	146.9, CH	3'	2.97 (d, 13.7) 3.13 (d, 13.7)	38.4, CH ₂
4	8.20 (d, 4.9)	117.1, CH	4'		75.5, C
4a		130.5, C	4'-OH	3.91 (s)	
5	7.94 (d, 2.5)	107.9, CH	5'	1.64 (q, 7.5) 1.69 (q, 7.5)	30.7, CH ₂
5a		127.3, C	6'	1.00 (t, 7.5)	7.8, CH ₃
6		159.1, C	7'		162.3, C
7	7.36 (dd, 8.8, 2.5)	118.6, CH	8'	3.37 (dd, 11.3, 5.6) 3.41 (dd, 11.3, 6.9)	66.6, CH ₂
8	8.54 (d, 8.8)	118.5, CH			
8a		134.5, C			
6-OCH ₃	3.97 (s)	56.2, CH ₃			

Plagranline A, a novel β -carboline from *Platycodon grandiflorus*




Figure 2. Calculated and experimental and ECD of **1**

Table 2. IC₅₀ values of alkaloids **1–2** inhibiting NO production in RAW 246.7 cells

Compound	IC ₅₀ (μM) ^a
1	36.18±1.02
2	37.45±1.75
L-NMMA^b	32.95±0.44

^aIC₅₀ values were expressed as mean ± SD (*n* = 3).

^bL-NMMA = L-NG-Monomethylarginine hydrochloride was used as the positive control.

Alkaloids **1** and **2** were assessed for their abilities to inhibit NO production in LPS-stimulated RAW 264.7 cells [12, 13]. The results (Table 2) indicated that compounds **1** and **2** displayed moderate inhibitory effects against NO production with IC₅₀ values of 36.18 and 37.45 μM, respectively.

Acknowledgments

This work was partially supported by the National Natural Science Fund for the National Natural Science Foundation of China (No. U21A20406).

Supporting Information

Supporting Information accompanies this paper on <http://www.acgpubs.org/journal/records-of-natural-products>

ORCID

Ju-Tao Wang: [0000-0001-8258-7658](https://orcid.org/0000-0001-8258-7658)

Lan Ding: [0000-0001-6042-1497](https://orcid.org/0000-0001-6042-1497)

Ren-Nai Yang: [0000-0003-0627-4777](https://orcid.org/0000-0003-0627-4777)

Liang-Ping Zha: [0000-0002-4644-5852](https://orcid.org/0000-0002-4644-5852)

Yang Yu: [0000-0002-9477-2313](https://orcid.org/0000-0002-9477-2313)

Shuang-Ying Gui: [0000-0002-3855-6632](https://orcid.org/0000-0002-3855-6632)

References

- [1] L. Zhang, Y. L. Wang, D. W. Yang, C. H. Zhang, N. Zhang, M. H. Li and Y. Z. Liu (2015). *Platycodon grandiflorus* an ethnopharmacological, phytochemical and pharmacological review, *J. Ethnopharmacol.* **164**, 147-161.
- [2] W. Li, W. Zhang, L. Xiang, Z. Wang, Y. N. Zheng, Y. P. Wang and L. Chen (2010). Platycoside n: a new oleanane-type triterpenoid saponin from the roots of *Platycodon grandiflorum*, *Molecules* **15**, 8702-8708.
- [3] L. Zhang, Z. H. Liu and J. K. Tian (2007). Cytotoxic triterpenoid saponins from the roots of *Platycodon grandiflorum*, *Molecules* **12**, 832-841.
- [4] Y. H. Choi, D. S. Yoo, M. R. Cha, C. W. Choi, Y. S. Kim, S. U. Choi and S. Y. Ryu (2010). Antiproliferative effects of saponins from the roots of *Platycodon grandiflorum* on cultured human tumor cells *Molecules, J. Nat. Prod.* **73**, 1863-1867.
- [5] I. Mazol, M. Glensk and W. Cisowski (2004). Polyphenolic compounds from *Platycodon grandiflorum* A. DC, *Acta. Pol. Pharm.* **61**, 203-208.
- [6] W. Li and H. J. Yang (2022). Phenolic Constituents from *Platycodon grandiflorum* root and their anti-inflammatory activity, *Molecules* **26**, 4530.
- [7] W. Li, W. Zhang, L. Xiang, Z. Wang, Y. N. Zheng, Y. P. Wang and L. Chen (2010). Platycoside n: a new oleanane-type triterpenoid saponin from the roots of *Platycodon grandiflorum*, *Molecules* **15**, 8702-8708.
- [8] X. D. Dong, Y. N. Liu, S. S. Yu, H. Y. Ji, Y. Y. Feng, A. J. Liu and J. Yu (2021). Extraction, optimization, and biological activities of a low molecular weight polysaccharide from *Platycodon grandiflorus*, *Ind. Crop. Prod.* **165**, 113427.
- [9] C. Z. Wang, N. Q. Zhang, Z. Z. Wang, Z. Qi, B. Z. Zheng and P. Y. Li (2017). Rapid characterization of chemical constituents of *Platycodon grandiflorum* and its adulterant *Adenophora stricta* by UPLCQTOF-MS/MS, *J. Mass Spectrom.* **52**, 643–656.
- [10] J. T. Wang, H. Y. Li, R. Rao, Y. J. Yue, G. K. Wang and Y. Yu (2021). (±)-Stagonosporopsin A, stagonosporopsin B and stagonosporopsin C, antibacterial metabolites produced by endophytic fungus *stagonosporopsis oculihominis*, *Phytochem. Lett.* **45**, 157–160.
- [11] N. Aimi, M. Kitajima, N. Oya, W. Nitta, H. Takayama, S.-I. Sakai, I. Kostenyuk, Y. Gleba, S. Endreb and J. Stockigt (1996). Isolation of alkaloids from cultured hybrid cells of *Rauwolfia serpentina* and *Rhazya stricta*, *Chem. Pharm. Bull.* **44**, 1637-1639.
- [12] Y. Lu and Y.F. Chen (2021). Eudesmane Sesquiterpenoids from *Salvia plebeia*, *Rec.Nat.Prod.***15**, 613-616.
- [13] Y. J. Yue and Y. Yu (2022). Isolation and identification of endophytic fungi from *Dendrobium huoshanense* with their antibacterial and anti-inflammatory activities, *Pak. J. Pharm. Sci.* **35**, 1143-1151

A C G
publications

© 2022 ACG Publications