

## Polyketides and Alkaloids from the Fungus *Penicillium* sp.

Yifang Chen , Ying Lu , Jianyong Zhao  and Qian Chen \*

The First People's Hospital of Linping District, Hangzhou, Zhejiang 311100, China

(Received October 03, 2022; Revised October 18, 2022; Accepted October 20, 2022)

**Abstract:** The marine-derived fungus SCZ-1 was cultured on rice solid medium. The fermented materials was extracted using ethyl acetate to afford an extract, which was separated by repeated column chromatography to afford three polyketides and five alkaloids, including 2 new polyketides (**1** and **2**). The known compounds were identified to be penipyrrol A (**3**), cyclopenin (**4**), cyclophenol (**5**), terretrione D (**6**), viridicatol (**7**), and viridicatin (**8**). The structures of **1** and **2** were determined by extensive analyses of spectroscopic data (1D and 2D NMR, and HRESIMS). The absolute configuration of **2** was assigned by comparison of the experimental ECD spectrum with calculated ECD spectrum. Compound **1**, bearing a two-carbon side chain at C-2, is seldom found among natural isocoumarins. All metabolites were screened for their inhibitory effects toward  $\alpha$ -glucosidase, only compound **8** showed weak activity with an IC<sub>50</sub> value of 548  $\mu$ M.

**Keywords:** marine-derived fungus; *Penicillium* sp.; new polyketides. © 2022 ACG Publications. All rights reserved.

### 1. Plant Source

The fungal strain SCZ 4-1 was isolated from sediments collected from the Mirs Bay of Shenzhen. The strain was identified as *Penicillium* sp. by analyses of the ITS region of the rDNA sequence with those of a standard record. The ITS sequence has been sent to the GenBank (<http://www.ncbi.nlm.nih.gov>) with the accession number KY445779.

### 2. Previous Studies

In recent years, the metabolites of fungi have been deeply excavated by natural product chemists, leading to the discovery of thousands of molecules with various structures and profound activity [1-6]. The *Penicillium* is an outstanding member which have been proved to be productive. Previous chemical investigation of *Penicillium* strains led to identification of meroterpenoids [7-9], alkaloids [10, 11], polyketides [12, 13], sesquiterpenes and diterpenes [14-17].

In our study, we found that the <sup>1</sup>H NMR data of the fermented materials of the marine-derived *Penicillium* sp. SCZ4-1 showed resonances that were probable for alkaloids. Subsequent chromatographic separation of the EtOAc extract led to the isolation of eight compounds including three polyketides (**1–3**) and five alkaloids (**4–8**). Herein, we reported the isolation and structural identification of these metabolites.

### 3. Present Study


The strain was cultured on solid rice medium (80 g rice and 90 mL water) in 20 erlenmeyer

\*Corresponding author: E- Mail: [18072889003@163.com](mailto:18072889003@163.com) (Q. Chen)

flasks (500 mL). The fermentation was carried out for 4 weeks under static conditions at 25 °C. The fermented materials were extracted with EtOAc (1000 mL) for three times and afforded an EtOAc extract (2.1 g), which was separated on a silica gel with petroleum ether/EtOAc (95:5 to 50:50) as eluent to give five fractions (Fr.I–Fr.V). Fra. II was chromatographed over ODS silica gel (MeOH/H<sub>2</sub>O = 20:80 to 100:0) to give five fractions (Fr.IIa–Fr.IIe). Fr.IIb was purified by HPLC (a YMC-pack ODS-A column, 2 mL/min) with MeCN/H<sub>2</sub>O (50:50) as eluent to give **5** (8.5 mg), **8** (1.8 mg), and **7** (2.3 mg). Fr.IIc was purified on HPLC using 60% acetonitrile/water as eluent to give **2** (2.7 mg), **4** (5.7 mg), and **6** (2.3 mg). Fr.III was applied to the ODS gel (MeOH/H<sub>2</sub>O = 20:80 to 100:0) to give five fractions (Fr.IIIa– Fr.IIIe). Fr.IIIb was separated by HPLC (MeCN/H<sub>2</sub>O = 45:55) to yield **1** (1.5 mg) and **3** (2.5 mg).

*Peniketide A (1)*: Light yellow oil, UV (MeOH)  $\lambda_{\max}$  (log  $\epsilon$ ) 239 (4.36), 244 (4.39) nm; <sup>1</sup>H NMR and <sup>13</sup>C NMR data, see Table 1; HRESIMS  $m/z$ : 237.0388 [M + H]<sup>+</sup> (calcd. for C<sub>11</sub>H<sub>9</sub>O<sub>6</sub><sup>+</sup>, 237.0394).

*Methyl ester of penipyrol A (2)*: Colorless oil; [ $\alpha$ ]<sub>D</sub><sup>25</sup> –34 (c = 0.1, CH<sub>3</sub>OH); UV (MeOH)  $\lambda_{\max}$  291 (3.77) nm; ECD (c 7.1 × 10<sup>-4</sup> M, MeOH)  $\lambda_{\max}$  ( $\Delta\epsilon$ ) 314 (–0.49), 254 (–1.87), 209 (+1.27) nm; <sup>1</sup>H and <sup>13</sup>C NMR data, Table 1; HRESIMS  $m/z$  275.1287 [M – H]<sup>–</sup> (calcd. for C<sub>16</sub>H<sub>19</sub>O<sub>4</sub><sup>–</sup>, 275.1289)




**Figure 1.** Structures of compounds **1–8** from the strain *Penicillium* sp.

Compound **1** had the molecular formula of C<sub>11</sub>H<sub>8</sub>O<sub>6</sub> as assigned by the HRESIMS (237.0388 [M + H]<sup>+</sup>, calcd. for C<sub>11</sub>H<sub>9</sub>O<sub>6</sub><sup>+</sup>, 237.0394), indicating eight sites of unsaturation. The <sup>1</sup>H NMR spectrum displayed the signals for three aromatic protons [ $\delta_{\text{H}}$  6.47 (1H, br s), 6.34 (2H, br s)] and two protons for a methylene [ $\delta_{\text{H}}$  3.52 (2H, br s)]. The <sup>13</sup>C NMR exhibited 11 carbon resonances in total, which were attributed by HSQC spectrum to three sp<sup>2</sup> methine ( $\delta_{\text{C}}$  107.8, 104.0, 102.9), seven sp<sup>2</sup> non-protonated carbons ( $\delta_{\text{C}}$  173.0, 167.5, 167.3, 164.6, 152.6, 141.0, 99.8), and a sp<sup>3</sup> methylene carbon ( $\delta_{\text{C}}$  40.3). The above-mentioned data indicated that the structure of **1** contained a benzene ring, a carboxyl carbon, an ester carbon, and two carbons for a double bond. As the unsaturated groups accounted for seven of the eight sites of unsaturation, the remaining one required the presence of an additional ring in the structure. These data were very similar to those of a reported natural analog (**1a**) except for the presence of a methylene ( $\delta_{\text{H}}$  3.52;  $\delta_{\text{C}}$  40.3) and an additional carbonyl carbon ( $\delta_{\text{C}}$  173.0) instead of the allylic methyl ( $\delta_{\text{H}}$  2.22;  $\delta_{\text{C}}$  19.0) in **1a** [18], suggesting that the methyl in **1a** was replaced by an acetic acid moiety in **1**. The deduced structure was further confirmed by 2D NMR analyses (Figure 2) and MS data, especially the HMBC correlations from methylene protons H<sub>2</sub>-11 ( $\delta_{\text{H}}$  3.52) to C-2 ( $\delta_{\text{C}}$  152.6), and the carboxylic carbon C-12 ( $\delta_{\text{C}}$  173.0) and from the olefinic proton H-1 ( $\delta_{\text{H}}$  6.47) to C-2, C-11 ( $\delta_{\text{C}}$  40.3), and C-12. Thus, the structure of **1** was determined as 7,9-dihydroxy-

3-acetic acidisocoumarin, interestingly, isocoumarin bearing a two-carbon side chain at C-2 was rarely found in nature. Compound **1** was named peniketide A.

**Table 1.**  $^1\text{H}$  (400 Hz) and  $^{13}\text{C}$  NMR (100 Hz) Data of **1–3** and **1a** in methanol- $d_4$  ( $\delta$  in ppm)


No.	<b>1</b>		<b>1a</b>		<b>2</b>		<b>3</b>	
	$\delta_{\text{H}}$	$\delta_{\text{C}}$	$\delta_{\text{H}}$	$\delta_{\text{C}}$	$\delta_{\text{H}}$	$\delta_{\text{C}}$	$\delta_{\text{H}}$	$\delta_{\text{C}}$
1	6.47, br s	107.8	6.47	102.1				
2		152.6		155.7		165.2		165.1
3						124.6		124.4
4		167.3		166.9	7.32, d (6.9)	142.4	7.30, d (6.9)	142.3
5		99.8		99.8	6.26, d (6.9)	106.7	6.25, d (6.9)	106.6
6		164.6		165.0		159.0		159.0
7	6.34, br s	102.9	6.30	103.8	6.25, d (15.8)	119.6	6.23, d (15.8)	119.4
8		167.5		167.0	7.04, d (15.8)	138.7	7.04, d (15.8)	138.8
9	6.34, br s	104.0	6.33	105.9		137.0		136.6
10		141.0		141.8	5.75, d (9.9)	136.0	5.78, d (9.8)	136.6
11	3.52, br s	40.3	2.22	19.0	3.42, ddd (9.9, 7.2, 7.2)	48.0	3.34, ddd (9.8, 7.2, 7.2)	47.1
12		173.0			1.83, m	27.1	1.83, m	27.2
					1.61, m		1.61, m	
13					0.93, t (7.4)	11.9	0.94, t (7.4)	12.0
14						175.7		177.2
15					1.88, s	12.7	1.89, s	12.7
16					2.06, s	16.6	2.06, s	16.7
OCH <sub>3</sub>					3.70, s	52.4		



**Figure 2.** Key HMBC ( $\rightarrow$ ), COSY ( $\rightarrow$ ), NOESY ( $\dashrightarrow$ ) correlations of **1** and **2**

The molecular formula of compound **2** was assigned as  $\text{C}_{16}\text{H}_{20}\text{O}_4$  on the basis of the HRESIMS (275.1287  $[\text{M} - \text{H}]^-$ , calcd. for  $\text{C}_{16}\text{H}_{19}\text{O}_4^-$ , 275.1289) and NMR data (Table 1), requiring seven degrees of unsaturation. The  $^1\text{H}$  NMR spectrum showed the resonances for two olefinic methyls ( $\delta_{\text{H}}$  2.06 and 1.89), a triplet of methyl ( $\delta_{\text{H}}$  0.94), a singlet of methoxy group ( $\delta_{\text{H}}$  3.70), five olefinic protons ( $\delta_{\text{H}}$  7.30, 7.04, 6.25, 6.23, 5.78) including two for a *trans* double bond [ $\delta_{\text{H}}$  7.04 (15.8 Hz), 6.23 (15.8 Hz)], and three aliphatic protons ( $\delta_{\text{H}}$  3.34, 1.83, 1.61), while the  $^{13}\text{C}$  NMR and HSQC spectra presented 15 carbon resonances totally, involving eight olefinic carbons ( $\delta_{\text{C}}$  165.2, 142.4, 138.7, 137.0, 136.0, 124.6, 106.7, 119.6), two carbonyl carbons ( $\delta_{\text{C}}$  175.7, 159.0), four methyls ( $\delta_{\text{C}}$  16.6, 12.7, 11.9), a methoxy ( $\delta_{\text{C}}$  52.4), one methylene ( $\delta_{\text{C}}$  27.1), one methine ( $\delta_{\text{C}}$  48.0). The aforementioned data were similar to the co-isolated know compound penipyrol A (**3**) with only distinction owing to the presence of a methoxy ( $\delta_{\text{H}}$  3.70,  $\delta_{\text{C}}$  52.4), suggesting **2** was the methyl derivative of **3** [19]. The speculated structure of **2** was confirmed by 2D NMR analyses. Specifically, the HMBC correlation from the methoxy protons to the carbonyl carbon at  $\delta_{\text{C}}$  175.7 established the methyl ester moiety. The geometries of  $\Delta^7$  and  $\Delta^{10}$  in **2** were both assigned as *E*-configuration by NOESY experiments (Figure 2) and by comparison of its NMR data with those of **3**. The absolute configuration of **2** was

determined by comparison of experimental ECD spectrum with the calculated ECD spectrum, which led to the assignment of the 11*R* configuration for **2** (Figure 3).



**Figure 3.** Experimental ECD spectrum of **2** in MeOH and the calculated ECD spectra of *R*-**2** at the b3lyp/6-31+g(d,p) level

Besides, the other known compounds were elucidated to be penipyrrol A (**3**) [19], cyclopinin (**4**) [20], cyclopinol (**5**) [20], terretrione D (**6**) [21], viridicatol (**7**) [22], viridicatin (**8**) [20], by comparing their <sup>1</sup>H and <sup>13</sup>C NMR data with reported data in the literature.

All the isolated metabolites were tested for the activity toward the  $\alpha$ -glucosidase according to the literature [23], only compound **8** showed weak inhibitory effect with an IC<sub>50</sub> value of 548  $\mu$ M (the positive control acarbose: 217  $\mu$ M).

## Supporting Information

Supporting Information accompanies this paper on <http://www.acgpubs.org/journal/records-of-natural-products>

## ORCID

Yifang Chen: [0000-0003-3120-7218](https://orcid.org/0000-0003-3120-7218)

Ying Lu: [0000-0002-4385-0529](https://orcid.org/0000-0002-4385-0529)

Jianyong Zhao: [0000-0002-1379-0940](https://orcid.org/0000-0002-1379-0940)

Qian Chen: [0000-0003-1687-5090](https://orcid.org/0000-0003-1687-5090)

## References

- [1] A. R. Carroll, B. R. Copp, R. A. Davis, R. A. Keyzers and M. R. Prinsep (2022). Marine natural products, *Nat. Prod. Rep.* **39**, 1122-1171.
- [2] H. A. Raja, A. N. Miller, C. J. Pearce and N. H. Oberlies (2017). Fungal identification using molecular tools: a primer for the natural products research community, *J. Nat. Prod.* **80**, 756-770.
- [3] S.-K. Daley and G. A. Cordell (2021). Biologically significant and recently isolated alkaloids from endophytic fungi, *J. Nat. Prod.* **84**, 871-897.
- [4] T. El-Elimat, H. A. Raja, M. Figueroa, A. H. Al Sharie, R. L. Bunch and N. H. Oberlies (2021). Freshwater fungi as a source of chemical diversity: a review, *J. Nat. Prod.* **84**, 898-916.
- [5] Y. Li, W. Liu, W. Xu, X. Zeng, Z. Cheng and Q. Li (2020). Aspterrics A and B, new sesquiterpenes from deep sea-derived fungus *Aspergillus terreus* YPGA10, *Rec. Nat. Prod.* **14**, 18-22.

- [6] R. Yao, S. Yao and A. Hua (2020). Aromatic compounds from the marine-derived fungus *Aspergillus versicolor*, *Rec. Nat. Prod.* **14**, 427-430.
- [7] B. W. Qi, N. Li, B. B. Zhang, Z. K. Zhang, W. J. Wang, X. Liu, J. Wang, T. Awakawa, P. F. Tu, I. Abe, S. P. Shi and J. Li (2022). A multifunctional cytochrome P450 and a meroterpenoid cyclase in the biosynthesis of fungal meroterpenoid atlantinone B, *Org. Lett.* **24**, 2526-2530.
- [8] Y. Z. Wu, G. Y. Xia, H. Xia, L. Y. Wang, Y. N. Wang, L. Li, H. C. Shang and S. Lin (2022). Seco and nor-seco isodhilarane-type meroterpenoids from *Penicillium purpurogenum* and the configuration revisions of related compounds, *J. Nat. Prod.* **85**, 248-255.
- [9] J. W. Tang, L. M. Kong, W. Y. Zu, K. Hu, X. N. Li, B. C. Yan, W. G. Wang, H. D. Sun, Y. Li and P. T. Puno (2019). Isopenicins A–C: two types of antitumor meroterpenoids from the plant endophytic fungus *Penicillium* sp. sh18, *Org. Lett.* **21**, 771-775.
- [10] H. Y. Cao, J. Y. Zhao, C. Yi, S. F. Sun, K. L. Chen, Z. M. Qin and Y. B. Liu (2022). Undescribed meleagrins alkaloids from the endophytic fungus *Penicillium commune*, *Phytochemistry* **204**, 113441.
- [11] F. H. Yao, X. Liang, X. H. Lu, X. Cheng, L. X. Luo and S. H. Qi (2022). Pyrrospirones K–Q, decahydrofluorene-class alkaloids from the marine-derived fungus *Penicillium* sp. SCSIO 41512, *J. Nat. Prod.* **85**, 2071-2081.
- [12] Q. Li, W. Xu, R. Z. Fan, J. Zhang, Y. L. Li, X. W. Wang, S. Y. Han, W. Liu, M. H. Pan and Z. B. Cheng (2020). Penithoketone and penithochromones A–L, polyketides from the deep-sea-derived fungus *Penicillium thomii* YPGA3, *J. Nat. Prod.* **83**, 2679-2685.
- [13] Y. Zang, Y. Gong, Z. Shi, C. Qi, C. Chen, Q. Tong, J. Liu, J. Wang, H. Zhu and Y. Zhang (2022). Multioxidized aromatic polyketides produced by a soil-derived fungus *Penicillium canescens*, *Phytochemistry* **193**, 113012.
- [14] F. Li, W. Sun, S. Zhang, W. Gao, S. Lin, B. Yang, C. Chai, H. Li, J. Wang, Z. Hu and Y. Zhang (2020). New cyclopiane diterpenes with anti-inflammatory activity from the sea sediment-derived fungus *Penicillium* sp. TJ403-2, *Chin. Chem. Lett.* **31**, 197-201.
- [15] Z. Cheng, Y. Li, W. Xu, W. Liu, L. Liu, D. Zhu, Y. Kang, Z. Luo and Q. Li (2019). Three new cyclopiane-type diterpenes from a deep-sea derived fungus *Penicillium* sp. YPGA11 and their effects against human esophageal carcinoma cells, *Bioorg. Chem.* **91**, 103129.
- [16] Z. Hu, Y. Zhu, J. Chen, J. Chen, C. Li, Z. Gao, J. Li and L. Liu (2022). Sesquiterpenoids with phytotoxic and antifungal activities from a pathogenic fungus *Aspergillus alabamensis*, *J. Agric. Food Chem.* **70**, 12065-12073.
- [17] G. P. Yin, M. Gong, G. M. Xue, T. Gong, X. Cao, W. Wang and C. H. Hu (2021). Penispidins A–C, aromatic sesquiterpenoids from *Penicillium virgatum* and their inhibitory effects on hepatic lipid accumulation, *J. Nat. Prod.* **84**, 2623-2629.
- [18] R. Ben Ameer Mehdi, S. Sioud, L. Fourati Ben Fguira, S. Bejar and L. Mellouli (2006). Purification and structure determination of four bioactive molecules from a newly isolated *Streptomyces* sp. TN97 strain, *Process Biochem.* **41**, 1506-1513.
- [19] W. Guo, X. Kong, T. Zhu, Q. Gu and D. Li (2015). Penipyrols A–B and peniamidones A–D from the mangrove derived *Penicillium solitum* GWQ-143, *Arch. Pharmacol. Res.* **38**, 1449-1454.
- [20] R. P. Hodge, C. M. Harris and T. M. Harris (1988). Verrucofortine, a major metabolite of *Penicillium verrucosum* var. cyclopium, the fungus that produces the mycotoxin verrucosidin, *J. Nat. Prod.* **51**, 66-73.
- [21] L. A. Shaala and D. T. A. Youssef (2015). Identification and bioactivity of compounds from the fungus *Penicillium* sp. CYE-87 isolated from a marine tunicate, *Mar. Drug.* **13**, 1698-1709.
- [22] L. J. Fremlin, A. M. Piggott, E. Lacey and R. J. Capon (2009). Cottoquinazoline A and cotteslosins A and B, metabolites from an Australian marine-derived strain of *Aspergillus versicolor*, *J. Nat. Prod.* **72**, 666-670.
- [23] H. Kiziltas, A. C. Goren, Z. Bingol, S. H. Alwasel and I. Gulcin (2021). Anticholinergic, antidiabetic and antioxidant activities of *Ferula orientalis* L. Determination of its polyphenol contents by LC-HRMS, *Rec. Nat. Prod.* **15**, 513-528.