

# Mitigating Fluoride, Lead, Arsenic and Cadmium Toxicities in Laboratory Animals and Ruminants through Natural Products

Abdellatif Rahim <sup>1,2</sup>, Fatma Aydogmus-Öztürk <sup>3</sup>, Cansel Cakir <sup>4</sup>,

Abdelkhalid Essamadi <sup>2</sup> and Bouchra El Amiri <sup>\*,1</sup>

<sup>1</sup>Animal Production Unit, Regional Center Agricultural Research of Settat, National Institute for Agricultural Research (INRA), Avenue Ennasr, P.O. Box 415 Rabat Principal, 10090 Rabat, Morocco

<sup>2</sup>Hassan First University of Settat, Faculty of Sciences and Techniques, Laboratory of Biochemistry, Neurosciences, Natural Ressources and Environment, P.O. Box 577, 26000 Settat, Morocco

<sup>3</sup>Köycegiz Vocational School, Mugla Sitki Kocman University, 48121 Mugla, Türkiye

<sup>4</sup>Department of Chemistry, Faculty of Science, Mugla Sitki Kocman University, 48121 Mugla, Türkiye

(Received February 24, 2022; Revised April 17, 2022; Accepted April 17, 2022)

**Abstract:** Environmental pollutants are considered a serious health problem for humans and animals mainly in ruminants for several regions of the world. Previously, many studies have investigated the mechanisms of toxicity of these pollutants on laboratory animals. Afterward, other studies have demonstrated that exposure to environmental pollutants may cause several adverse effects on the ruminant organs, influencing their performance and leading to socio-economic problems for breeders. Fluoride, lead, arsenic, and cadmium are the most common poisonings in ruminants, they can cause several irreversible toxic effects in many organs depending on the mode of action. The adverse effects of fluoride, lead, arsenic, and cadmium toxicities in laboratory animals and ruminants have been clearly summarized in this review. In addition, several results on protective or ameliorative effects by means of natural products against these toxicities have been illustrated.

**Keywords:** Environmental pollutants; laboratory animals; ruminants; toxicity; mitigation; natural products.  
© 2022 ACG Publications. All rights reserved.

## 1. Introduction

Livestock occupies a prominent place in several regions of the world. About 1.3 billion habitats worldwide live in developing countries where their livelihood depends directly or indirectly on livestock [1]. It plays an important socio-economic and cultural role in the well-being of rural households, such as food supply, source of income, asset saving, source of employment, soil fertility, livelihoods, transport, agricultural traction, agricultural diversification, and sustainable agricultural production [2]. Ruminants including cattle, sheep, and goats are the domestic animals that dominate global livestock production [3] due to their adaptability to various climatic conditions [4]. However, in several regions,

\* Corresponding author: E-Mail : [bouchraelamiri@hotmail.com](mailto:bouchraelamiri@hotmail.com); Phone: +212523729300/07; Fax: +212523729306/ +212523720927

## Mitigating fluoride, lead, arsenic and cadmium toxicities

ruminants are continuously exposed to environmental poisonings which are harmful at specific doses [5]. Besides, it was highlighted that the most toxic environmental poisonings for ruminants are fluoride, lead, arsenic, and cadmium [6]. Furthermore, the ingestion and/or inhalation of these toxic elements can cause several adverse effects on the animal body, influencing its performance, and resulting in socio-economic problems. Moreover, these pollutants lead to irreversible toxic effects [7, 8]. Accordingly, to facilitate the understanding of the toxicity mechanisms of these poisonings and to highlight preventive solutions, several studies have been performed first on laboratory animals and then on ruminants.

To prevent the effects of all environmental pollutants cited above, several synthetic metal chelators also have been reported with encouraging degrees of success [9]. However, these synthetic chelators have toxic effects [10, 8]. Recently, research have been directed towards the use of natural biomolecules originating from medicinal plants, rich in natural minerals which can chelate these environmental pollutants, and contain a wide range of antioxidants that can reduce the oxidative stress [7, 8]. This review focuses on investigating the sources, the harmful effects, and the mechanisms of fluoride, lead, arsenic, and cadmium toxicities in both laboratory animals and ruminants. In addition, this review addresses the dietary strategies pursued to prevent the toxic effects of these elements, and the results obtained by the previous studies are illustrated to provide more information to specialist readers.

## 2. Sources of Environmental Pollutants

Environmental pollutants are widely distributed [11] and are produced by natural factors including geological degradation of bedrock materials, volcanic eruptions, salt spray, forest fires, rock weathering, biogenic sources, and wind-blown soil particles [12]. These latter are also produced by anthropogenic inputs that cause disturbances and acceleration of the natural geochemical cycle [13]. More precisely, the intensive and non-optimal use of agrochemical fertilizers, pesticides, wastewater irrigation, supplements of sewage sludge, higher atmospheric deposition by industrial units, and the combustion of fossil fuels have led to a high level of inorganic pollutants in the air and soil [14]. Compared to natural sources, anthropogenic sources are generally considered to be the major causes of increasing environmental pollutants [15]. They have the potential to contaminate both soils, the surface water and the air. Soil pollution by environmental pollutants is a critical global environmental problem [16]. In addition, the presence of environmental pollutants in soils can contaminate many crops and plant species [14]. Therefore, their consumption poses risks to the health of humans and animals [17]. On the other hand, drinking water is of the essential need for human and animal survival. However, anthropogenic inputs can contaminate it with excess environmental pollutants. In addition, industrialization and urbanization are considered as a significant contributors to the increase in the level of water contamination by environmental pollutants [12]. Environmental pollutants are transported by runoff from industries, municipalities, and urban areas. Most of them eventually accumulated in the soil and sediment of water bodies [18] and are also released into the atmosphere [19] through the emission of industrial gas combustions, mining, automobile exhaust, dust, etc. The atmosphere accumulates these pollutants [20] and then transmit them to ground through precipitation [21] or contaminate the soil and water subsequently entering into the food chain of animals [19, 13].

## 3. Toxicity of Environmental Pollutants on Laboratory Animals and Ruminants

Chronic exposure to environmental pollutants by humans and animals, either through inhalation or through consumption of contaminated plants and water, can result in toxic effects. Fluoride, lead, arsenic, and cadmium, are the most popular pollutants for humans and animals [6]. Table 1 and 2 summarize the main clinical signs of these pollutants in laboratory animals and ruminants respectively, and causes adverse effects on different organs such as liver, kidneys, hard tissues, reproductive organs as well as growth and biochemical parameters.

**Table 1.** Toxic effects of fluoride, lead, arsenic and cadmium on laboratory animals

Element	Effects	References
Fluoride	Impaired tooth enamel development and the presence of white chalk-like incisors with broken tips in rats and mice	[28, 29]
	Reproductive system and fertility in mice and rabbits	[34, 35, 133]
	Altered liver function in mice	[23]
	Oxidative stress and disturbance in liver and kidney functions in rabbits, mice and rats	[134, 135]
	Heart troubles in rabbits	[136]
Lead	Reproduction and oxidative stress in rats	[57]
	Oxidative stress in rats and mice	[58]
	Hepatic and renal toxicities in rats	[55, 137]
	Systolic blood pressure and bone mineral density in rats	[56]
	Weight gain and risk of obesity in rats	[138]
Arsenic	Carcinogenesis in rats and mice	[69]
	Liver damage and fibrogenesis oxidative stress in mice	[73]
	Exacerbates atherosclerotic lesion formation and inflammation in mice	[139]
	Oxidative stress and hepatorenal toxicity in rats and mice	[140],
	Reduced plasma levels of testosterone and gonadotropin in male mice and rats	[84, 141]
Cadmium	Fetal malformations in female mice and rats	[72, 142]
	Carcinogenesis in rats and mice	[92, 93, 143]
	Increased uterine wet weight, promoted growth and development of the mammary glands in female rats	[94]
	Reproduction disturbance	[96, 98]
	Pathological testicular alterations, and liver and kidney damage in rats	[99]
	Increased lipid peroxidation in liver and kidney tissues in rats	[100]

### 3.1. Fluoride

Fluoride is the 13th most abundant halogen in nature and in the world [22]. It has many physiological roles for animals [23] but it may lead to toxic effects in critical doses. Among domestic animals, ruminants have less fluoride tolerance than simple-stomach animals [24]. Fluoride enters the ruminants body by several routes, including inhalation, consumption of contaminated water, or plants growing in contaminated soil leading to fluorosis [8]. More precisely, its bioaccumulation in teeth and bones may cause several injurious effects in the form of dental and skeletal fluorosis [24, 25]. Moreover, increasing the duration of exposure to fluoride has been shown to produce adverse effects in other tissues, leading to oxidative stress, DNA damage, apoptosis, and necrosis [26].

#### 3.1.1. Effects of Fluoride on Laboratory Animals

Rabbits, rats, and mice are all very sensitive to fluoride toxicity [27]. For this reason, these species are widely used in experimental studies to analyze the fluoride toxicity [24]. Oral administration of sodium fluoride altered the development of tooth enamel and revealed chalk-like white incisors with broken tips in rats and mice [28, 29]. On the other hand, chronic fluoride intoxication has caused bone damage, osteosclerosis, osteomalacia, and osteoporosis in rats [30, 31]. Moreover, several studies reported that

## Mitigating fluoride, lead, arsenic and cadmium toxicities

**Table 2.** Toxic effects of fluoride, lead, arsenic and cadmium on ruminants

Element	Effects	References
Fluoride	Skeletal fluorosis in buffaloes, cattle, camels, goats, and sheep.	[37]
	Dental and skeletal fluorosis in sheep.	[144]
	Mandibular lesions in sheep.	[145]
	Alteration of some serum, kidney and liver biochemical parameters in cattle.	[38]
	Increase in fluoride milk yield in goats.	[146]
	Disturbance of Cu, Ca, Mg, Mn, P, and Zn levels in sheep bones.	[41]
	Adverse effects on the antioxidant system of kidneys, liver and muscles in sheep.	[40]
	Liver disturbances and nephropathy in sheep.	[36]
Lead	Increase of concentrations lead in blood and tissues in cattle.	[65]
	Severe depression, paresis of hypoglossal nerve, ataxia, muscle twitching in cattle.	[147]
	Convulsion in cattle.	[62]
	Decrease in triiodtironine (T3) and thyroxine (T4) concentration, increase in estradiol plasma concentration in cattle.	[46]
	Paralysis of hindlimbs, lameness, osteoporosis, abortion and transient infertility in sheep.	[147]
	Nephrosis, osteoporosis, and fractures in lambs.	[60]
	Moderate excitation, ataxia, weight loss and pale mucous membranes in cattle.	[63]
Arsenic	Weakness, congestive mucous membranes, emaciation and several ulcers in different parts of the skin in ruminants.	[75]
	Weight loss, weakness, languor, ill-thrift, dry and alopecic coat that readily comes off, focal skin lesions, congested mucous membranes and stomatitis in sheep.	[56]
	Buccal erosion, stomatitis, cutaneous ulcers and serous atrophy of fat deposits were observed in ewes.	[75]
	Abnormal growth in wool, and biochemical, hematological and histopathological disturbances in sheep.	[76]
	Hepatotoxicity in goats and sheep.	[148, 149]
	Severe clinical signs of toxicity and toxico-pathological changes in goats.	[150]
	Weight loss, changes in heart and respiratory rhythms, hair loss on the flanks, and keratosis of the nose and mouth in goats.	[151]
	Presence of leukopenia, anemia and an increased rate of erythrocyte fragility in goats.	[152]
Cadmium	Disruption of some biochemical parameters in cattle.	[78]
	Meat quality affected in goats.	[102]
	Decrease in the levels of zinc and copper in sheep and cattle.	[103]
	Cadmium toxicity associated with copper deficiency in cattle.	[51]

fluoride results in liver and kidney function disturbances associated with oxidative stress in rabbits, mice, and rats [32, 33]. Furthermore, chronic fluoride intoxication caused a significant decrease in the weight of the testes, epididymis, and ventral prostate. It also negatively influenced sperm quality

parameters and fertility rate in mice [34, 35]. Fluoride also affected blood and genetic parameters in rats and mice [23].

### 3.1.2. *Effects of Fluoride on Ruminants*

Several studies in many endemic areas reported the harmful effects of fluoride on ruminants such as cattle, sheep, goats, camels, and buffaloes [36, 37, 24]. In endemic areas of India, about 31.2% of mature animals and 10.7% of immature animals such as buffaloes, cattle, camels, goats, and sheep, showed signs of skeletal fluorosis like periosteal exostoses, intermittent lameness, and tendon stiffness in the legs [37]. Moreover, dental and skeletal fluorosis have been highlighted in this species in many other endemic areas such as Morocco, Turkey, and other countries [36, 38, 39, 40, 41, 42]. On the other hand, many studies reported the negative effects of fluorosis in soft tissue, genetic parameters, teratogenic effects, apoptosis, genotoxic effects, reproductive organs, and growth in ruminants [5, 43, 44].

## 3.2. *Lead*

Lead is a common heavy metal found in low concentrations in the Earth's crust [45], it has many biological properties in ruminants' bodies however it is highly toxic when it exceeded the recommended dose [46]. It is accumulated in the environment naturally and by anthropogenic activities [47, 48]. Lead toxicity has been reported in several countries around the world such as Nigeria, Mexico, India, China, France, and United States [49]. Exposure to lead in industrial regions induces several clinic pathological changes through toxicity occurring in the kidney and endocrine systems. Moreover, a high level of lead in ruminants resulted in reproductive failure [50]. The cattle are poisoned more frequently, followed by sheep and goats [51, 48]. Additionally, the bioaccumulation (a blood concentration  $\geq 0.20 \mu\text{g}$  of Pb/mL) of this metal in meat and milk of some ruminants can be considered as a major risk for human public health [48, 52, 51].

### 3.2.1. *Effects of Lead on Laboratory Animals*

The researchers have reported various lead-induced toxic effects on laboratory animals [53]. The research on lead poisoning focuses primarily on its toxic effects on the kidneys, liver [54, 55], heart [56], and reproductive organs [50, 57]. In addition, the relationship between the toxicity of lead and the generation of oxidative stress [58, 59] has also been reported.

### 3.2.2. *Effects of Lead on Ruminants*

Toxicological studies reported several clinical symptoms in cattle, sheep, and goats [48]. The clinical signs generally presented in poisoned cattle are ill-thrift, emaciation, muscle wastage and developmental abnormalities in fetuses [60], opaque hair, thickening of phalange epiphyses, and moderate anemia [51]. Moreover, other studies have observed that lead poisoned cattle present severe depression, paresis of the hypoglossal nerve, ataxia, muscle twitching [61], convulsion, coma, respiratory failure, and death [62]. Swarup et al. [46] have reported a decrease in triiodothyronine (T3) and thyroxine (T4) concentration, and an increase in estradiol plasma concentration in some ruminants. On the other hand, lead poisoned sheep presented paralysis of hindlimbs, lameness, osteoporosis, abortion and transient infertility [61], nephrosis, osteoporosis, and fractures [60]. Whereas, goats presented moderate excitation, ataxia, weight loss, and pale mucous membranes [63].

## 3.3. *Arsenic*

Arsenic is a carcinogenic metalloid found ubiquitously in the form of organic and inorganic pollutants in water, soil, and air arising from natural and anthropogenic sources [64]. It is the 20th most abundant element on earth. According to the National Academy of Sciences Committee on Medical and Biological Effects of Environmental Pollutants, the average of arsenic content in the earth's crust is 2.5

### Mitigating fluoride, lead, arsenic and cadmium toxicities

mg/kg with a concentration in soil ranging from 0.1 to 50 mg/kg [65]. In addition, the World Health Organization has recommended that the maximum allowable concentration of arsenic in drinking water is 10 µg/L. However, its concentration in some countries such as Bangladesh, China, India, Mexico, Argentina, Chile, the United States, and Brazil has been found in the range of 50-3200 µg/L [66]. Furthermore, arsenic is readily soluble in groundwater due to pH, redox conditions, temperature, and dissolution composition [67]. Accordingly, the exposition of this metal can lead to severe clinical manifestations [68].

#### *3.3.1. Effects of Arsenic on Laboratory Animals*

Since arsenic is a toxic agent, significant efforts have been made in recent decades to test its various toxic effects using animal models, especially rodents (rats and mice). Several studies reported that chronic exposure to arsenic induces carcinogenesis in rats and mice [69, 70]. Additionally, a large number of experimental toxicological studies on both sexes of rats and mice have shown remarkable toxic effects of arsenic on reproduction [71, 72]. Similarly, a study reported that chronic arsenic exposure resulted in the fatty liver with serum aminotransferase and alanine aminotransferase levels followed by hepatic fibrosis in mice [73]. Histological and hematological disturbances have been observed in the same species [74].

#### *3.3.2. Effects of Arsenic on Ruminants*

Many species including sheep, goats, and cattle have been documented to be susceptible to arsenic poisonings [5]. Due to the high level of arsenic contamination in a village located in northwestern Iran, sheep showed various clinical signs such as weakness, atrophy, and inappropriate integument. In the same study, microscopic observations revealed hyperemia in the kidneys and heavy parasite infestation of the abomasum wall [75]. In another study, biochemical, hematological, and histopathological disturbances were observed in the same species, after with the administration of 6.6 mg/kg of body weight of sodium arsenite mixed with water for 133 days [76]. Moreover, the effect of chronic arsenic poisoning in sheep was investigated in another study [61]. It revealed that this metal causes weight loss, weakness, languor, dry coat and alopecia, focal skin lesions, congested mucous membranes, and stomatitis. On the other hand, Selby et al. [77] reported that cattle showed signs of arsenic poisoning, such as severe irritation of the digestive tract. It also reported that cattle are more sensitive to arsenic poisoning than other animals [77, 35]. In the Ghentugachi village of Nadia district, West Bengal, India, the high level of arsenic in the drinking water showed arsenic concentration was above the permissible limit, cattle exhibited decreased superoxide dismutase and catalase activities, decreased erythrocyte level and plasma nitrite level, increased rate of lipid peroxidation, protein carbonyl, and blood arsenic level in cattle, compared to those raised in free areas [78].

#### *3.4. Cadmium*

Cadmium is a non-essential heavy metal [79], which can induce toxic effects in many animals. It is considered to be one of the most toxic elements in the environment, with a wide range of organ toxicity and a long elimination half-life [80]. Anthropogenic sources, resulting in contamination of forage, feed, and water, are sources of cadmium exposure in farmed ruminants in many regions [81]. Moreover, some geographic areas are also associated with high natural concentrations of cadmium such as Ireland [82, 83], Poland, Germany Sardinia [84]. Long-term chronic exposure to cadmium has been associated with anemia, anosmia, and cardiovascular disease [85]. In affected areas, chronic ingestion or inhalation of cadmium can result in several adverse effects in ruminants, mainly in the liver and kidneys [86, 87]. Furthermore, this metal can affect growth rate [88] and reproduction [89]. Chronic ingestion of 2-5 mg/kg body weight per day for one year produced subclinical effects [90]. For that, Wilkinson et al. [91] proposed that the livers and kidneys of adult ruminants reared in contaminated areas for more than one grazing season be removed from the human food chain to reduce the risk of cadmium intake by the human population.

### 3.4.1. Effects of Cadmium on Laboratory Animals

Several experimental studies reported that cadmium results in carcinogenesis of various organs mainly kidney, lung, liver, and prostate cancer [92, 93]. In another study, exposure to cadmium increased uterine wet weight, promoted growth, and development of the mammary glands, and induced hormone-regulated genes in female rats [94]. Other toxic effects of cadmium on reproduction have been reported by Zhu *et al.* [95] such as damaged seminiferous tubules [96], production of reactive oxygen species, a decrease of spermatogonia number, and a decrease in sperm motility and count [97, 98]. On the other hand, it showed pathological testicular alterations, and liver and kidney damage after chronic exposure to cadmium in rats [99]. Moreover, other studies reported that the administration of cadmium inhibited superoxide dismutase (SOD) activity, increased endogenous levels of lipid peroxides, and increased lipid peroxidation in liver and kidney tissues in rats [100].

### 3.4.2. Effects of Cadmium on Ruminants

A study reported that cadmium poisoned sheep results in a significant reduction in humoral immunity [101]. Moreover, high levels of cadmium have been recorded in the liver, kidneys and muscles of goats grazing on areas highly contaminated with cadmium in Nsukka-Nigeria, this also affected their meat quality [102]. On the sewage treatment field of Marrakech-Morocco, the high level of cadmium found in the muscles, liver, kidneys, and bones of sheep and cattle has led to a disruption of the normal metabolism of trace elements, and decreasing the levels of zinc and copper [103].

## 4. Mechanisms of Environmental Pollutants Toxicity

After their ingestion and distribution through food or water into the body of animals, the environmental poisons exert their effects by different pathways, according to their chemical activities, biological features, and cellular targets [15, 104]. The toxicity mechanisms of environmental pollutants toxicity are not yet clear. While some studies claim that these pollutants induce their toxicity in biological systems by binding the sulfhydryl groups and generating reactive oxygen species [104]. More precisely, these pollutants are acidified in the body and oxidized to their various oxidative states. Hence, they can readily bind to biological molecules such as cations, proteins, and enzymes to form stable and strong bonds [105]. For instance, as fluoride is highly electronegative with a strong affinity towards electropositive elements, it binds with calcium, reducing their absorption and causing dental fluorosis and skeletal fluorosis [7]. Moreover, fluoride has been reported to bind to other divalent cations and inhibit many enzymes involved in the pentose phosphate pathway, antioxidant defense system, and myosin-ATPase activity [8, 106]. On the other hand, arsenic, cadmium, and lead can bind to the thiol (-SH) groups of certain enzymes and proteins, and modify their cysteine residues. In addition, they can bind to lipids and nucleic acids altering several biological activities in animals [104, 107]. Arsenic, lead, and cadmium metal ions have the ability to substitute the divalent cations such as  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Fe}^{2+}$  and monovalent cations such as  $\text{Na}^{+}$  and  $\text{K}^{+}$ , and causes significant changes in various biological processes such as cell adhesion, intra-, and intercellular signaling, protein folding, maturation, apoptosis, ion transport, enzyme regulation and neurotransmitter release [53, 108, 109].

## 5. Prevention Dietary Strategies Against Environmental Pollutants

To prevent or reduce toxicity from environmental pollutants, several preclinical and clinical studies have tested heavy metals supplementation as chelating agents to promote pollutants excretion, or synthetic antioxidants to correct the oxidative stress generated. However, these therapeutic strategies are themselves reported to have a number of different safety and efficacy concerns. Hence, several recent studies have been directed towards medicinal plants (Table 3 and 4) as natural sources of biomolecules, such as essential metals and antioxidants, which can simultaneously exercise the two therapeutic strategies, chelation and free radical scavenging [110].

## Mitigating fluoride, lead, arsenic and cadmium toxicities

**Table 3.** Selected studies natural products having ameliorative effects against fluoride, lead, arsenic and cadmium toxicities in laboratory animals

Element	Natural product	Effects	References
Fluoride	<i>Spirulina platensis</i>	Ameliorates behavioral changes, neuronal damage, and thyroid dysfunction in rats	[117]
	<i>Embllica officinalis</i>	Protective effect against hyperlipidemia and oxidative stress in rats	[153]
	<i>Tamarindus indicia</i>	Increase in urinary excretion and decrease in retention of fluoride in bones rats	[154]
Lead	<i>Lycopersicon esculentum</i>	Prevention of oxidative stress in blood plasma and kidney in rats	[155]
	<i>Camellia sinensis</i>	Alleviates the oxidative stress in the liver, and regulates hepatic metabolism in rats	[156]
	<i>Camellia sinensis</i>	Reduces lead levels and improves renal functions in rats	[157]
	<i>Allium sativum</i> and <i>Zingiber officinale</i>	Reduces blood lead levels, and regulates the hepatic and testicular cytoarchitecture in rats	[158]
Arsenic	<i>Withania somnifera</i>	Maintains the cellular integrity of testicular cells and improves sperm quality parameters in rats	[126]
	<i>Ipomea aquatica</i>	Amelioration of antioxidant parameters in liver, kidney, heart, brain and testis in mice	[127]
	<i>Tamarindus indica</i>	Regulation of serum hepatic and oxidative stress markers in rats	[128]
	<i>Triticum aestivum</i>	Attenuates arsenic-induced oxidative stress, normalizes body weight, organ weight, hematological profiles, serum biochemical profile and modulates liver and kidney biochemical parameters in rats	[159]
	<i>Moringa oleifera</i>	Reduces the elevation of serum triglycerides, glucose, urea, alkaline phosphatase, aspartate aminotransferase and alanine aminotransferase activities in mice	[132]
Cadmium	Glycine max	Ameliorates cardiac, and aorta oxidative stress in rats	[122]
	<i>Sesamum indicum</i>	Reduces serum levels, and cardiac biochemical parameters in rats	[123]
	<i>Satureja hortensis</i>	Prevents the cadmium induced lesions in hepatic functions in rats	[161]
	<i>Withania somnifera</i>	Reduces oxidative stress in testis in rats	[162]
	Royal jelly	Attenuates cadmium induced nephrotoxicity in mice	[129]
	<i>Terminalia chebula</i>	Decreases serum uric acid, urea, creatinine and total protein levels, and regulates serum lipid profiles and liver biomarkers in rats	[130]

Moreover, these natural resources provide a great variety of other nutrients, such as protein, and secondary metabolites, which have been reported to have beneficial effects against environmental pollutants [111]. Firstly, several preclinical studies on the protective effects of natural products against



these pollutants were carried out on laboratory animals. Then, other studies addressed some ruminants with encouraging levels of success. Because of their low cost, availability, and low or no side effects even at high doses [24], several preclinical studies have tested herbal medicine such as *Tamarindus indica* [112, 113], *Terminalia arjuna* [114], *Mangifera indica* [115], *Curcuma longa* [116], *Spirulina platensis* [117] and *Aloe vera* [118]. These natural products revealed encouraging results against environmental pollutants toxicities.

**Table 4.** Studies on the effects of natural products against fluoride, lead and arsenic toxicities in ruminants

Element	Natural product	Effects	References
Fluoride	<i>Tamarindus indica</i>	Reduce in serum fluoride and urinary hydroxyproline concentrations, increase in serum calcium level in cattle.	[119]
	<i>Tamarindus indica</i>	Increase in calcium level and serum alkaline phosphatase activity, decrease in serum hydroxyproline level in cattle.	[7]
	Plants rich in vitamins C and D	Reduces dental and bone damage in sheep.	[163]
Lead	<i>Allium sativum</i>	Reduces serum lead concentration and increases urine lead concentration in goats.	[121]
	<i>Allium sativum</i>	Reduces blood, kidney, bone and ovary lead levels in sheep.	[120]
Arsenic	<i>Curcuma Longa</i> and <i>Passiflora foetida</i>	Reduce the absorption of arsenic from the body, restore normal liver and kidney function, protect against oxidative stress and prevent DNA damage in sheep.	[116]
	<i>Curcuma longa</i> and <i>Zingiber officinale</i>	Remove arsenic from the body and protect against potential damage in calve.	[132]
	<i>Curcuma longa</i> and <i>Amaranthus spinosus</i>	Remove arsenic from the body and protect against DNA fragmentation in cattle.	[131]

### 5.1. Fluoride Toxicity Prevention

*Tamarindus indica* contains has a wide range of antioxidants (L-ascorbic acid, alpha-tocopherol, carotenes), essential minerals, proteins, sugars, phytosterols, and triterpenes [113]. Additionally, its beneficial ameliorative effects against fluorosis in laboratory animals was reported [112, 113]. Recent studies have tested its effects in fluorotic cattle [7, 119]. Gupta et al. [119] reported that the supplementation of fluorotic cattle with 100 g of dried powder of *Tamarindus indica* fruit pulp for 90 days, resulted in a significant increase in serum calcium and alkaline phosphatase levels, and a decrease in urinary hydroxyproline compared to fluorotic cattle not supplemented [119]. Furthermore, it was shown that chronic fluoride intoxication caused a significant decrease in plasma copper, zinc and iron concentration in cattle, while these parameters were increased in fluoridated cattle supplemented with dried powder of *Tamarindus indica* fruit pulp [7].

### 5.2. Lead toxicity Prevention

To prevent lead poisoning several dietary strategies have been tested in the laboratory animals, like *Allium sativum*, *Lycopersicon esculentum*, *Camellia sinensis*, and *Zingiber officinale*, and it was concluded that intake of diets rich in natural products may be helpful in preventing lead poisoning [111]. Furthermore, the ameliorative effect of *Allium sativum* against this intoxication has been evaluated in small ruminants, and it has been reported that this plant reduced kidney, bone, and ovary lead contents in sheep [120]. Additionally, it reduced bones, lungs, heart, liver, kidneys and skeletal muscles lead levels in goats [121].

## Mitigating fluoride, lead, arsenic and cadmium toxicities

### 5.3. Arsenic and Cadmium Toxicity Prevention

A large number of researches confirmed the beneficial effect of *Glycine max* [122], *Sesamum indicum* [123], *Satureja hortensis* [124] *Withania somnifera* [125, 126], *Ipomea aquatica* [127], *Tamarindus indica* [128], Royal jelly [129], and *Terminalia chebula* [130] on arsenic and cadmium toxicities reduction. Concerning the effects of natural products on arsenic toxicity in ruminants, it was reported that *Curcuma longa*, *Zingiber officinale*, and *Amaranthus spinosus* have eliminated the absorption of arsenic from the bovine body and protected against DNA fragmentation in these species [131, 132]. In a recent study, *Curcuma longa* powder and *Paederia foetida* powder were effectively able to remove arsenic from the body of arsenicosis induced by sodium arsenite in sheep. These two plants also helped to protect against arsenic damage, restored normal liver and kidney function, protected against oxidative stress, and prevented DNA damage [116]. In addition, the improving effects of *Curcuma longa* powder were better than those of *Paederia foetida* powder [116]. Contrastingly, until now there are no studies carried out on the evaluation of the ameliorative effects of natural plants on cadmium toxicity in ruminants.

## 6. Conclusion

This study has highlighted the importance of using natural products in the prevention and reduction of the toxic effects of fluoride, lead, arsenic, and cadmium in laboratory animals as well as in ruminants. Plant species (*Tamarindus indica* and *Allium sativum*) have been reported as promising new approaches to prevent of these poisonings. Other less-researched natural products may offer promising starting points for future studies, such as *Spirulina platensis* as a rich source of protein, minerals, and antioxidants. We expect that this review will stimulate further research, especially in endemic countries. On the other hand, we suggest more toxicity studies of these natural products in ruminants to prevent their indiscriminate use. In addition, it is absolutely necessary to raise public awareness of the importance of these precious resources in the treatment of these toxicities in ruminants. Finally, studies must also be oriented towards the integration of these natural products in the diet intended for endemic regions to facilitate their uses and applications.

ORCID 

Abdellatif Rahim: [0000-0001-5832-1710](https://orcid.org/0000-0001-5832-1710)

Fatma Aydogmus-Öztürk: [0000-0001-6070-7452](https://orcid.org/0000-0001-6070-7452)

Cansel Cakir: [0000-0002-6175-9008](https://orcid.org/0000-0002-6175-9008)

Abdelkhalid Essamadi: [0000-0003-4537-1684](https://orcid.org/0000-0003-4537-1684)

Bouchra El Amiri: [0000-0003-3443-5988](https://orcid.org/0000-0003-3443-5988)

## References

- [1] P.K. Thornton (2010). Livestock production: recent trends, future prospects, *Phil. Trans. R. Soc. B* **365**, 2853–2867.
- [2] F.J.C. Swanepoel, A. Stroebe and S. Moyo (2010). The role of livestock in developing communities: enhancing multifunctionality, International Livestock Research Institute (ILRI), University of the Free State, CTA Wageningen, The Netherlands.
- [3] A. Mottet, C. de Haan, A. Falcucci, G. Tempio, C. Opio and P. Gerber (2017). Livestock: On our plates or eating at our table? A new analysis of the feed/food debate, *Glob. Food Sec.* **14**, 1–8.
- [4] A. Sandhage-Hofmann (2016). Rangeland management, *Reference Module In Earth Systems And Environmental Sciences*, Elsevier, University of Bonn, Bonn, Germany.
- [5] S.M. Gwaltney-Brant (2018). Epidemiology of Animal Poisonings in the United States, In: Veterinary toxicology, Basic and clinical principles, ed: R.C. Gupta, Academic Press, United States, pp. 37-44.
- [6] A. Bhargava, F. F., Carmona, M., Bhargava and S. Srivastava (2012). Approaches for enhanced phytoextraction of heavy metals. *The J. Environ. Managm.* **105**, 103-120.

- [7] P. Samal, R.C. Patra, A.R. Gupta, S.K. Mishra, D. Jena and D. Satapathy (2016). Effect of *Tamarindus indica* leaf powder on plasma concentrations of copper, zinc, and iron in fluorotic cows, *Vet. World* **9**, 1121–1124.
- [8] A. Rahim, A. Essamadi and B. El Amiri (2022). A comprehensive review on endemic and experimental fluorosis in sheep: Its diverse effects and prevention, *Toxicology* **465**, 153025.
- [9] M.J. Kosnett (2013). The role of chelation in the treatment of arsenic and mercury poisoning, *J. Med. Toxicol.* **9**, 347–354.
- [10] G.B. van der Voet, O. Schijns and F.A. de Wolff (1999). Fluoride enhances the effect of aluminium chloride on interconnections between aggregates of hippocampal neurons, *Arch. Physiol. Biochem.* **107**, 15–21.
- [11] J. Wei, X. Zheng, J. Liu, G. Zhang, Y. Zhang, C. Wang and Y. Liu (2021). The levels, sources, and spatial distribution of heavy metals in soils from the drinking water sources of Beijing, China, *Sustainability* **13**, 3719.
- [12] V. Masindi and K.L. Muedi (2018). Environmental contamination by heavy metals. Heavy Metals, IntechOpen. <https://doi.org/10.5772/intechopen.76082>.
- [13] M. Jin, H. Yuan, B. Liu, J. Peng, L. Xu and D. Yang (2020). Review of the distribution and detection methods of heavy metals in the environment, *Anal. Methods* **12**, 5747–5766.
- [14] G.P.S. Sidhu (2016). Heavy metal toxicity in soils: sources, remediation technologies and challenges, *Adv. Plants Agric. Res.* **5**, 445-446.
- [15] P.B. Tchounwou, C.G. Yedjou, A.K. Patlolla and D.J. Sutton (2012). Heavy metals toxicity and the environment, In: Molecular, Clinical and Environmental Toxicology, ed: A. Luch, Experientia Supplementum, Springer, Basel, pp.133–164. [https://doi.org/10.1007/978-3-7643-8340-4\\_6](https://doi.org/10.1007/978-3-7643-8340-4_6).
- [16] Y. Xie, J. Fan, W. Zhu, E. Amombo, Y. Lou, L. Chen and J. Fu (2016). Effect of heavy metals pollution on soil microbial diversity and *Bermudagrass* genetic variation, *Front. Plant Sci.* **7**, 755.
- [17] U. Okerefor, M. Makhatha, L. Mekuto, N. Uche-Okerefor, T. Sebola and V. Mavumengwana (2020). Toxic metal implications on agricultural soils, plants, animals, aquatic life and human health, *Int. J. Environ. Res. Public Health* **17**, 2204.
- [18] J. Musilová, J. Árvay, A. Vollmannova, T. Tóth and J. Tomáš (2016). Environmental contamination by heavy metals in region with previous mining activity, *Bull. Environ. Contam. Toxicol.* **97**, 569-75.
- [19] J. Briffa, E. Sinagra and R. Blundell (2020). Heavy metal pollution in the environment and their toxicological effects on humans, *Heliyon* **6**, e04691.
- [20] S. Gautam, L.K. Pandey, V. Vinayak and A. Arya (2017). Morphological and physiological alterations in the diatom *Gomphonema pseudoaugur* due to heavy metal stress, *Ecol. Indic.* **72**, 67–76.
- [21] L.N. Suvarapu and S.O. Baek (2017). Determination of heavy metals in the ambient atmosphere: A review, *Toxicol. Ind. Health* **33**, 79–96.
- [22] M.G. García and L. Borgnino (2015). CHAPTER 1 Fluoride in the context of the environment, In: Fluorine: Chemistry, Analysis, Function and Effects, ed: Victor R Preedy, pp.3–21.
- [23] E. Perumal, V. Paul, V. Govindarajan and L. Panneerselvam (2013). A brief review on experimental fluorosis, *Toxicol. Lett.* **223**, 236–251.
- [24] R. Ranjan and A. Ranjan (2015). Fluoride toxicity in animals. Springer Cham.
- [25] S.L. Choubisa, G.V. Mishra, Z. Sheikh, B. Bhardwaj, P. Mali and V.J. Jaroli (2011). Food, fluoride, and fluorosis in domestic ruminants in the Dungarpur district of Rajasthan, India, *Fluoride* **7**, 70-76.
- [26] F. Urut, S. Dede, V. Yuksek, S. Cetin, A. Usta and M. Taspinar (2021). In vitro evaluation of the apoptotic, autophagic, and necrotic molecular pathways of fluoride, *Biol. Trace Elem. Res.* **199**, 3700–3706.
- [27] J.L. Shupe and A.E. Olson (1971). Clinical aspects of fluorosis in horses, *J. Am. Vet. Med. Assoc.* **158**, 167-74.
- [28] A.K. Susheela and M. Bhatnagar (1993). Fluoride toxicity: a biochemical and scanning electron microscopic study of enamel surface of rabbit teeth, *Arch. Toxicol.* **67**, 573–579.
- [29] P. Ekambaram and V. Paul (2002). Modulation of fluoride toxicity in rats by calcium carbonate and by withdrawal of fluoride exposure, *Pharmacol. Toxicol.* **90**, 53–58.
- [30] B. Cadir, M. Kürkcü, Í.A. Oz and M.E. Benlidayi (2009). Effects of vitamin K1 on fluoride-induced bone changes in growing rats: a histomorphometric and radiodensitometric study, *Arch. Oral Biol.* **54**, 512–517.
- [31] C.H. Turner, I. Owan, E.J. Brizendine, W. Zhang, M.E. Wilson and A.J. Dunipace (1996). High fluoride intakes cause osteomalacia and diminished bone strength in rats with renal deficiency, *Bone* **19**, 595–601.
- [32] P.M. Basha, P. Rai and S. Begum (2011). Evaluation of fluoride-induced oxidative stress in rat brain: a multigeneration study, *Biol. Trace Elem. Res.* **142**, 623–637.
- [33] D. Giri, R.C. Ghosh, D. Kashyap, G. Dewangan and S.K. Maiti (2015). Haemato-biochemical alterations in subacute oral toxicity of sodium fluoride in wistar rats, *J. Anim. Res.* **5**, 595-598.

## Mitigating fluoride, lead, arsenic and cadmium toxicities

- [34] A. Elbetieha, H. Darmani and A. Al-Hiyasat (2000). Fertility effect of sodium fluoride in male mice, *Fluoride* **33**, 128–134.
- [35] R.S. Gupta, T.I. Khan, D. Agrawal and J.B.S. Kachhawa (2007). The toxic effects of sodium fluoride on the reproductive system of male rats, *Toxicol. Ind. Health* **23**, 507–513.
- [36] M. Kessabi, A. Hamliri, J.P. Braun and A.G. Rico (1985). Experimental acute sodium fluoride poisoning in sheep: renal, hepatic, and metabolic effects, *Fundam Appl. Toxicol.* **5**, 1025–1033.
- [37] S. Choubisa (2013). Fluoride toxicosis in immature herbivorous domestic animals living in low fluoride water endemic areas of Rajasthan, India: an observational survey, *Fluoride* **46**, 19–24.
- [38] M. Kessabi, M. Khouzaimi, J.P. Braun and A. Hamliri (1983). Serum biochemical effects of fluoride on cattle in the *Darmous area*, *Vet. Hum. Toxicol.* **25**, 403–406.
- [39] A. Bennis, M. Kessabi, A. Hamliri, F. Farge and J.P. Braun (1993). Plasma biochemistry of adult goats with chronic fluoride poisoning in Morocco, *Fluoride* **26**, 241–246.
- [40] F. Yur, N. Mert, S. Dede, Y. Değer, A. Ertekin, H. Mert, S. Yasar, I. Dogan and A. Isik (2013). Evaluation of serum lipoprotein and tissue antioxidant levels in sheep with fluorosis, *Fluoride* **46**, 90–96.
- [41] N. Aydin, S. Dede and P. Tanritanir (2014). The distribution of minerals in some tissues of sheep with fluorosis, *Fluoride* **47**, 43–48.
- [42] M.J.K. Simon, F.T. Beil, W. Rüther, B. Busse, T. Koehne, M. Steiner, P. Pogoda, A. Ignatius, M. Amling and R. Oheim (2014). High fluoride and low calcium levels in drinking water is associated with low bone mass, reduced bone quality and fragility fractures in sheep, *Osteoporos. Int.* **25**, 1891–1903.
- [43] C.J. Botha, T.W. Naude, P.P. Minnaar, S.R. Van Amstel and S.D. Janse van Rensburg (1993). Two outbreaks of fluorosis in cattle and sheep, *J. S. Afr. Vet. Assoc.* **64**, 165–168.
- [44] S. Çetin and F. Yur (2016). Levels of trace elements in muscle and kidney tissues of sheep with fluorosis, *Biol. Trace Elem. Res.* **174**, 82–84.
- [45] H.C. Kim, T.W. Jang, H.J. Chae, W.J. Choi, M.N. Ha, B.J. Ye, B.G. Kim, M.J. Jeon, S.Y. Kim and Y.S. Hong (2015). Evaluation and management of lead exposure, *Ann. Occup. Environ. Med.* **27**, 1–9.
- [46] D. Swarup, R. Naresh, V.P. Varshney, M. Balagangatharathilagar, P. Kumar, D. Nandi and R.C. Patra (2007). Changes in plasma hormones profile and liver function in cows naturally exposed to lead and cadmium around different industrial areas, *Res. Vet. Sci.* **82**, 16–21.
- [47] R.C. Patra, D. Swarup, M.C. Sharma and R. Naresh (2006). Trace mineral profile in blood and hair from cattle environmentally exposed to lead and cadmium around different industrial units, *J. Vet. Med. A Physiol. Pathol. Clin. Med.* **53**, 511–517.
- [48] L. Reis, P. Pardo, A. Camargos and E. Oba (2010). Mineral element and heavy metal poisoning in animals, *Afr. J. Med. Med. Sci.* **1**, 560–579.
- [49] E. Obeng-Gyasi (2019). Sources of lead exposure in various countries, *Environ. Health Rev.* **34**, 25–34.
- [50] M.A. Assi, M.N.M. Hezmee, A.W. Haron, M.Y.M. Sabri and M.A. Rajion (2016). The detrimental effects of lead on human and animal health, *Vet. World* **9**, 660–671.
- [51] M. Miranda, M. López-Alonso, P. García-Partida, J. Velasco and J.L. Benedito (2006). Long-term follow-up of blood lead levels and haematological and biochemical parameters in heifers that survived an accidental lead poisoning episode, *J. Vet. Med. A Physiol. Pathol. Clin. Med.* **53**, 305–310.
- [52] C. Waldner, S. Checkley, B. Blakley, C. Pollock and B. Mitchell (2002). Managing lead exposure and toxicity in cow-calf herds to minimize the potential for food residues, *J. Vet. Diagn. Invest.* **14**, 481–486.
- [53] G. Flora, D. Gupta and A. Tiwari (2012). Toxicity of lead: A review with recent updates, *Interdiscip. Toxicol.* **5**, 47–58.
- [54] M.K. Singh, S. Dwivedi, S.S. Yadav, P. Sharma, S. Khattri (2014). Arsenic-induced hepatic toxicity and its attenuation by fruit extract of *Embllica officinalis* (Amla) in mice, *Indian J. Clin. Biochem.* **29**, 29–37.
- [55] W.H. El-Tantawy (2016). Antioxidant effects of *Spirulina* supplement against lead acetate-induced hepatic injury in rats, *J. Tradit. Complement. Med.* **6**, 327–331.
- [56] D. Bagchi and H.G. Preuss (2005). Effects of acute and chronic oval exposure of lead on blood pressure and bone mineral density in rats, *J. Inorg. Biochem.* **99**, 1155–1164.
- [57] R.A.R. Elgawish and H.M.A. Abdelrazek (2014). Effects of lead acetate on testicular function and caspase-3 expression with respect to the protective effect of cinnamon in albino rats, *Toxicol. Rep.* **1**, 795–801.
- [58] A.C.B.A. Lopes, T.S. Peixe, A.E. Mesas and M.M.B. Paoliello (2016). Lead exposure and oxidative stress: a systematic review, *Rev. Environ. Contam. Toxicol.* **236**, 193–238.
- [59] A.L. Wani, A. Ara and J.A. Usmani (2015). Lead toxicity: a review, *Interdiscip. Toxicol.* **8**, 55–64.
- [60] J. Payne and C. Livesey (2010). Lead poisoning in cattle and sheep, *In Practice* **32**, 64–69.
- [61] O.M. Radostits, C.C. Gay, K.W. Hinchcliff, and P.D. Constable (2006). Veterinary medicine: a textbook of the diseases of cattle, horses, sheep, pigs and goats, Elsevier Saunders.

- [62] M.E. Gilbert, M.E. Kelly, T.E. Samsam and J.H. Goodman (2005). Chronic developmental lead exposure reduces neurogenesis in adult rat hippocampus but does not impair spatial learning, *Toxicol. Sci.* **86**, 365–374.
- [63] A.M. Soliman, E.A. Abu-Basha, S.A.H. Youssef, A.M. Amer, P.A. Murphy, C.C. Hauck, R. Gehring and W.H. Hsu (2013). Effect of chronic lead intoxication on the distribution and elimination of amoxicillin in goats, *J. Vet. Sci.* **14**, 395–403.
- [64] M.A. El-Ghiaty and A.O.S. El-Kadi (2021). Arsenic: various species with different effects on cytochrome P450 regulation in humans, *Excli. J.* **20**, 1184–1242.
- [65] R. Kapp (2016). Arsenic: toxicology and health effects, In: *Encyclopedia of Food and Health*, ed: B. Caballero, P.M. Finglas and F. Toldra, Elsevier Publishing, pp.256-265.
- [66] S.H. Frisbie, R. Ortega, D.M. Maynard and B. Sarkar (2002). The concentrations of arsenic and other toxic elements in Bangladesh's drinking water, *Environ. Health Perspect.* **110**, 1147–1153.
- [67] I. Herath, M. Vithanage, J. Bundschuh, J.P. Maity and P. Bhattacharya (2016). Natural arsenic in global groundwaters: distribution and geochemical triggers for mobilization, *Curr. Pollution Rep.* **2**, 68–89.
- [68] K.C.M. Kwok, L.F. Koong, G. Chen and G. McKay (2014). Mechanism of arsenic removal using chitosan and nanochitosan, *J. Colloid Interface Sci.* **416**, 1–10.
- [69] J.P. Wang, L. Qi, M.R. Moore and J.C. Ng (2002). A review of animal models for the study of arsenic carcinogenesis, *Toxicol. Lett.* **133**, 17–31.
- [70] M.R. Garry, A.B. Santamaria, A.L. Williams and J.M. DeSesso (2015). *In utero* arsenic exposure in mice and early life susceptibility to cancer, *Regul. Toxicol. Pharmacol.* **73**, 378–390.
- [71] K. Tewari and D. Jhala (2004). Fluoride and/or arsenic toxicity in mice testis with formation of giant cells and subsequent recovery by some antidotes, *Fluoride* **37**, 172–184.
- [72] Y.J. Kim and J.M. Kim (2015). Arsenic toxicity in male reproduction and development, *J. Reprod. Dev.* **19**, 167–180.
- [73] A. Santra, A. Maiti, S. Das, S. Lahiri, S.K. Charkaborty and D.N. Mazumder (2000). Hepatic damage caused by chronic arsenic toxicity in experimental animals, *J. Toxicol. Clin. Toxicol.* **38**, 395–405.
- [74] R. Ferzand, J.A. Gadahi, S. Saleha and Q. Ali (2008). Histological and haematological disturbance caused by arsenic toxicity in mice model, *Pak. J. Biol. Sci.* **11**, 1405–1413.
- [75] J. Ashrafihelan, J.S. Amoli, M. Alamdari, T.A. Esfahani, M. Mozafari, A.R. Nourian and A.A. Bahari (2013). Arsenic toxicosis in sheep: The first report from Iran, *Interdiscip. Toxicol.* **6**, 93–98.
- [76] C. Maji, S. Sarkar, S. Biswas, P.H. Patra, B.K. Datta, S. Bandyopadhyay, T.K. Biswas, C. Jana and T.K. Mandal (2016). Experimental assessment of arsenic toxicity in garole sheep in India, *Emerg. Contam.* **2**, 128–134.
- [77] L.A. Selby, A.A. Case, G.D. Osweiler and H.M. Hayes (1977). Epidemiology and toxicology of arsenic poisoning in domestic animals, *Environ. Health Perspect.* **19**, 183–189.
- [78] T. Rana, A.K. Bera, S. Das, D. Bhattacharya, S. Bandyopadhyay, D. Pan and S. Kumar Das (2010). Effect of chronic intake of arsenic-contaminated water on blood oxidative stress indices in cattle in an arsenic-affected zone, *Ecotoxicol. Environ. Saf.* **73**, 1327–1332.
- [79] P. Loganathan, S. Vigneswaran, J. Kandasamy and R. Naidu (2012). Cadmium sorption and desorption in soils: a review, *Crit. Rev. Environ. Sci. Technol.* **42**, 489–533.
- [80] L. Patrick (2003). Toxic metals and antioxidants: Part II the role of antioxidants in arsenic and cadmium toxicity, *Altern. Med. Rev.* **8**, 106–128.
- [81] Y. Li, Y. Wang, X. Shen and F. Liu (2021). The combinations of sulfur and molybdenum fertilizations improved antioxidant capacity of grazing Guizhou semi-fine wool sheep under copper and cadmium stress, *Ecotoxicol. Environ. Saf.* **222**, 112520.
- [82] M.J. Canty, S. McCormack, E.A. Lane, D.M. Collins and S.J. More (2011). Essential elements and heavy metal concentrations in a small area of the Castlecomer Plateau, Co. Kilkenny, Ireland: Implications for animal performance, *Irish J. Agric. Food Res.* **50**, 223–238.
- [83] M.J. Canty, A. Scanlon, D.M. Collins, G. McGrath, T.A. Clegg, E. Lane, M.K. Sheridan and S.J. More (2014). Cadmium and other heavy metal concentrations in bovine kidneys in the Republic of Ireland, *Sci. Total Environ.* **485**, 223–231.
- [84] N. Pant, R. Kumar, R.C. Murthy and S.P. Srivastava (2001). Male reproductive effect of arsenic in mice, *Biometals.* **14**, 113–117.
- [85] J.T. Ayodele and F.B. Suleiman (2004). Determination of iodine in human milk and urine, *Ife J. Sci.* **6**, 79–83.
- [86] S. Roggeman, G. de Boeck, H. De Cock, R. Blust and L. Bervoets (2014). Accumulation and detoxification of metals and arsenic in tissues of cattle (*Bos taurus*), and the risks for human consumption, *Sci. Total Environ.* **466**, 175–184.
- [87] E.A. Lane, M.J. Canty and S.J. More (2015). Cadmium exposure and consequence for the health and productivity of farmed ruminants, *Res. Vet. Sci.* **101**, 132–139.

## Mitigating fluoride, lead, arsenic and cadmium toxicities

- [88] M. Anke, T. Masaoka, B. Groppel, G. Zervas and W. Arnhold (1989). The influence of sulphur, molybdenum and cadmium exposure on the growth of goat, cattle and pig, *Arch. Tierernahr.* **39**, 221–228.
- [89] I.A. Kreis, M. de Does, J.A. Hoekstra, C. de Lezenne Coulander, P.W. Peters and G.H. Wentink (1993). Effects of cadmium on reproduction, an epizootologic study, *Teratology* **48**, 189–196.
- [90] S.H. Prankel, R.M. Nixon and C.J.C. Phillips (2004). Meta-analysis of feeding trials investigating cadmium accumulation in the livers and kidneys of sheep, *Environ. Res.* **94**, 171–183.
- [91] J.M. Wilkinson, J. Hill and C.J.C. Phillips (2003). The accumulation of potentially-toxic metals by grazing ruminants, *Proc. Nutr. Soc.* **62**, 267–277.
- [92] S.A. Gunn, T.C. Gould and W.A. Anderson (1963). Cadmium-induced interstitial cell tumors in rats and mice and their prevention by zinc, *J. Natl. Cancer Inst.* **31**, 745–759.
- [93] E. Löser (1980). A 2-year oral carcinogenicity study with cadmium on rats, *Cancer Lett.* **9**, 191–198.
- [94] M.D. Johnson, N. Kenney, A. Stoica, L. Hilakivi-Clarke, B. Singh, G. Chepko, R. Clarke, P.F. Sholler, A.A. Lirio, C. Foss, R. Reiter, B. Trock, S. Paik and M.B. Martin (2003). Cadmium mimics the *in vivo* effects of estrogen in the uterus and mammary gland, *Nat. Med.* **9**, 1081–1084.
- [95] Q. Zhu and X. Li, R.S. Ge (2020). Toxicological effects of cadmium on mammalian testis, *Front. Genet.* **11**, 527.
- [96] R.P. Leite, F.S. Predes, J.C. Monteiro, K.M. Freitas, R.S. Wada and H. Dolder (2013). Advantage of Guaraná (*Paullinia cupana* Mart) supplementation on cadmium-induced damages in testis of adult Wistar rats, *Toxicol. Pathol.* **41**, 73–79.
- [97] S. Amara, H. Abdelmelek, C. Garrel, P. Guiraud, T. Douki, J.L. Ravanat, A. Favier, M. Sakly and K. Ben Rhouma (2008). Preventive effect of zinc against cadmium-induced oxidative stress in the rat testis, *J. Reprod. Dev.* **54**, 129–134.
- [98] R. Mahmoudi, A. Azizi, S. Abedini, V. Hemayatkhah Jahromi, H. Abidi and M. Jafari Barmak (2018). Green tea improves rat sperm quality and reduced cadmium chloride damage effect in spermatogenesis cycle, *J. Med. Life* **11**, 371–380.
- [99] S. Saygi, G. Deniz, O. Kutsal and N. Vural (1991). Chronic effects of cadmium on kidney, liver, testis, and fertility of male rats, *Biol. Trace Elem. Res.* **31**, 209–214.
- [100] T. Hussain, G.S. Shukla and S.V. Chandra (1987). Effects of cadmium on superoxide dismutase and lipid peroxidation in liver and kidney of growing rats: *in vivo* and *in vitro* studies, *Pharmacol. Toxicol.* **60**, 355–358.
- [101] Z. Wang, Y. Sun, W. Yao, Q. Ba and H. Wang (2021). Effects of cadmium exposure on the immune system and immunoregulation, *Front Immunol.* **12**, 695484.
- [102] C.O.B. Okoye, J.N. Ugwu and J.N. Ugwu (2010). Impact of environmental cadmium, lead, copper and zinc on quality of goat meat in Nigeria, *Bull. Chem. Soc. Ethiop.* **24**, 133–138.
- [103] A. Sedki, N. Lekouch, S. Gamon and A. Pineau (2003). Toxic and essential trace metals in muscle, liver and kidney of bovines from a polluted area of Morocco, *Sci. Total Environ.* **317**, 201–205.
- [104] M. Balali-Mood, K. Naseri, Z. Tahergorabi, M.R. Khazdair and M. Sadeghi (2021). Toxic mechanisms of five heavy metals: mercury, lead, chromium, cadmium, and arsenic, *Front. Pharmacol.* **12**, 643972.
- [105] G.A. Engwa, P.U. Ferdinand, F.N. Nwalo and M. Unachukwu (2019). Mechanism and health effects of heavy metal toxicity in humans, *Poisoning in The Modern World-New Tricks For An Old Dog*, IntechOpen. <https://doi.org/10.5772/intechopen.82511>.
- [106] S. Park, K. Ajtai and T.P. Burghardt (1999). Inhibition of myosin ATPase by metal fluoride complexes, *Biochim. Biophys. Acta- Protein Struct. Mol. Enzym.* **1430**, 127–140.
- [107] N. Burford, M.D. Eelman and K. Groom (2005). Identification of complexes containing glutathione with As(III), Sb(III), Cd(II), Hg(II), Tl(I), Pb(II) or Bi(III) by electrospray ionization mass spectrometry, *J. Inorg. Biochem.* **99**, 1992–1997.
- [108] P. Faller, K. Kienzler and A. Krieger-Liszkay (2005). Mechanism of Cd<sup>2+</sup> toxicity: Cd<sup>2+</sup> inhibits photoactivation of Photosystem II by competitive binding to the essential Ca<sup>2+</sup> site, *Biochim. Biophys. Acta - Bioenerg.* **1706**, 158–164.
- [109] X. Wang, A.K. Mandal, H. Saito, J.F. Pulliam, E.Y. Lee, Z.J. Ke, J. Lu, S. Ding, L. Li, B.J. Shelton, T. Tucker, B.M. Evers, Z. Zhang and X. Shi (2012). Arsenic and chromium in drinking water promote tumorigenesis in a mouse colitis-associated colorectal cancer model and the potential mechanism is ROS-mediated Wnt/β-catenin signaling pathway, *Toxicol. Appl. Pharmacol.* **262**, 11–21.
- [110] A. Rahim, C. Çakir, M. Ozturk, B. Şahin, A. Soulaïmani, M. Sibaoeuih, B. Nasser, R. Eddoha, A. Essamadi and B. El Amiri (2021). Chemical characterization and nutritional value of *Spirulina platensis* cultivated in natural conditions of Chichaoua region (Morocco), *S. Afr. J. Bot.* **141**, 235–242.
- [111] Q. Zhai, A. Narbad and W. Chen (2015). Dietary strategies for the treatment of cadmium and lead toxicity, *Nutrients* **7**, 552–571.

- [112] R. Ranjan, D. Swarup, R.C. Patra and V. Chandra (2009). *Tamarindus indica* L. and *Moringa oleifera* M. extract administration ameliorates fluoride toxicity in rabbits, *Indian J. Exp. Biol.* **47**, 900–905.
- [113] E. Perumal, T. Namitha, S. Bhuvaneswari, A. Subramaniam, V. Dhakshinamoorthy and M. Coimbatore (2010). Therapeutic efficacy of *Tamarindus indica* (L) to protect against fluoride-induced oxidative stress in the liver of female rats, *Fluoride* **43**, 134.
- [114] M. Sinha, P. Manna and P.C. Sil (2007). Aqueous extract of the bark of *Terminalia arjuna* plays a protective role against sodium-fluoride-induced hepatic and renal oxidative stress, *J. Nat. Med.* **61**, 251–260.
- [115] R.A. Vasant and A.V.R.L. Narasimhacharya (2011). Amelioration of fluoride induced oxidative stress by *Mangifera indica* L fruit, *Spatula DD* **1**, 181–188.
- [116] C. Maji, S. Biswas, P.K. Sarkar, P.H. Patra, S. Bandyopadhyay, S. Sarkar, A. Chattopadhyay and T.K. Mandal (2020). Evaluation of ameliorative effect of two selected plant drugs on experimentally induced arsenic toxicity in sheep, *Environ. Sci. Pollut. Res. Int.* **27**, 36744–36753.
- [117] D. Banji, O. Banji, N. Pratusha and A.R. Annamalai (2013). Investigation on the role of *Spirulina platensis* in ameliorating behavioural changes, thyroid dysfunction and oxidative stress in offspring of pregnant rats exposed to fluoride, *Food Chem.* **140**, 321–31.
- [118] T.M. da Silva, B.M. da Fonseca, A.L.L.S. Sales, P. Holleben, M.C. Valera and M.A.M. de Araújo (2016). Effects of fluoride and *Aloe vera* Tooh gel in artificial white spot lesions *in vitro*, *RGO, Rev. Gaúch. Odontol* **64**, 56–61.
- [119] S. Dey, A. Gupta, D.Swarup, M. Saini, A. Saxena and A. Dan (2013). Ameliorative effect of *Tamarindus indica* L. on biochemical parameters of serum and urine in cattle from fluoride endemic area, *Vet. Arh.* **83**, 487–496.
- [120] V. Najamezhad, M.R. Aslani, M. Balali-Mood (2008). Evaluation of allicin for the treatment of experimentally induced subacute lead poisoning in sheep, *Biol. Trace Elem. Res.* **126**, 141–7.
- [121] K. Badiei, K. Mostaghni, A. Nowrooziasl and A. Tabatabaei Naeini (2006). Ameliorated effects of *Allium Sativum* on subclinical lead toxicity in goats, *Pak. Vet. J.* **26**, 184–186.
- [122] M.F.F. Pérez Díaz, M. Acosta, F.H. Mohamed, M.L. Ferramola, L.B. Oliveros and M.S. Gimenez (2013). Protective effect of soybeans as protein source in the diet against cadmium-aorta redox and morphological alteration, *Toxicol. Appl. Pharmacol.* **272**, 806–815.
- [123] B.E. Oyinloye, A.F. Adenowo, F.O. Osunsanmi, B.I. Ogunyinka, S.O. Nwozo, A.P. Kappo (2016). Aqueous extract of *Monodora myristica* ameliorates cadmium-induced hepatotoxicity in male rats, *SpringerPlus* **5**, 641.
- [124] E. Moghtadai Khorasgani, A. Ghasemi Pirbalouti and S. Adibi (2015). Histopathological evidences for effect beneficial of *Satureja hortensis* extract on hepatic lesion by cadmium–induced in Rat, *J. Med Herb.* **6**, 101–107.
- [125] M. Chopra, S. Rani and V. Sharma (2013). Protective effect of *Withania somnifera* and vitamin E against cadmium induced toxicity in mice spleen, *Int. J. Pharmacol. Toxicol. Sci.* **3**, 32–42.
- [126] A. Kumar, R. Kumar, M.S. Rahman, M.A. Iqubal, G. Anand, P.K. Niraj and M. Ali (2015). Phytoremedial effect of *Withania somnifera* against arsenic-induced testicular toxicity in Charles Foster rats, *Avicenna J. Phytomed* **5**, 355–364.
- [127] T.K. Dua, S. Dewanjee, M. Gangopadhyay, R. Khanra, M. Zia-Ul-Haq and V. De Feo (2015). Ameliorative effect of water spinach, *Ipomea aquatica* (Convolvulaceae), against experimentally induced arsenic toxicity, *J. Transl. Med.* **13**, 81.
- [128] M. Roy, S. SushovanRoy, B. Roopali, H.K. Ratre, N. Singh, J. Singh and R. Pathak (2020). Ameliorative potential of *Tamarindus indica* seed coat against arsenic-induced hepatotoxicity in wistar rats, *J. Entomol. Zool. Stud.* **8**, 1125-1129.
- [129] R.S. Almeer, G.I. AlBasher, S. Alarifi, S. Alkahtani, D. Ali and A.E. Abdel Moneim (2019). Royal jelly attenuates cadmium-induced nephrotoxicity in male mice, *Sci. Rep.* **9**, 1-12.
- [130] S.H. Negm and S.S. El-Soadaa (2020). Effect of *Terminalia chebula* on cadmium-induced nephrotoxicity and lipid profiles in rats, *Biosci. Res.* **17**, 1535-1544.
- [131] J.M. Hazarika, P.K. Sarkar, A. Chattopadhyay, T.K. Mandal and S. Sarkar (2015). Evaluation of some selected herbs on arsenic-affected cattle in Nadia District, West Bengal, India, *Environ. Sci. Pollut. Res.* **22**, 4942-4948
- [132] S. Biswas, C. Maji, P.K. Sarkar, S. Sarkar, A. Chattopadhyay and T.K. Mandal (2016). Ameliorative effect of two Ayurvedic herbs on experimentally induced arsenic toxicity in calves, *J. Ethnopharmacol.* **197**, 266-273.
- [133] N. Kumar, S. Sood, B. Arora and M. Singh, Beena (2010). Effect of duration of fluoride exposure on the reproductive system in male rabbits, *J. Hum. Reprod. Sci.* **3**, 148–152.
- [134] S. Aggarwal (2000). Histopathology of fluoride-induced hepatotoxicity in rabbits, *Sci. Rep.* **34**, 34–42.

## Mitigating fluoride, lead, arsenic and cadmium toxicities

- [135] A.E. Azab, M.O. Albasha, J.M. Jbireal and A.A. Adwas (2018). Sodium fluoride induces hepato-renal oxidative stress and pathophysiological changes in experimental animals, *Open J. Apoptosis* **7**, 1-23.
- [136] M.P. Basha and N.S. Sujitha (2011). Chronic fluoride toxicity and myocardial damage: antioxidant offered protection in second generation rats, *Toxicol. Int.* **18**, 99–104.
- [137] F. Aziz, I. Maulood and M.A.H. Chawsheen (2012). Effects of melatonin, vitamin C and E alone or in combination on lead-induced injury in liver and kidney organs of rats, *IOSR J. Pharm.* **2**, 13-18.
- [138] A. Lynda, M. Hamadouche, A. Touabti and S. Khennouf (2011). Effect of long-term exposure to low or moderate lead concentrations on growth, lipid profile and liver function in albino rats, *Adv. Biol. Res.* **5**, 339–347.
- [139] S. Srivastava, E.N. Vladykovskaya, P. Haberzettl, S.D. Sithu, S.E. D'Souza and J.C. States (2009). Arsenic exacerbates atherosclerotic lesion formation and inflammation in ApoE<sup>-/-</sup> mice, *Toxicol. Appl. Pharmacol.* **241**, 90–100.
- [140] R. Rathor, D. Dubey and G. Kannan (2007). Concomitant administration of Moringa oleifera seed powder in the remediation of arsenic-induced oxidative stress in mouse, *Cell Biol. Int.* **31**, 44–56.
- [141] M. Sarkar, G.R. Chaudhuri, A. Chattopadhyay and N.M. Biswas (2003). Effect of sodium arsenite on spermatogenesis, plasma gonadotrophins and testosterone in rats, *Asian J. Androl.* **5**, 27–31.
- [142] D.G. Stump, J.F. Holson, T.L. Fleeman, M.D. Nemec and C.H. Farr (1999). Comparative effects of single intraperitoneal or oral doses of sodium arsenate or arsenic trioxide during in utero development, *Teratology* **60**, 283–291.
- [143] M.P. Waalkes, T.P. Coogan and R.A. Barter (1992). Toxicological principles of metal carcinogenesis with special emphasis on cadmium, *Crit. Rev. Toxicol.* **22**, 175–201.
- [144] U.N. Samal and B.N. Naik (2009). The fluorosis problem in tropical sheep, *Fluoride* **25**, 183-190.
- [145] M. Kessabi, B. Assimi and J.P. Braun (1984). The effects of fluoride on animals and plants in the South Safi zone, *Sci. Total Environ.* **38**, 63–68.
- [146] V. Kant, A. Srivastava, P. Verma and S. Uppal (2009). Ameliorative effect of aluminium sulphate on the milk fluoride levels in goats, *Isr. J. Vet. Med.* **64**, 45-46.
- [147] O.M. Radostits, C.C. Gay, D.C. Blood, K.W. Hinchcliff, J.H. Arundel, D.E. Jacobs, and C. Valledor Martínez (2000). Bovine mastitis: In: Veterinary Medicine A Textbook of the Diseases of Cattle, Sheep, Pigs, Goats and Horses. 9th Edition, W.B. Saunders Company Ltd., London.
- [148] P.H. Patra, S. Bandyopadhyay, M.C. Bandyopadhyay and T.K. Mandal (2013). Immunotoxic and genotoxic potential of arsenic and its chemical species in goats, *Toxicol. Int.* **20**, 6–10.
- [149] B. Keshavarzi, A. Seradj, Z. Akbari, F. Moore, A. Rahmani Shahraki and M. Pourjafar (2015). Chronic arsenic toxicity in sheep of kurdistan province, western Iran, *Arch. Environ. Contam. Toxicol.* **69**, 44-53.
- [150] U. Biswas, S. Sarkar, M.K. Bhowmik, A.K. Samanta and S. Biswas (2000). Chronic toxicity of arsenic in goats: clinicobiochemical changes, pathomorphology and tissue residues, *Small Rumin Res.* **38**, 229–235.
- [151] N.B. Ortega-Morales, J.A. Cueto-Wong, E. Barrientos-Juárez, G. García-Vargas, H. Salinas-González, A. Buendía Garcia and J. Morán-Martínez (2020). Toxicity in goats exposed to arsenic in the region Lagunera, Northern Mexico, *Vet. Sci.* **7**, 59.
- [152] B.K. Biswas, R.K. Dhar, G. Samanta, B.K. Mandal, D. Chakraborti, I. Faruk, K.S. Islam, M.M. Chowdhury, A. Islam and S. Roy (1998). Detailed study report of samta, one of the arsenic-affected villages of Jessore district, Bangladesh, *Curr. Sci.* **74**, 134–145.
- [153] R.A. Vasant and A.V. Narasimhacharya (2012). Alleviatory effects of *Embllica officinalis* g. as a food supplement in fluoride induced hyperlipemia and oxidative stress, *Int. J. Pharm. Pharm. Sci.* **4** (1), 404-408.
- [154] S. Dey, D. Swarup, A. Saxena and A. Dan (2011). In vivo efficacy of tamarind (*Tamarindus indica*) fruit extract on experimental fluoride exposure in rats, *Res. Vet. Sci.* **91**, 422–425.
- [155] E.O. Salawu, O.A. Adeeyo, O.P. Falokun, U.A. Yusuf, A. Oyerinde and A.A. Adeleke (2009). Tomato (*Lycopersicon esculentum*) prevents lead-induced testicular toxicity, *J. Hum. Reprod. Sci.* **2**, 30-4.
- [156] E.S. Mehana, A. Meki and K. Fazili (2010). Ameliorated effects of green tea extract on lead induced liver toxicity in rats, *Exp. Toxicol. Pathol.* **64**, 291–5.
- [157] A.M.H. Abdel-Moneim, A.R. Meki, A.M.A.S. Gabr, A.A. Mobasher and M.F. Lutfi (2014). The protective effect of green tea extract against lead toxicity in rats kidneys, *Asian J. Biomed. Pharm. Sci.* **4**, 30–34.
- [158] Y.I. Oboma, B. Sylvanus, P.N. Okara, F.A. Tamuno-omie and O.E. Ibiang (2018). Protective effect of combined aqueous extracts of *Allium sativum* and *Zingiber officinale* against lead acetate induced hepatotoxicity and testicular damage in Rattus norvegicus, *MOJ Anat. Physiol.* **5**, 306-313.
- [159] B.V.S. Lakshmi, M. Sudhakar, F.J. Sudha and M.V. Gopal (2015). Ameliorative effect of *Triticum aestivum* Linn against experimentally induced arsenic toxicity in male albino rats, *Der Pharm. Lett.* **7**, 202–211.



- [160] A. Sheikh, F. Yeasmin, S. Agarwal, M. Rahman, K. Islam, E. Hossain, S. Hossain, M.R. Karim, F. Nikkon, Z.A. Saud and K. Hossain (2014). Protective effects of *Moringa oleifera* Lam. leaves against arsenic-induced toxicity in mice, *Asian Pac. J. Trop. Biomed.* **4**, 353-358.
- [161] E.M. Khorasgani, I.S. Haghdooost, R. Sedaghat, P. Mortazavi and M. Roghani (2013). *Satureja hortensis* L. alcoholic extract ameliorates cadmium-induced pancreatic damage in rats, *Middle East J. Sci. Res.* **15**, 32–35.
- [162] E. Prithiviraj, S. Suresh, N.V. Lakshmi, M. Karthik, L. Ganesh and P. Seppan (2013). Protective effect of *Withania somnifera* (Linn.) on cadmium-induced oxidative injury in rat testis, *Ethnopharmacol.* **4**, 269-90.
- [163] S.L. Choubisa (2010). Natural amelioration of fluoride toxicity (fluorosis) in goats and sheep, *Curr. Sci.* **99**, 1331–1332.

**A C G**  
**publications**

© 2022 ACG Publications