

# Analysis of Phenolic Compounds by LC-HRMS and Determination of Antioxidant and Enzyme Inhibitory Properties of *Verbascum speciosum* Schrad

Hatice Kızıldaş <sup>1</sup>, Zeynebe Bingöl <sup>2</sup>, Ahmet C. Gören <sup>3</sup>,  
Saleh H. Alwasel <sup>4</sup> and İlhami Gülçin <sup>5,\*</sup>

<sup>1</sup>Vocational School of Health Services, Van Yuzuncu Yil University, 65080-Van, Turkiye

<sup>2</sup>Vocational School of Health Services, Tokat Gaziosmanpasa University, 60250-Tokat, Turkiye

<sup>3</sup>Department Chemistry, Faculty of Sciences, Gebze Technical University, 41400-Gebze-Kocaeli, Turkiye

<sup>4</sup>Department of Zoology, College of Science, King Saud University, Riyadh, Saudi Arabia

<sup>5</sup>Department of Chemistry, Faculty of Sciences, Atatürk University, 25240-Erzurum, Turkiye

(Received October 14, 2022; Revised November 23, 2022; Accepted December 01, 2022)

**Abstract:** Studies have shown an inverse correlation between age-related illnesses like coronary heart disease and cancer and fruit and vegetable intake. Given the probable health benefits of natural antioxidants from plants, research on them has increased. *Verbascum* L is a large genus of Scrophulariaceae family and 323 species distributed worldwide. Since *Verbascum* L. species are plants that are grown in many regions of Turkiye and are used in folk medicine, it is important to evaluate the biological activity of these species. In this study antioxidant properties of *Verbascum speciosum* Schrad. were investigated. The antioxidant capacities of water and ethanol-based extracts obtained from air parts were evaluated with Fe<sup>3+</sup> reducing, CUPRAC, FRAP, DPPH<sup>•</sup>, and ABTS<sup>•+</sup> scavenging antioxidant methods. Extracts were also examined to determine their AChE,  $\alpha$ -glycosidase and  $\alpha$ -amylase enzyme inhibitions. This investigation could be a basis for further phytochemical investigations of *Verbascum speciosum* Schrad.

**Keywords:** *Verbascum speciosum*; antioxidant activity; reducing power; radical scavenging; enzyme inhibition.  
© 2022 ACG Publications. All rights reserved.

## 1. Introduction

Oxidation is essential to fuel the biological processes of living organisms. However, it can cause uncontrolled production of oxygen-induced free radicals [1]. Free radicals are highly unstable, reactive, short-lived atoms or molecules with unpaired electrons. Therefore, they tend to attack and bind with other molecules to gain a more stable form [2,3]. During the functioning of aerobic metabolism, it is inevitable for cells to produce free radicals and especially reactive oxygen species (ROS) [4]. The

\* Corresponding author: E-Mail: [igulcin@atauni.edu.tr](mailto:igulcin@atauni.edu.tr) (I Gulcin); Phone:+904422314375, Fax:+904422314109

## Secondary metabolites and antioxidant and enzyme inhibitory activities of *Verbascum speciosum*

balance between ROS formation and reduction is controlled by the antioxidant system [5-7]. But, exposure to UV light, smoking, and other environmental pollutants and cell metabolism disorders also increase the body's free radical level and a defect in the cell's redox balance becomes [8,9]. Increased ROS attacks membranes, enzymes, lipids, carbohydrates, proteins, and DNA causing various disorders and diseases [10]. There are studies confirming the role of free radicals in the generation of diseases such as atherosclerosis, cancer, cardiovascular diseases, and brain dysfunction [11]. Antioxidants consist of some enzymes, vitamins E,  $\beta$ -carotene, or organic substances such as phenols and amines, which can stop radical chain reactions and counter the harmful effects of oxidation in tissues [12,13]. Antioxidants delay or completely inhibit the oxidation process of unstable molecules. In this way, antioxidants are molecules that can prevent the oxidation of biomolecules caused by free radicals and related cell damage [14]. Insufficient intake and production of antioxidants can cause oxidative stress in case of excessive formation of free radicals [15,16]. Natural antioxidants are created in living cell metabolism to maintain the oxidation-reduction balance [17]. Synthetic antioxidants are often used nowadays [1]. Recently, synthetic antioxidants are suspected to have carcinogenic effects and therefore their use has been limited, so there is significant attention on the discovery of natural antioxidants that can be used as substitutes [18-20]. Plants constitute an important source of active compounds and thus can show strong antioxidant effects. The fact that plants contain these active compounds that create physiological effects in the human body adds medicinal value to them [21,22]. Medicinal plants are recognized as a resource for the prevention of many diseases [23]. They attract the attention of researchers in the therapy of some diseases such as metabolic inflammations and cancer. Some phytochemicals are the subject of cancer therapy research. Many phenolic and flavonoid compounds are frequently used in drug development processes [24-26]. Studies have determined that phenolic compounds in plants reduce oxidative damage and prevent from free radicals [27]. Therefore, the commercial importance of plants containing these secondary compounds is increasing [28-30]. We believe *Verbascum speciosum* (*V. speciosum*) is one of those medicinal plants.

The genus *V. speciosum* belongs to the family Scrophulariaceae and includes 360 species distributed worldwide [31]. It is represented by 132 additional hybrids and 257 species in Türkiye among them, 202 species are endemic [32]. *Verbascum* L. genus is known as mullein. It has been reported that drugs prepared from different parts of *Verbascum* L. are used for skin, lung, and intestinal diseases, cough, headaches, and chills in Europe, Asia, Africa, and North America [33]. Traditionally, all parts of *Verbascum* L. species are used in Türkiye for the treatment of eczema and other skin disorders and for drying wounds [34]. *V. speciosum* is one of the well-known species of *Verbascum* L. It has been reported to be used traditionally for different diseases including hemorrhoids and coughs [33].

In the present study, ethanol extract of aerial parts of *V. speciosum* and lyophilized water extract of aerial part of *V. speciosum* (WEVS) were evaluated by several bioanalytical antioxidant methods including  $\text{Fe}^{3+}$ -TPTZ reducing capacity (FRAP),  $\text{Fe}^{3+}$  reducing,  $\text{Cu}^{2+}$  reducing ability (CUPRAC), DPPH and ABTS radical scavenging. Furthermore, total phenol and flavonoid contents were also investigated. Quantitative determination of phenolic substances was made by LC-HRMS. EEVS and WEVS were examined to determine their  $\alpha$ -amylase, acetylcholinesterase (AChE), and  $\alpha$ -glycosidase enzyme inhibitions.

## 2. Materials and Methods

### 2.1. Chemicals

All chemicals for antioxidant activity and enzyme inhibition such as  $\alpha$ -tocopherol, neocuproine (2,9-dimethyl-1,10-phenanthroline), BHT (butylated hydroxytoluene), DPPH, ascorbic acid, ABTS, 3-(2-pyridyl)-5,6-bis(4-phenyl-sulfonic acid)-1,2,4-triazine (Ferrozine), ascorbic acid, trichloroacetic acid (TCA),  $\alpha$ -tocopherol were obtained from Sigma (Sigma-Aldrich GmbH, Steinheim, Germany). The standard phenolics for LC-HRMS including ascorbic acid ( $\geq 99\%$ ), chlorogenic acid ( $\geq 95\%$ ), fumaric acid ( $\geq 99\%$ ), caffeic acid ( $\geq 98\%$ ), vanillic acid ( $\geq 97\%$ ), naringin ( $\geq 90\%$ ), rutin ( $\geq 94\%$ ), syringic acid ( $\geq 95\%$ ), rosmarinic acid ( $\geq 96\%$ ), p-coumaric acid ( $\geq 98\%$ ), quercetin ( $\geq 95\%$ ), salicylic acid ( $\geq 98\%$ ), naringenin ( $\geq 95\%$ ), luteolin (95%), emodin (90%) and chrysin ( $\geq 96\%$ ) were purchased from Sigma-

## Secondary metabolites and antioxidant and enzyme inhibitory activities of *Verbascum speciosum*

Aldrich, (-)-epigallocatechin (>97%), (-)-epigallocatechin gallate (>97%), orientin (>97%), (+)-trans-taxifolin (>97%), luteolin 7-glucoside (>97%), hyperoside (>97%), ellagic acid (>97%), quercitrin (>97%), apigenin (>97%), hispidulin (>97%), acacetin (>97%) and hederagenin (>97%) were purchased from TRC Canada, verbascoside (86.31%) was obtained from HWI Analytik GMBH. Luteolin-7-rutinoside (>97%) was purchased from Carbosynth limited. Hesperidin ( $\geq$ 98%) was purchased from J&K, dihydrokaempferol (>97%), nepetin-7-glucoside (>97%), isosakuranetin (>97%), and penduletin (>97%) was purchased from Phytolab, apigenin 7-glucoside (>97%) was purchased from EDQM CS. Myricetin (>95%) was purchased from Carl Roth GmbH+Co, nepetin (98%) was purchased from Supelco and caffeic acid phenethyl ester ( $\geq$ 97%) was purchased from European Pharmacopoeia.

### 2.2. Plant Materials

*Verbascum speciosum* Schrad. was collected from (A9) Kars: Sarıkamış, Soğuksu in June 2019 (Location: 40°18'19"K, 42°39'32"D, 2070 m. It was identified by botanist Dr. Süleyman Mesut PINAR, Van Yüzüncü Yıl University, Faculty of Science, Biology Department. The voucher specimen (voucher code: VANF 164145) is deposited at Van Yüzüncü Yıl University, Faculty of Science, Herbarium of the Biology Department (VANF), Van, Turkey.

### 2.3. Plant Extraction

To prepare WEVS, 40 g aerial parts *V. speciosum* were taken and shade-dried. It is powdered and boiled for 15 minutes with 500 mL distilled water, then, it is lyophilized in a lyophilizer (Labconco, Freezone 1L) at -50°C at 5 mm-Hg a pressure setting and stored at -20 °C [35,36]. The preparation of the EEVS was first started by pulverizing *V. speciosum* (25 g). The ground plant material was soaked with 0.5 L of ethanol, then the ethanol was evaporated via a rotary evaporator (Heidolph Hei-VAP HL, Germany) [37,38].

### 2.4. Radical Scavenging Assays

For evaluating the DPPH radical scavenging activity of EEVS and WEVS prepared and 0.5 mL of DPPH radical solution (0.1 mM) was transferred to each sample [39]. After 30 minutes of incubation, then, the absorbance was measured at 517 nm [40,41]. To determine the ABTS<sup>•+</sup> scavenging activity of EEVS and WEVS the method in a previous study was used [42]. First, 2.45 mM persulfate solution was transferred to 2 mM ABTS solution to generate ABTS radicals. The absorbance of the ABTS<sup>•+</sup> radical control solution was adjusted to  $0.750 \pm 0.025$  nm at 734 nm with a phosphate buffer of 0.1 M and pH 7.4. Then 1 mL of ABTS<sup>•+</sup> solution was added to different extract quantities and after 30 min incubation absorbances were recorded at 734 nm [43].

### 2.5. Reducing Ability

The Fe<sup>3+</sup> reducing capability of EEVS and WEVS was measured by a method that can measure the reducing ability of an extract by direct reduction of Fe[(CN)<sub>6</sub>]<sub>3</sub> to Fe[(CN)<sub>6</sub>]<sub>2</sub> [44]. As a result, the Perl-Prussian blue complex, which possesses absorbance at 700 nm, leads to the formation of Fe<sub>4</sub>[Fe(CN)<sub>6</sub>]<sub>3</sub> [45]. To determine the CUPRAC of EEVS and WEVS, a previous method was realized with some changes [46]. The FRAP assay is based on the reduction of the TPTZ-Fe<sup>3+</sup> complex [47].

### 2.6. Total Phenolics and Flavonoids

Quantification of phenolic compounds present in EEVS and WEVS was performed as described in a prior study [48]. The total amount of flavonoids found in EEVS and WEVS was carried out as described previously [49].

## Secondary metabolites and antioxidant and enzyme inhibitory activities of *Verbascum speciosum*

### 2.7. Enzyme Inhibition Assays

The enzyme inhibitory properties of EEVS were measured in accordance with previous studies [50,51]; Firstly, the AChE enzyme inhibitory property of EEVS was measured using electric eel as the source of AChE enzyme [52] and then, the inhibitory effect of EEVS on  $\alpha$ -glycosidase and  $\alpha$ -amylase was realized [53-55]. The IC<sub>50</sub> value is the concentration of extracts causing 50% inhibition. It was obtained from activity (%) against extract plots [56-58].

### 2.8. LC-HRMS Analysis

The phenolic contents of EEVS and WEVS were made according to the LC-HRMS analysis method [59]. LC-HRMS experiments were realized on a Thermo ORBITRAP Q-EXACTIVE mass spectrometry equipped with a Troyasil C18 column (3  $\mu$ m particle size, 150 x 3 mm i.d.) [60]. Acidified methanol and water gradient by HPLC assay was determined as the finest mobile phase [61,62]. The identification of the compounds was made by comparing the retention times of the standards (in the range of purity 95-99% see section chemicals) and HRMS data of Bezmialem Vakıf University, Drug Application and Research Centre Library (ILMER). Dihydrocapsaicin was used as an internal standard (IS) for LC-HRMS analysis for the reduction of repeatability problems caused by external effects including ionization repeatability, in mass spectrometry measurements. Dihydrocapsaicin solution (100 mg/L) was used as an IS. The mass values of the target phenolics are summarized in Table 4. Method Validation data and LC-HRMS chromatograms of the extracts were given as Table S1, Figure S1, and Figure S2 in Supporting Information.

### 2.9. Statistical Analyses

The results of the experiments were determined by averaging the triplicate analyses. For statistical analysis one-way analysis of variance (ANOVA) test was used. The differences between groups were made with Duncan's correction. The significance level was taken as  $p < 0.05$ ;  $p < 0.001$  was very significant and SPSS statistical software version 19.0 (SPSS Inc, Chicago, III, USA) package was used for analysis.

## 3. Results and Discussion

It is important to choose the most appropriate method when determining the antioxidant capacity of plants. In this study, two different free radical scavenging methods (DPPH<sup>•</sup> and ABTS<sup>•+</sup>) were used to determine the antioxidant ability of the extracts [63,64]. The basis of the DPPH method is based on the DPPH<sup>•</sup> scavenging percentage of antioxidants in the plant extract. ABTS assay is based on the percentage of antioxidants in the plant extract to scavenge ABTS<sup>•+</sup> radicals [65,66]. DPPH<sup>•</sup> scavenging activities of EEVS and WEVS and standard antioxidant compounds were investigated. The IC<sub>50</sub> values EEVS and WEVS and standard antioxidants demonstrated in the following order: Ascorbic acid (16.12 $\pm$ 0.003,  $r^2$ : 0.9566) <  $\alpha$ -Tocopherol (23.10 $\pm$ 0.032,  $r^2$ :0.9825) < EEVS (24.70 $\pm$ 0.009,  $r^2$ : 0.9878) < BHT (31.50 $\pm$ 0.011,  $r^2$ : 0.9754) < WEVS (173.25 $\pm$ 0.029,  $r^2$ : 0.9704) (Table 1 and Figure 1). The lower IC<sub>50</sub> values suggest an effective scavenged DPPH<sup>•</sup> scavenging activity [67]. Ascorbic acid was found to be the compound with the most powerful DPPH<sup>•</sup> scavenging activity. EEVS was found to have a free radical scavenging ability close to standard compounds. DPPH radical scavenging activity of the methanol extract of *V. speciosum* was determined in a study and the IC<sub>50</sub> value was found as 32.35  $\mu$ g/mL [68]. In another study, DPPH<sup>•</sup> scavenging of water extract of *V. speciosum* aerial parts was found to be 48.27  $\pm$  4.12  $\mu$ g/mL [33]. In the current study, it was determined that especially EEVS exhibited a strong DPPH<sup>•</sup> scavenging activity. When all these results were interpreted, it was observed that the IC<sub>50</sub> value levels were low in all studies. According to ABTS<sup>•+</sup> scavenging activity methods, a stable form of the radical is produced in the experiment and forms blue-green ABTS<sup>•+</sup> by reacting with an antioxidant, and decolorization specifies the rate of ABTS<sup>•+</sup> inhibition [45,69]. The IC<sub>50</sub> values of ABTS<sup>•+</sup> removing for EEVS, WEVS, and standards were determined in the following order:  $\alpha$ -

Secondary metabolites and antioxidant and enzyme inhibitory activities of *Verbascum speciosum*

Tocopherol ( $15.400 \pm 0.003$ ,  $r^2: 0.9866$ ) < EEVS ( $19.8 \pm 0.017$ ,  $r^2: 0.9956$ ) < Ascorbic acid ( $23.10 \pm 0.001$ ,  $r^2: 0.9998$ ) < BHT ( $26.654 \pm 0.008$ ,  $r^2: 0.9717$ ) < WEVS ( $34.65 \pm 0.010$ ,  $r^2: 0.9969$ ) (Table 1 and Figure 1). It was found that EEVS has an efficient ABTS<sup>+</sup> scavenging activity close to standard compounds. In a study conducted with *Verbascum euphraticum* and *Verbascum oocarpum*, ABTS<sup>+</sup> scavenging activity of methanol extract was found as  $90.52 \pm 0.05$  and  $90.62 \pm 0.06$   $\mu\text{g/mL}$ , respectively [34]. In the present study, the IC<sub>50</sub> amounts obtained for ABTS<sup>+</sup> scavenging activity were found to be more effective when compared with the different species mentioned above.


**Table 1.** DPPH radical and ABTS radical scavenging capability of the EEVS, WEVS, and standard compounds

Antioxidants	DPPH <sup>•</sup> scavenging		ABTS <sup>+</sup> scavenging	
	IC <sub>50</sub>	r <sup>2</sup>	IC <sub>50</sub>	r <sup>2</sup>
Ascorbic acid	$16.116 \pm 0.003$	0.9566	$23.10 \pm 0.001$	0.9998
$\alpha$ -Tocopherol	$23.1 \pm 0.032$	0.9825	$15.400 \pm 0.003$	0.9866
BHT	$31.500 \pm 0.011$	0.9754	$26.654 \pm 0.008$	0.9717
EEVS	$24.75 \pm 0.009$	0.9878	$19.8 \pm 0.017$	0.9956
WEVS	$173.25 \pm 0.029$	0.9704	$34.65 \pm 0.010$	0.9969

The antioxidant profile of extracts of *V. speciosum* was characterized using the ferric ions (Fe<sup>3+</sup>) reducing, CUPRAC, and FRAP assays [70], shown in Table 2 and Figure 2. Reduction capacity is an important factor in determining whether a molecule has antioxidant activity [71,72]. The first method used was to reduce Fe<sup>3+</sup> to Fe<sup>2+</sup> in Fe[(CN)<sub>6</sub>]<sup>3+</sup> solution, which is one of the common methods. The reaction system is based on the reduction of Fe<sup>3+</sup> in potassium ferricyanide to Fe<sup>2+</sup> with the addition of an antioxidant can be recorded at 700 nm [73,74]. The concentrations of positive controls EEVS and WEVS were increased steadily ( $\mu\text{g/mL}$ ). The reducing power of EEVS, WEVS, and standard antioxidants were as follows: Ascorbic acid ( $1.520 \pm 0.028$ ,  $r^2: 0.9970$ ) > BHT ( $1.269 \pm 0.005$ ,  $r^2: 0.9880$ ) > EEVS ( $1.177 \pm 0.018$ ,  $r^2: 0.9805$ ) >  $\alpha$ -Tocopherol ( $0.990 \pm 0.007$ ,  $r^2: 0.9942$ ) > WEVS ( $0.409 \pm 0.005$ ,  $r^2: 0.9964$ ) (Table 2 and Figure 2a). According to the results, it was determined that EEVS has a strong ability to reduce ferric ions (Fe<sup>3+</sup>) ( $p < 0.001$ ) (Table 2).

In the CUPRAC test, the absorbance measurement of the stable complex formed between copper ions and neocuproine is observed at 450 nm. High absorbance values indicate high reducing ability [75,76]. The CUPRAC of 30  $\mu\text{g/mL}$  of EEVS, WEVS, and standards are demonstrated in Table 2 and Figure 2b. The CUPRAC of EEVS, WEVS, and standards were as following order: BHT ( $1.561 \pm 0.089$ ,  $r^2: 0.9978$ ) > Ascorbic acid ( $1.069 \pm 0.007$ ,  $r^2: 0.9722$ ) >  $\alpha$ -Tocopherol ( $0.785 \pm 0.061$ ,  $r^2: 0.9986$ ) > EEVS ( $0.744 \pm 0.016$ ,  $r^2: 0.9858$ ) > WEVS ( $0.236 \pm 0.023$ ,  $r^2: 0.9536$ ) ( $p < 0.001$ , Table 2). According to the results, the reducing power of EEAC, Cu<sup>2+</sup> ions was high, although not as strong as standard compounds.

The FRAP method is based on measuring the power of a sample with antioxidant properties to reduce oxidant ferric iron to ferrous form [77,78]. The FRAP of EEVS, WEVS, and standard antioxidants decreased as follows: Ascorbic acid ( $1.624 \pm 0.015$ ,  $r^2: 0.9930$ ) > BHT ( $0.909 \pm 0.006$ ,  $r^2: 0.9874$ ) >  $\alpha$ -Tocopherol ( $0.755 \pm 0.075$ ,  $r^2: 0.9867$ ) > EEVS ( $0.584 \pm 0.002$ ,  $r^2: 0.9288$ ) > WEVS ( $0.361 \pm 0.005$ ,  $r^2: 0.9567$ ) (Table 2 and Figure 2c). According to the method, higher absorbances represent the higher reduction capability of the Fe<sup>3+</sup>-TPTZ. EEVS and WEVS demonstrated an effective FRAP reduction ability although not as strong as standard compounds ( $p < 0.001$ , Table 2). In a study, the FRAP of water extract from *V. speciosum* was found to be  $754.41 \pm 10.39$   $\mu\text{g/mL}$  [33]. In conclusion, it was found that extracts of *V. speciosum* showed a value close to the standard compounds in all antioxidant activity tests.


Secondary metabolites and antioxidant and enzyme inhibitory activities of *Verbascum speciosum*

**Figure 1.** Radical removing activity of EEVS and WEVS. **A.** DPPH radical removing activity. **B.** ABTS radical removing activity (10-30  $\mu\text{g/mL}$ )

**Table 2.** The reducing ability of the EEVS, WEVS, and standards by  $\text{Fe}^{3+}$  (120  $\mu\text{g/mL}$ ),  $\text{Cu}^{2+}$  (30  $\mu\text{g/mL}$ ) and  $\text{Fe}^{3+}$ -TPTZ (150  $\mu\text{g/mL}$ ) reducing ability

Antioxidants	$\text{Fe}^{3+}$ - $\text{Fe}^{2+}$ reducing		$\text{Cu}^{2+}$ - $\text{Cu}^{+}$ reducing		$\text{Fe}^{3+}$ -TPTZ reducing	
	$\lambda_{700}$	$r^2$	$\lambda_{450}$	$r^2$	$\lambda_{593}$	$r^2$
$\alpha$ -Tocopherol <sup>a</sup>	1.520 $\pm$ 0.028 <sup>b,c,d,e</sup>	0.9970	1.069 $\pm$ 0.007 <sup>b,d,e</sup>	0.9722	1.624 $\pm$ 0.015 <sup>b,c,d,e</sup>	0.9930
Ascorbic acid <sup>b</sup>	0.990 $\pm$ 0.007 <sup>e</sup>	0.9942	0.785 $\pm$ 0.061 <sup>d,e</sup>	0.9986	0.755 $\pm$ 0.075 <sup>d,e</sup>	0.9867
BHT <sup>c</sup>	1.269 $\pm$ 0.005 <sup>b,d,e</sup>	0.9880	1.561 $\pm$ 0.089 <sup>a,b,d,e</sup>	0.9978	0.909 $\pm$ 0.006 <sup>b,d,e</sup>	0.9874
EEVS <sup>d</sup>	1.177 $\pm$ 0.018 <sup>b,e</sup>	0.9805	0.744 $\pm$ 0.016 <sup>e</sup>	0.9858	0.584 $\pm$ 0.002 <sup>e</sup>	0.9288
WEVS <sup>e</sup>	0.409 $\pm$ 0.005	0.9964	0.236 $\pm$ 0.023	0.9536	0.361 $\pm$ 0.005	0.9567

\*Different letters in the same column were significantly different ( $p < 0.05$ ). *a*:  $\alpha$ -Tocopherol, *b*: Ascorbic acid, *c*: BHT, *d*: EEVS, *e*: WEVS.

Secondary metabolites and antioxidant and enzyme inhibitory activities of *Verbascum speciosum*

**Figure 2.** The reducing abilities of EEVS, WEVS and standard compounds.

## Secondary metabolites and antioxidant and enzyme inhibitory activities of *Verbascum speciosum*

The antioxidant effects of plants are due to the secondary metabolites such as phenolic and flavonoid compounds they contain [79]. Phenolic compounds are among the main plant secondary metabolites [80]. It was determined that the foods consumed rich in phenolic have a protective effect against cardiovascular diseases and cancer [81]. Phenolics and flavonoids have anticancer, anti-allergic, antibacterial, and anti-inflammatory properties and are free radical scavengers. Flavonoids are an important class of secondary metabolites found in plants. Numerous phenolic hydroxyl groups attached to the ring structures of flavonoids give them antioxidant activity [82]. Due to their strong free radical scavenging properties, flavonoids show antioxidant activities such as metal chelation and reduction [83].

The total phenolic contents of EEVS and WEVS were found to be  $24.545 \pm 4.45$   $\mu\text{g}$  GAE and  $16.364 \pm 0.01$   $\mu\text{g}$  GAE, respectively. The total flavonoid contents of EEVS and WEVS were found to be  $5.328 \pm 0.75$   $\mu\text{g}$  QE and  $16.750 \pm 0.69$   $\mu\text{g}$  QE, respectively. It was found a positive correlation between total phenolics and flavonoids in EEVS and WEVS and antioxidant abilities. In a study, the methanol extract of *Verbascum speciosum* was investigated and the total phenol and flavonoid contents were found as  $82.00 \pm 6.43$  mg/g GAE and  $30.79 \pm 0.50$  (mg/g) rutin equivalent, respectively [84]. In a study conducted in 2020, total phenolic and flavonoid contents in the *V. speciosum* methanol extract were found to be  $45.01 \pm 1.44$  mg GAE and  $14.26 \pm 1.81$  mg QE, respectively [33]. *Verbascum euphraticum* and *Verbascum oocarpum* species were examined in a study and their total phenolic contents were respectively;  $50.29 \pm 0.29$  (mg GAE/g) and  $61.23 \pm 1.74$  (mg GAE/g) and total flavonoid contents were found as  $28.47 \pm 0.25$  (mg RE/g) and  $49.97 \pm 0.32$  (mg RE/g), respectively [34]. The results obtained in the previous studies mentioned above were found to be sometimes higher and sometimes lower than our results. The reason for this is thought to be due to differences in the ecological and soil structure of the region where the plant is grown, analysis methods, solvents, and extraction conditions.

**Table 3.** The total phenolics and flavonoids in EEVS and WEVS

Extracts	Total phenolics ( $\mu\text{g}/\text{mL}$ extract)	Total flavonoid ( $\mu\text{g}/\text{mL}$ extract)
EEVS	$24.545 \pm 4.45$	$5.328 \pm 0.75$
WEVS	$16.364 \pm 0.01$	$16.750 \pm 0.69$

Various iridoids, phenyletanoids, phenylpropanoids, saponins, and flavonoids were determined as secondary metabolites in *Verbascum* species [85]. In this study, the phenolic compound contents of EEVS and WEVS were determined by the LC-HRMS technique and thirty-eight polyphenols were quantified and identified (Table 4). Verbascoside (13740.0 mg/kg), p-coumaric acid (10983.89 mg/kg), and fumaric acid (2221.61 mg/kg) are the main polyphenols identified in 1 mg of EEVS while verbascoside (4390.71 mg/kg), rutin (561.82 mg/kg) and apigenin (561.54 mg/kg) are the main polyphenols in 1 mg of WEVS. It was determined that the ethanolic extracts showed the highest compound content from both extracts. Among the 38 compounds, 17 molecules were found in both EEVS and WEVS, and verbascoside was most abundant in the EEVS and WEVS. Verbascoside is a phenylethanoid glycoside and has been isolated from many different plant species, *Verbascum sinuatum* being one of them. Verbascoside has been the subject of many studies due to its pharmacological effects such as antioxidant, anti-inflammatory, neuroprotective, and anticancer effects. Polyphenols, which are secondary plant metabolites, contain thousands of structural variations from simple phenolics to tannins [86-88]. The main dietary polyphenols are phenolic acids, flavonoids, tannins, catechins, lignans, stilbenes, and anthocyanidin phytochemicals. It has been reported that more than 8,000 of the many dietary polyphenols are of plant origin. Studies have shown that polyphenolics in the diet are helpful in preventing diseases [89,90]. Polyphenols have antioxidant properties, so they can significantly lower the risk of developing many chronic diseases, particularly cancer, cardiovascular disease, inflammatory diseases, diabetes, kidney failure, and neurodegenerative disorders [81,82]. Plants have been used for a long time because of the natural products they contain such as antioxidants [93]. Plants have also been used for medicinal applications from time immemorial. Since the level of side effects, they show is low, their use in the pharmaceutical and food industry is increasing today [94]. One of the reasons for this



Secondary metabolites and antioxidant and enzyme inhibitory activities of *Verbascum speciosum*

increase is that thanks to scientific developments, the presence of phytochemical compounds exhibiting pharmacological properties of medicinal plants can be detected so that the therapeutic properties of these plants can be verified [95].

**Table 4.** The quantity of phenolics (mg/kg extract) in EEAC and WEAC by LC-HRMS

Compounds	EEVS	WEVS
Ascorbic acid	68.30	302.93
(-)-Epigallocatechin	<LOD	<LOD
(-)-Epigallocatechin gallate	<LOD	-
Chlorogenic acid	31.25	86.95
Fumaric acid	2221.61	-
(-)-Epicatechin	<LOD	-
(-)-Epicatechin gallate	<LOD	-
Verbascoside	13740.00	4390.71
Orientin	0.42	<LOD
Caffeic acid	33.65	302.76
(+)- <i>trans</i> taxifolin	0.02	-
Luteolin-7-rutinoside	261.72	99.70
Naringin	<LOD	<LOD
Luteolin 7-glucoside	1305.78	191.67
p-Coumaric acid	10983.89	-
Hesperidin	<LOD	-
Rutin	805.58	561.82
Syringic acid	<LOD	-
Rosmarinic acid	10.20	13.08
Hyperoside	590.93	83.68
Dihydrokaempferol	0.05	-
Apigenin 7-glucoside	26.51	6.15
Ellagic acid	<LOD	-
Quercitrin	38.93	<LOD
Myricetin	0.02	-
Quercetin	2.63	1.86
Salicylic acid	6.72	5.53
Naringenin	3.09	<LOD
Luteolin	409.17	371.33
Nepetin	0.65	-
Apigenin	77.97	561.54
Hispidulin	188.79	56.17
Isosakuranetin	<LOD	<LOD
Penduletin	3.98	-
Caffeic asit phenethyl ester	0.14	-
Chrysin	1.32	4.74
Acacetin	3.42	2.82
Hederagenin	<LOD	-

Alzheimer's disease (AD) is a common and irreversible age-related neurodegenerative disease and is the most common cause of dementia [98]. Memory loss, daily life disorders, neuropsychiatric symptoms, and other behavioral disorders are among the clinical symptoms of AD [99]. Decreased acetylcholine (ACh) levels are among the etiological features of AD. The enzyme acetylcholinesterase (AChE) hydrolyzes ACh to acetate and choline [100]. By using AChE inhibitors, inhibition of ACh

Secondary metabolites and antioxidant and enzyme inhibitory activities of *Verbascum speciosum*

hydrolysis can be achieved, increasing neurotransmitter levels in synapses and improving cholinergic and cognitive neurotransmission capacity [101]. In the current study, the results of AD-related cholinesterase inhibition were evaluated and a high anti-AChE activity measured with the ethanol acetate extract of *V. speciosum* ( $IC_{50}$ : 1.516  $\mu\text{g/mL}$ ,  $r^2$ : 0.9719) (Table 5). Tacrine was used with an  $IC_{50}$  value of 0.124  $\mu\text{M}$  against AChE (Table 5) [19]. In a previous study, *Verbascum euphraticum* and *Verbascum oocarpum* species were examined for their AChE inhibition, the results were  $1.93 \pm 0.21$  (mg Galantamine/g) and  $0.81 \pm 0.16$  (mg Galantamine/g), respectively [34].

**Table 5.** The half-maximal enzyme inhibition values ( $IC_{50}$ ,  $\mu\text{g/mL}$ ) of EEVS against  $\alpha$ -glycosidase, acetylcholinesterase, and  $\alpha$ -amylase

Enzymes	EEVS		Standards
	$IC_{50}$ ( $\mu\text{g/mL}$ )	$r^2$	$IC_{50}$ ( $\mu\text{M}$ )
$\alpha$ -Glycosidase <sup>a</sup>	1.120	0.9441	22.80
$\alpha$ -Amylase <sup>a</sup>	0.859	0.9811	10.01
Acetylcholinesterase <sup>b</sup>	1.516	0.9719	0.124

<sup>a</sup>Acarbose was used as a standard for  $\alpha$ -glycosidase and  $\alpha$ -amylase enzymes and taken from the reference of [96] and [97], respectively.

<sup>b</sup>Tacrine was used as a standard for acetylcholinesterase enzyme and taken from our previous study [41].

Diabetes mellitus (DM) is a main cause of mortality and morbidity worldwide. DM is a chronic disease that includes markers of hyperglycemia and insulin deficiency. WHO reported that deaths from diabetes mellitus will double between 2005-2030 [102].  $\alpha$ -Glycosidase is a crucial enzyme, which catalyzes the final step in the digestive mechanism of carbohydrates.  $\alpha$ -Glycosidase inhibitors inhibit enzyme activity and prevent the rapid breakdown of sugar units, thus keeping the blood sugar level under control. Therefore,  $\alpha$ -glucosidase inhibitors are used clinically to treat diabetes, but most of these inhibitors are synthetic drugs with some harmful reactions [103]. Investigation of the inhibitory effects of dietary polyphenols on  $\alpha$ -glucosidases is of great interest [104].  $\alpha$ -Amylase is an enzyme that helps break down large, insoluble molecules such as starch into soluble ones.  $\alpha$ -Amylase inhibitors delay the breakdown of carbohydrates and prevent the sudden increase of postprandial glucose in the blood [105]. The  $\alpha$ -amylase enzyme is under investigation specifically to treat diabetes and hyperlipidemia. Inhibitory effects of polyphenols for  $\alpha$ -amylases have been widely reported [106].

The determination of the inhibition of antidiabetic enzymes of  $\alpha$ -amylase and  $\alpha$ -glycosidase enzymes was evaluated for the determination of antidiabetic profiles of *V. speciosum*. The  $IC_{50}$  values of  $\alpha$ -glycosidase were 1.120  $\mu\text{g/mL}$  ( $r^2$ : 0.9441) for EEAC and 22.80  $\mu\text{M}$  for acarbose (Table 5). Also, the  $IC_{50}$ s of  $\alpha$ -amylase were 0.859  $\mu\text{g/mL}$  ( $r^2$ : 0.9811) for EEVS and 10.01  $\mu\text{M}$  for acarbose, respectively (Table 5) [97]. According to the results, it was shown that EEVS has a very high affinity towards enzymes including  $\alpha$ -glycosidase and  $\alpha$ -amylase. It was more effective than the standard inhibitor acarbose. In a previous study, *Verbascum euphraticum* was examined for its  $\alpha$ -glycosidase inhibition, the result was  $0.65 \pm 0.01$  (mmol Acarbose equivalent/g), and also at the same research *Verbascum euphraticum* and *Verbascum oocarpum* examined for their  $\alpha$ -amylase inhibition and the results were as  $0.31 \pm 0.02$  (mmol Acarbose E/g) and  $0.31 \pm 0.01$  (mmol Acarbose equivalent/ g), respectively [34].

#### 4. Conclusions

In this study, data on the phytochemical bioactivity and properties of *V. speciosum* extracts were evaluated for their antioxidant ability, phenolics, and flavonoids, as well as their anti-Alzheimer and anti-diabetic properties. The ethanol extract of the aerial part of *V. speciosum* was found to have a value close to the standard antioxidants, also, a high content of total flavonoids and phenolics. In addition, it was also evaluated that there was a statistically significant difference between the averages of the groups. It was determined by LC-HRMS for the main phenolic compound in both extracts was verbascoside.

## Secondary metabolites and antioxidant and enzyme inhibitory activities of *Verbascum speciosum*

Also, *V. speciosum* showed high inhibition effects against AChE,  $\alpha$ -glycosidase, and  $\alpha$ -amylase enzymes. EEAC has higher antioxidant activity and inhibition effect compared to WEAC as it takes both hydrophobic and hydrophilic components. Values obtained from this study have shown that *V. speciosum* as a source of potential bioactive components is a candidate for use in drug design processes for the treatment of DM and AD, due to its potent pharmacological activity.

### Acknowledgment:

S. Alwasel would like to extend his sincere appreciation to the Researchers Supporting Project (RSP-2023/59), King Saud University, Saudi Arabia, for support. Also, for his support in the collection and identification of plant samples, we would like to thank Assoc. Dr. Süleyman Mesut Pinar.

### ORCID

Hatice Kızıldağ : 0000-0002-0833-3358

Zeynebe Bingöl: 0000-0003-3373-779X

Ahmet C. Gören: 0000-0002-5470-130X

Saleh H. Alwasel: 0000-0002-0626-2306

İlhami Gülçin: 0000-0001-5993-1668

### References

- [1] I. Gulcin (2020). Antioxidants and antioxidant methods-An updated overview, *Arch. Toxicol.* **94**, 651-715.
- [2] M. Işık, S. Beydemir, A. Yılmaz, M.E. Naldan, H.E. Aslan and İ. Gülçin (2017). Oxidative stress and mRNA expression of acetylcholinesterase in the leukocytes of ischemic patients, *Biomed. Pharmacother.* **87**, 561-567.
- [3] F.M. Kandemir, S. Kucukler, E. Eldutar, C. Caglayan and İ. Gülçin (2017). Chrysin protects rat kidney from paracetamol-induced oxidative stress, inflammation, apoptosis, and autophagy: A multi-biomarker approach, *Sci. Pharmaceut.* **85(1)**, 4.
- [4] E. Köksal, E. Bursal, E. Dikici, F. Tozoğlu and İ. Gülçin (2011). Antioxidant activity of *Melissa officinalis* leaves, *J. Med. Plants Res.* **5(2)**, 217-222.
- [5] A. Baghiani, N. Boussoualim, H. Trabsa, S. Aouachria and L. Arrar (2013). In vivo free radical scavenging, Antihemolytic activity and antibacterial effects of *Anchusa azurea* extracts, *Int. J. Med. Sci.* **46**, 1113-1118.
- [6] I. Gulcin, (2012). Antioxidant activity of food constituents: An overview, *Arch. Toxicol.* **86(3)**, 345-391.
- [7] V. Morocho, D. Sarango, C. Cruz-Erazo, N. Cumbicus, L. Cartuche and A.I. Suárez (2020). Chemical constituents of *Croton thurifer* Kunth as  $\alpha$ -glucosidase inhibitors, *Rec. Nat. Prod.* **14(1)**, 31-41
- [8] P. Shu, L. Zhang, W. Liu, Y. Fei, M. Sun, Y. Lou, A. Liu, M. Yu, J. Li, X. Wei and N. Sun (2021). Chemical constituents from *Typhonium giganteum* rhizome and their antioxidant, tyrosinase inhibitory activities, *Rec. Nat. Prod.* **15**, 53-58.
- [9] İ. Gülçin (2008). Measurement of antioxidant ability of melatonin and serotonin by the DMPD and CUPRAC methods as trolox equivalent, *J. Enzyme Inhib. Med. Chem.* **23(6)**, 871-876.
- [10] M. Elmastas, İ. Türkekel, L. Öztürk, I. Gulcin, Ö. Işıldak and H.Y. Aboul-Enein (2006). The antioxidant activity of two wild edible mushrooms (*Morchella vulgaris* and *Morchella esculanta*), *Comb. Chem. High Throughput Screen.* **9(6)**, 443-448
- [11] V.N. Hristo, I.I. Iliana (2014). 8th Conference on medicinal and aromatic plants of southeast European countries, *Z. Arznei- Gewurzpfla.* **19(4)**, 293-299.
- [12] İ. Gülçin, V. Mshvildadze, A. Gepdiremen and R. Elias (2006). Antioxidant activity of a triterpenoid glycoside isolated from the berries of *Hedera colchica*: 3-O-( $\beta$ -D-glucopyranosyl)-hederagenin. *Phytother. Res.* **20(2)**, 130-134.
- [13] E. Köksal and İ. Gülçin (2008). Purification and characterization of peroxidase from cauliflower (*Brassica oleracea* L.) buds, *Protein Peptide Lett.* **15(4)**, 320-326.
- [14] N. Carsono, S.G. Tumilaar, D. Kurnia, D. Latipudin and M.H. Satari (2022). A review of bioactive compounds and antioxidant activity properties of piper species, *Molecules*, **27**, 6774.
- [15] İ. Gülçin, M. Oktay, E. Köksal, H. Şerbetçi, Ş. Beydemir and Ö.I. Küfrevioğlu (2008). Antioxidant and radical scavenging activities of uric acid, *Asian J. Chem.* **20(3)**, 2079-2090.

Secondary metabolites and antioxidant and enzyme inhibitory activities of *Verbascum speciosum*

- [16] Y. Çetinkaya, H. Göçer, A. Menzek and İ. Gülçin, (2012). Synthesis and antioxidant properties of (3,4-dihydroxyphenyl) (2,3,4-trihydroxyphenyl) methanone and its derivatives, *Arch. Pharm.* **345(4)**, 323-334. 172.
- [17] H. Kiziltas, A.C. Gören, S. Alwasel and İ. Gulçin (2022). Sahlep (*Dactylorhiza osmanica*): Phytochemical analyses by LC-HRMS, molecular docking, antioxidant activity and enzyme inhibition profiles, *Molecules*, **27(20)**, 6907.
- [18] A. Baran, E. Karakilic, Ö. Faiz and F. Özen (2020). Synthesis of chalcone-containing zinc and cobalt metallophthalocyanines; investigation of their photochemical, DPPH radical scavenging and metal chelating characters, *Org. Commun.* **13(2)**, 65-78.
- [19] E. Bursal, A. Aras, Ö. Kılıç, P. Taslimi, A.C. Gören and I. Gulcin (2019). Phytochemical content, antioxidant activity and enzyme inhibition effect of *Salvia eriophora* Boiss. & Kotschy against acetylcholinesterase,  $\alpha$ -amylase, butyrylcholinesterase and  $\alpha$ -glycosidase enzymes, *J. Food Biochem.* **43(3)**, e12776.
- [20] M. Topal, H. Gocer, F. Topal, P. Kalin, L. Polat Köse, İ. Gülçin, K. Cetin Çakmak, M. Küçük, L. Durmaz, A.C. Gören and S.H. Alwasel (2016). Antioxidant, antiradical and anticholinergic properties of cynarin purified from the illyrian thistle (*Onopordum illyricum* L.), *J. Enzyme Inhib. Med. Chem.* **31(2)**, 266-275.
- [21] I. Gulcin, Ş. Beydemir, İ.G. Şat and Ö.İ. Küfrevioğlu (2005). Evaluation of antioxidant activity of cornelian cherry (*Cornus mas* L.), *Acta Aliment. Hung.* **34(2)**, 193-202.
- [22] I. Gulcin and A. Daştan (2007). Synthesis of dimeric phenol derivatives and determination of in vitro antioxidant and radical scavenging activities, *J. Enzyme Inhib. Med. Chem.* **22(6)**, 685-695.
- [23] İ. Gülçin, İ.G. Şat, Ş. Beydemir and Ö.İ. Küfrevioğlu (2004). Evaluation of the in vitro antioxidant properties of extracts of broccoli (*Brassica oleracea* L.), *Ital. J. Food Sci.* **16(1)**, 17-30.
- [24] M. Şentürk, İ. Gülçin, Ş. Beydemir, Ö.İ. Küfrevioğlu and C.T. Supuran, (2011). In vitro inhibition of human carbonic anhydrase I and II isozymes with natural phenolic compounds, *Chem. Biol. Drugs Des.* **77(6)**, 494-499.
- [25] S.B. Öztürk Sarıkaya, F. Topal, M. Şentürk, İ. Gülçin and C.T. Supuran, (2011). In vitro inhibition of  $\alpha$ -carbonic anhydrase isozymes by some phenolic compounds, *Bioorg. Med. Chem. Lett.* **21(14)**, 4259-4262.
- [26] İ. Gülçin and S. Beydemir (2013). Phenolic compounds as antioxidants: Carbonic anhydrase isoenzymes inhibitors, *Mini Rev. Med. Chem.* **13(3)**, 408-430.
- [27] I. Gulcin (2005). The antioxidant and radical scavenging activities of black pepper (*Piper nigrum*) seeds, *Int. J. Food Sci. Nutr.* **56(7)**, 491-499.
- [28] Z. Huyut, S. Beydemir and İ. Gülçin (2016). In vitro and in vivo inhibitory effects of some phenolic compounds on the activities of carbonic anhydrase, *J. Enzyme Inhib. Med. Chem.* **31(6)**, 1234-1240.
- [29] H. Tohma, I. Gulcin, E. Bursal, A.C. Gören, S.H. Alwasel and E. Köksal (2017). Antioxidant activity and phenolic compounds of ginger (*Zingiber officinale* Rosc.) determined by HPLC-MS/MS, *J. Food Measure. Charac.* **11(2)**, 556-566.
- [30] H. Tohma, E. Köksal, Ö. Kılıç, Y. Alan, M.A. Yılmaz, İ. Gülçin, E. Bursal and S.H. Alwasel, (2016). RP-HPLC/MS/MS analysis of the phenolic compounds, antioxidant and antimicrobial activities of *Salvia* L. species, *Antioxidants* **5**, 38, 84.
- [31] V.H. Heywood, Flowering plants of the world. Oxford Univ Press. 1993.
- [32] M. Firat (2022). *Verbascum zerdust* (Scrophulariaceae), a new species from Bitlis province (Turkey) belonging to section Bothrosperma, *Nord. J. Bot.* doi:10.1111/njb.03649.
- [33] M. Selseleh, S.N. Ebrahimi, A. Aliahmadi, A. Sonboli and M.H. Mirjalili (2020). Metabolic profiling, antioxidant, and antibacterial activity of some Iranian *Verbascum* L. species, *Ind. Crops. Prod.* **153**, 112609.
- [34] G. Zengin, M. Fawzi, K. Ibrahime, N. Sadeer, F. Maggi, G. Caprioli, S. Angeloni, A. Mollica, A. Stefanucci, G. Ak, U. Cakılcıoglu, R. Polat and H. Akan (2021). Evaluation of chemical constituents and biological properties of two endemic *Verbascum* species, *Proces. Biochem.* **108**, 110-120.
- [35] P. Kalin, I. Gulcin and A.C. Gören (2015). Antioxidant activity and polyphenol content of cranberries (*Vaccinium macrocarpon*), *Rec. Nat. Prod.* **9** 496-502.
- [36] İ. Gülçin, R. Elias, A. Gepdiremen, L. Boyer and E. Köksal (2007). A comparative study on the antioxidant activity of fringe tree (*Chionanthus virginicus* L.) extracts, *Afr. J. Biotechnol.* **6(4)**, 410-418.
- [37] F.M. Birdane, M. Cemek, Y.O. Birdane, İ. Gülçin and M.E. Büyükokuroğlu (2007). Beneficial effects of *Foeniculum vulgare* on ethanol-induced acute gastric mucosal injury in rats, *World J. Gastroenterol.* **13(4)**, 607-611.
- [38] K. Çetin Çakmak and I. Gulcin (2019). Anticholinergic and antioxidant activities of usnic acid-an activity-structure insight, *Toxicol. Rep.* **6**, 1273-1280.
- [39] S. Çakmakçı, E.F. Topdaş, P. Kalm, H. Han, P. Şekerci, L. Polat Kose and İ. Gülçin (2015). Antioxidant capacity and functionality of oleaster (*Elaeagnus angustifolia* L.) flour and crust in a new kind of fruity ice cream, *Int. J. Food Sci. Technol.* **50(2)**, 472-481.

Secondary metabolites and antioxidant and enzyme inhibitory activities of *Verbascum speciosum*

- [40] M. Akdeniz, I. Yener, A. Ertas, M. Firat, B. Resitoglu, N. Hasimi, S.I. Kandemir, M.A. Yilmaz, A.B. Demirkoz, U. Kolak, and S. Oksuz (2021). Biological and chemical comparison of natural and cultivated samples of *Satureja macrantha* CA Mey, *Rec. Nat. Prod.* **15(6)**, 568-584
- [41] P. Taslimi, E. Koksals, A.C. Gören, E. Bursal, A. Aras, O. Kılıç, S. Alwasel and I. Gulcin (2020). Anti-Alzheimer, antidiabetic and antioxidant potential of *Satureja cuneifolia* and analysis of its phenolic contents by LC-MS/MS, *Arab. J. Chem.* **13(3)**, 4528-4537.
- [42] İ. Gülçin, S. Beydemir, F. Topal, N. Gagua, A. Bakuridze, R. Bayram and A. Gepdiremen (2012). Apoptotic, antioxidant and antiradical effects of majdine and isomajdine from *Vinca herbacea* Waldst. and kit, *J. Enzyme Inhib. Med. Chem.* **27(4)**, 587-594.
- [43] I. Gulcin, A.C. Gören, P. Taslimi, S.H. Alwasel, O. Kilic and E. Bursal (2020). Anticholinergic, antidiabetic and antioxidant activities of Anatolian pennyroyal (*Mentha pulegium*)-analysis of its polyphenol contents by LC-MS/MS, *Biocat. Agric. Biotechnol.* **23**, 101441.
- [44] N. Bulut, U.M. Koçyiğit, I.H. Gecibesler, T. Dastan, H. Karci, P. Taslimi, S. Durna Dastan, İ. Gulçin and A. Cetin (2018). Synthesis of some novel pyridine compounds containing bis-1,2,4-triazole moiety and investigation of their antioxidant properties, carbonic anhydrase and acetylcholinesterase enzymes inhibition profiles, *J. Biochem. Mol. Toxicol.* **32(1)**, e22006.
- [45] M.H. Sehitoglu, H. Han, P. Kalin, İ. Gülçin, A. Ozkan and H.Y. Aboul-Enein (2015). Pistachio (*Pistacia vera* L.) gum: A potent inhibitor of reactive oxygen species, *J. Enzyme Inhib. Med. Chem.* **30(2)**, 264-269.
- [46] N. Eruygur, M. Ataş, M. Tekin, P. Taslimi, U.M. Koçyiğit and I. Gulcin (2019). In vitro antioxidant, antimicrobial, anticholinesterase and antidiabetic activities of Turkish endemic *Achillea cucullata* (Asteraceae) from ethanol extract, *S. Afr. J. Bot.* **120**, 141-145.
- [47] F. Turkan, M.N. Atalar, A. Aras, I. Gulcin and E. Bursal (2020). ICP-MS and HPLC analyses, enzyme inhibition and antioxidant potential of *Achillea schischkinii* Sosn, *Bioorg. Chem.* **94**, 103333.
- [48] H. Tohma, A. Altay, E. Koksals, A.C. Gören and İ. Gülçin (2019). Measurement of anticancer, antidiabetic and anticholinergic properties of sumac (*Rhus coriaria*) - Analysis of its phenolic compounds by LC-MS/MS, *J. Food Measure. Charac.* **13(2)**, 1607-1619.
- [49] E. Koksals, S.H. Tohma, Ö. Kılıç, Y. Alan, A. Aras, İ. Gulcin and E. Bursal (2017). Assessment of antimicrobial and antioxidant activities of *Nepeta trachonitica*-Analysis of its phenolic compounds using HPLC-MS/MS, *Sci. Pharm.* **15**, 85(2).
- [50] F. Özbey, P. Taslimi, İ. Gulcin, A. Maraş, S. Goksu and C.T. Supuran (2016). Synthesis, acetylcholinesterase, butyrylcholinesterase, carbonic anhydrase inhibitory and metal chelating properties of some novel diaryl ether, *J. Enzyme Inhib. Med. Chem.* **31(S2)**, 79-85.
- [51] E. Garibov, P. Taslimi, A. Sujayev, Z. Bingöl, S. Çetinkaya, İ. Gulcin, S. Beydemir, V. Farzaliyev, S.H. Alwasel and C.T. Supuran (2016). Synthesis of 4,5-disubstituted-2-thioxo-1,2,3,4-tetrahydropyrimidines and investigation of their acetylcholinesterase, butyrylcholinesterase, carbonic anhydrase I/II inhibitory and antioxidant activities, *J. Enzyme Inhib. Med. Chem.* **31(S3)**, 1-9.
- [52] H.İ. Gül, M. Tuğrak, H. Sakagami, P. Taslimi, İ. Gülçin and C.T. Supuran (2016). Synthesis and bioactivity studies on new 4-(3-(4-substitutedphenyl)-3a,4-dihydro-3h-indeno[1,2-c]pyrazol-2-yl) benzenesulfonamides, *J. Enzyme Inhib. Med. Chem.* **31(6)**, 1619-1624.
- [53] F. Erdemir, D. Barut Celepci, A. Aktaş, Y. Gök, R. Kaya, P. Taslimi, Y. Demir and İ. Gülçin (2019). Novel 2-aminopyridine liganded Pd (II) N-heterocyclic carbene complexes: synthesis, characterization, crystal structure and bioactivity properties, *Bioorg. Chem.* **91**, 103134.
- [54] A. Biçer, P. Taslimi, G. Yakali, İ. Gülçin, M.S. Gültekin and G. Turgut Cin (2019). Synthesis, characterization, crystal structure of novel bis-thiomethylcyclohexanone derivatives and their inhibitory properties against some metabolic enzymes, *Bioorg. Chem.* **82**, 393-404.
- [55] C. Çağlayan, Y. Demir, S. Küçükler, P. Taslimi, F.M. Kandemir and İ. Gulçin (2019). The effects of hesperidin on sodium arsenite-induced different organ toxicity in rats on metabolic enzymes as antidiabetic and anticholinergics potentials: A biochemical approach, *J. Food Biochem.* **43(2)**, e12720.
- [56] I. Gulcin, A. Scozzafava, C.T. Supuran, H. Akıncioğlu, Z. Koksals, F. Turkan and S. Alwasel (2016). The effect of caffeic acid phenethyl ester (CAPE) metabolic enzymes including acetylcholinesterase, butyrylcholinesterase, glutathione s-transferase, lactoperoxidase and carbonic anhydrase isoenzymes I, II, IX and XII, *J. Enzyme Inhib. Med. Chem.* **31(6)**, 1095-1101.
- [57] A. Maharramov, M. Kurbanova, P. Taslimi, Y. Demir, A. Safarova, E. Huseyinov, A. Sujayev, S.H. Alwasel and İ. Gulcin (2021). Synthesis, characterization, crystal structure and bioactivities of novel enamine and pyrrole derivatives endowed with acetylcholinesterase,  $\alpha$ -glycosidase and human carbonic anhydrase inhibition effects, *Org. Commun.* **14(2)**, 144-156
- [58] K. Küçüköglu, H.İ. Gül, P. Taslimi, İ. Gülçin and C.T. Supuran, (2019). Investigation of inhibitory properties of some hydrazone compounds on hCA I, hCA II and AChE enzymes, *Bioorg. Chem.* **86**, 316-321.

Secondary metabolites and antioxidant and enzyme inhibitory activities of *Verbascum speciosum*

- [59] Z. Özer, S. Çarıkçı, H. Yılmaz, T. Kılıç, T. Dirmenci and A.C. Gören (2020). Determination of secondary metabolites of *Origanum vulgare* subsp. *hirtum* and *O. vulgare* subsp. *vulgare* by LC-MS/MS, *J. Chem. Metrol.* **14**, 25-34.
- [60] M. Topal (2020). Secondary metabolites of ethanol extracts of *Pinus sylvestris* cones from eastern Anatolia and their antioxidant, cholinesterase and alpha-glucosidase activities, *Rec. Nat. Prod.* **14**, 129-138.
- [61] B. Magnusson and U. Örnemark (eds.) Eurachem Guide: The fitness for purpose of analytical methods-a laboratory guide to method validation and related topics, 2nd ed. 2014.
- [62] F. Akbaş, A. Özyayın, E. Polat and İ. Onaran (2020). *Lucilia sericata* larval secretions stimulating wound healing effects on rat dermal fibroblast cells, *Rec. Nat. Prod.* **14**(5), 340-354.
- [63] M. Isık, M. Korkmaz, E. Bursal, İ. Gülçin, E. Köksal and H. Tohma (2015). Determination of antioxidant properties of *Gypsophila bitlisensis*, *Int. J. Pharmacol.* **11**(4), 366-371.
- [64] M. Elmastaş, İ. Gülçin, L. Öztürk and İ. Gökçe, (2005). Investigation of antioxidant properties of spearmint (*Mentha spicata* L.), *Asian J. Chem.* **17**(1), 137-148.
- [65] J.H. Bae, Y.J. Park, J. Namiesnik, İ. Gülçin, T.C. Kim, H.C. Kim, B.G. Heo, S. Gorinstein and Y.G. Ku (2016). Effects of artificial lighting on bioactivity of sweet red pepper (*Capsicum annum* L.), *Int. J. Food Sci. Technol.* **51**(6), 1378-1385.
- [66] Z. Bingöl, H. Kızıldaş, A.C. Gören, L. Polat Köse, M. Topal, L. Durmaz, S.H. Alwasel and İ. Gulçin (2021). Antidiabetic, anticholinergic and antioxidant activities of aerial parts of shaggy bindweed (*Convolvulus betonicifolia* Miller subsp.)-profiling of phenolic compounds by LC-HRMS, *Heliyon* **7**, e06986.
- [67] H. Kızıldaş, Z. Bingol, A.C. Goren, S.H. Alwasel and İ. Gülçin (2021). Anticholinergic, antidiabetic and antioxidant activities of *Ferula orientalis* L.-Analysis of its polyphenol contents by LC-HRMS, *Rec. Nat. Prod.* **15**(6), 513-528.
- [68] K. Nofouzi, R. Mahmudi, K. Tahapour, E. Amini and K. Yousefi (2016). *Verbascum speciosum* methanolic extract: Phytochemical components and antibacterial properties, *J. Essent. Oil-Bearing Plant.* **19**(2), 499–505.
- [69] I. Gulcin, A.Z. Tel and E. Kirecci (2008). Antioxidant, antimicrobial, antifungal, and antiradical activities of *Cyclotrichium niveum* (BOISS.) Manden and Scheng, *Int. J. Food Prop.* **11**(2), 450-471.
- [70] D. Gupta (2015). Methods for determination of antioxidant capacity: A review, *Int. J. Pharm. Sci. Res.* **6**(2), 546–566.
- [71] H. Kızıldaş, Z. Bingol, A.C. Goren, S.M. Pınar, S.H. Alwasel and İ. Gülçin (2021). LC-HRMS profiling of phytochemicals, antidiabetic, anticholinergic and antioxidant activities of evaporated ethanol extract of *Astragalus brachycalyx* FISCHER. *J. Chem. Metrol.* **15**(2), 135-151.
- [72] L. Polat Köse and İ. Gulçin (2021). Evaluation of the antioxidant and antiradical properties of some phyto and mammalian lignans, *Molecules* **26**, 7099.
- [73] R. Apak, A. Calokerinos, S. Gorinstein, M.A. Segundo, D.B. Hibbert, İ. Gülçin, S. Demirci Çekiç, K. Güçlü, M. Özyürek, S. Esin Çelik, L.M. Magalhaes and P. Arancibia-Avila (2022). Methods to evaluate the scavenging activity of antioxidants toward reactive oxygen and nitrogen species, *Pure Appl. Chem.* **94**(1), 87-144.
- [74] İ. Gulçin, Z. Bingöl, P. Taslimi, A.C. Gören, S.H. Alwasel and A.Z. Tel (2022). Polyphenol contents, potential antioxidant, anticholinergic and antidiabetic properties of mountain mint (*Cyclotrichium leucotrichum*), *Chem. Biodivers.* **19**(3), e202100775.
- [75] L. Durmaz, A. Ertürk, M. Akyüz, L. Polat Köse, E.M. Uc, Z. Bingöl, R. Sağlamtaş, S. Alwasel and İ. Gulçin (2022). Screening of carbonic anhydrase, acetylcholinesterase, butyrylcholinesterase and  $\alpha$ -glycosidase enzymes inhibition effects and antioxidant activity of coumestrol, *Molecules* **27**, 3091.
- [76] M. Topal and İ. Gulçin (2022). Evaluation of the *in vitro* antioxidant, antidiabetic and anticholinergic properties of rosmarinic acid from rosemary (*Rosmarinus officinalis* L.), *Biocat. Agric. Biotechnol.* **43**, 102417.
- [77] L. Durmaz, H. Kiziltas, L. Guven, H. Karagecili, S. Alwasel and İ. Gulçin (2022). Antioxidant, antidiabetic, anticholinergic, and antiglaucoma effects of magnoflorine, *Molecules* **27**, 5902.
- [78] M.S. Özasan, R. Sağlamtaş, Y. Demir, Y. Genç, İ. Saraçoğlu and İ. Gulçin (2022). Isolation of some phenolic compounds from *Plantago subulata* L. and determination of its antidiabetic, anticholinesterase, antiepileptic and antioxidant activity, *Chem. Biodivers.* **19**(8), e202200280.
- [79] T. Ak and İ. Gülçin (2008). Antioxidant and radical scavenging properties of curcumin, *Chem. Biol. Interact.* **174**, 27-37.
- [80] İ. Gülçin, M. Elmastaş and H.Y. Aboul-Enein (2007). Determination of antioxidant and radical scavenging activity of basil (*Ocimum basilicum*) assayed by different methodologies, *Phytother. Res.* **21**(4), 354-361.
- [81] İ. Gülçin, V. Mshvildadze, A. Gepdiremen and R. Elias (2006). Screening of antioxidant and antiradical activity of monodesmosides and crude extract from *Leontice smirnowii* Tuber., *Phytomedicine*, **13**, 343-351.

Secondary metabolites and antioxidant and enzyme inhibitory activities of *Verbascum speciosum*

- [82] İ. Gülçin (2006). Antioxidant activity of caffeic acid (3,4-dihydroxycinnamic acid), *Toxicology*, **217**(2-3), 213-220.
- [83] İ. Gülçin (2006). Antioxidant and antiradical activities of L-Carnitine, *Life Sci.* **78**(8), 803–811.
- [84] K. Nofouzi (2015). Study on the antioxidant activity and *in vitro* antifungal activity of *Verbascum speciosum* methanolic extract, *Mycol. Res.* **2**(2), 97–103.
- [85] G. Öztürk, H.G. Ağalar, G. Yıldız, F. Göger and N. Kirimer (2019). Biological activities and luteolin derivatives of *Verbascum eskisehirensis* karavel., Ocak & Ekici, *Marmara Pharm. J.* **23**(3), 532–542.
- [86] H. Şenol, P. Tulay, M.Ç. Ergören, A. Hanoğlu, I. Çalış and G. Mocan (2021). Cytotoxic effects of verbascoside on MCF-7 and MDA-MB-231, *Turk. J. Pharm. Sci.* **18**(5), 637–644.
- [87] M. Galleano, O. Pechanova and G. Fraga (2010) Hypertension, nitric oxide, oxidants, and dietary plant polyphenols, *Curr. Pharm. Biotechnol.* **11**(8), 837-848.
- [88] E. Bursal, P. Taslimi, A. Gören and I. Gulcin (2020). Assessments of anticholinergic, antidiabetic, antioxidant activities and phenolic content of *Stachys annua*, *Biocat. Agric. Biotechnol.* **28**, 101711.
- [89] M. Rudrapal, S.J. Khairnar, J. Khan, A. Bin Dukayl, M.A. Ansari, M.N. Alomary, F.M. Alshabrmi, S. Palai, P.K. Deb and R. Devi (2022). Dietary polyphenols and their role in oxidative stress-induced human diseases: Insights into protective effects, antioxidant potentials and mechanism(s) of action, *Front. Pharmacol.* **13**, 806470.
- [90] A. Innocenti, S.B. Öztürk Sarıkaya, İ. Gülçin and C.T. Supuran (2010). Carbonic anhydrase inhibitors. Inhibition of mammalian isoforms I-XIV with a series of natural product polyphenols and phenolic acids, *Bioorg. Med. Chem.* **18**(6), 2159–2164.
- [91] Y. Chtourou, M. Morjen, R. Ammar, R. Mhiri, M. Jemaa, I. ELBini-Dhouib, H. Fetoui, N. Srairi-Abid, N. Marrakchi and J. Jebali (2022). Investigation of the renal protective effect of combined dietary polyphenols in streptozotocin-induced diabetic aged rats, *Nutrients* **14**(14), 2867.
- [92] I. Gulcin, E. Bursal, H.M. Şehitoğlu, M. Bilsel and A.C. Gören (2010). Polyphenol contents and antioxidant activity of lyophilized aqueous extract of propolis from Erzurum, Turkey, *Food Chem. Toxicol.* **48** (8-9), 2227-2238.
- [93] İ. Gülçin, Ö.İ. Küfrevioğlu, M. Oktay and M.E. Büyükokuroğlu, (2004). Antioxidant, antimicrobial, antiulcer and analgesic activities of nettle (*Urtica dioica* L.), *J. Ethnopharmacol.* **90**, 205-215.
- [94] İ. Gülçin, İ.G. Şat, Ş. Beydemir, M. Elmastaş and Ö.İ. Küfrevioğlu (2004). Comparison of antioxidant activity of clove (*Eugenia caryophyllata* Thunb) buds and lavender (*Lavandula stoechas* L.), *Food Chem.* **87**, 393-400.
- [95] T. Artunç, A. Menzek, P. Taslimi, I. Gulcin, C. Kazaz and E. Şahin (2020). Synthesis and antioxidant activities of phenol derivatives from 1,6-bis(dimethoxyphenyl)hexane-1,6-dione, *Bioorg. Chem.* **100**, 103884.
- [96] H. Teng, L. Chen, T. Fang, B. Yuan and Q. Lin (2017). Rb2 inhibits  $\alpha$ -glucosidase and regulates glucose metabolism by activating AMPK pathways in HepG2 cells, *J. Funct. Foods.* **28**, 306–313.
- [97] M. Torres-Naranjo, G. Suárez A, Gilardoni, L. Cartuche, P. Flores and V. Morocho (2016). Chemical constituents of *Muehlenbeckia tamnifolia* (Kunth) meisn (Polygonaceae) and its *in vitro*  $\alpha$ -amilase and  $\alpha$ -glucosidase inhibitory activities, *Molecules* **21**(11), 1461.
- [98] M, Gümüş, Ş.N. Babacan, Y. Demir, Y. Sert, İ. Koca and İ. Gülçin (2022). Discovery of sulfadrag-pyrrole conjugates as carbonic anhydrase and acetylcholinesterase inhibitors, *Arch. Pharm.* **355**(1), e2100242.
- [99] P. Taslimi and İ. Gulçin (2018). Antioxidant and anticholinergic properties of olivetol, *J. Food Biochem.* **42**(3), e12516.
- [100] S. Bal, Ö. Demirci, B. Şen, P. Taslimi, A. Aktaş, Y. Gök, M., Aygun and İ. Gülçin (2021). PEPPSI type Pd (II)NHC complexes bearing chloro-/fluorobenzyl group: Synthesis, characterization, crystal structures,  $\alpha$ -glycosidase and acetylcholinesterase inhibitory properties, *Polyhedron* **198**, 115060.
- [101] Y. Kaya, A. Erçağ, Y. Zorlu, Y. Demir and İ. Gülçin (2022). New Pd (II) complexes of the bithiocarbohydrazones derived from isatin and disubstituted salicylaldehydes: Synthesis, characterization, crystal structures and inhibitory properties against some metabolic enzymes, *J. Biol. Inorg. Chem.* **27**(2), 271-281.
- [102] R.E. Bora, H. Genç Bilgiçli, E.M. Üç, M.A. Alagöz, M. Zengin and İ. Gulçin (2022). Synthesis, characterization, evaluation of metabolic enzyme inhibitors and *in silico* studies of thymol based 2-amino thiol and sulfonic acid compounds, *Chem. Biol. Interact.* **366**, 110134.
- [103] A. Aktaş, G. Yaklı, Y. Demir, İ. Gulçin, M. Aygün and Y. Gök (2022). The palladium-based complexes bearing 1,3-dibenzylbenzimidazolium with morpholine, triphenylphosphine, and pyridine derivate ligands: synthesis, characterization, structure and enzyme inhibitions, *Heliyon* **8**, e10625.
- [104] Z. Xiao, R. Storms and A. Tsang (2006). A quantitative starch-iodine method for measuring alpha-amylase and glucoamylase activities, *Anal. Biochem.* **351**(1), 146–148.

Secondary metabolites and antioxidant and enzyme inhibitory activities of *Verbascum speciosum*

- [105] D. Aktaş Anıl, M.F. Polat, R. Sağlamtas, A.H. Tarikoğulları, M.A. Alagöz, İ. Gülçin, O. Algül and S. Burmaoğlu, (2022). Exploring enzyme inhibition profiles of novel halogenated chalcone derivatives on some metabolic enzymes: Synthesis, characterization and molecular modeling studies, *Comp. Biol. Chem.* **100**, 107748.
- [106] İ. Gulçin, O.V. Petrova, P. Taslimi, S.F. Malysheva, E.Y. Schmidt, L.N. Sobenina, N.K. Gusarova, B.A. Trofimov, B. Tuzun, V.M. Farzaliyev, E. Sujayev, S. Alwasel and A.R. Sujayev, (2022). Synthesis, characterization, molecular docking, acetylcholinesterase and  $\alpha$ -glycosidase inhibition profiles of nitrogen-based novel heterocyclic compounds, *ChemistrySelect*, **7(19)**, e20220037.

**A C G**  
**publications**

© 2021 ACG Publications