

Eupatorione A, an Unusual Sesquiterpenoid from the Aerial Parts of *Eupatorium adenophorum*

Hao Geng ^{*1}, Wen-Jie Gu ¹, Jia-Hui Luo ¹, Yong-Xun Yang ^{*2},
and Yang Yu ^{*3,4}

¹Department of Sciences, Xichang University, Xichang, Sichuan 615000, P. R. China

²Department of Animal Sciences, Xichang University, Xichang, Sichuan 615000, P. R. China

³School of Pharmacy, Anhui University of Chinese Medicine, Hefei, 230012, P. R. China

⁴Institute of Medicinal Chemistry, Anhui Academy of Chinese Medicine, Hefei 230012, P. R.
China

(Received June 22, 2023; Revised August 24, 2023; Accepted August 27, 2023)

Abstract: Herein, the chemical analysis of petroleum ether extract of *Eupatorium adenophorum* was conducted which led to the identification of a previously unreported sesquiterpenoid, named eupatorione A (**1**). The structure of eupatorione A (**1**) was elucidated through comprehensive spectroscopic analysis, including 1D, 2D-NMR techniques, as well as HRMS. Additionally, X-ray diffraction was employed to further confirm the structure. The anti-inflammatory activity of compound **1** was assessed by measuring its inhibitory effects on the production of nitric oxide (NO), induced by lipopolysaccharide (LPS) in RAW264.7 LPS-activated macrophages.

Keywords: Compositae; *Eupatorium adenophorum*; sesquiterpenoid; chemical structure. © 2023 ACG Publications. All rights reserved.

1. Plant Source

The aerial parts of *Eupatorium adenophorum* Spreng. (Compositae) were procured from Xichang, located at GPS coordinates of 30°52'N/104°44'E, within Sichuan province, during the wet season of July 2021. The plant was identified by Prof. Qing-Shan Yang from Anhui University of Chinese Medicine. An authentic sample with the designation EA20210711 was preserved at Anhui University of Traditional Chinese Medicine for reference.

2. Previous Studies

Plants can produce a multitude of natural products, many of which play a significant role in developing lead compounds in the treatment of diseases and the production of plant-derived

* Corresponding authors: E- Mail: genghao1709@163.com (H. Geng), xcc03100032@xcc.edu.cn (Y. X. Yang), yyang71@ahctm.edu.cn (Y. Yu).

Eupatorione A, an unusual sesquiterpenoid

insecticides. The fascinating structures and diverse biological activities of these compounds have garnered substantial interest from researchers in both the fields of natural products and medicinal chemistry [1–2]. Terpenoids, characterized by an extensive array of chemical structures and biological functions, are the most abundant compounds in the family of natural products. To date, over 95000 terpenoids have been documented [3–4]. *Eupatorium adenophorum* Spreng., a member of the Compositae family (also known as Asteraceae), is a significant invasive plant [5]. However, this plant exhibits a rich content of terpenoids. A previous phytochemical study revealed that the aerial parts of this plant possessed an abundance of sesquiterpenoids. These sesquiterpenoids demonstrated noteworthy activities such as antibacterial and antiviral activities [6–9]. In our effort to discover bioactive sesquiterpenoids from this plant, we have identified a sesquiterpenoid possessing an unprecedented 5/5/6 tricyclic rare skeleton as depicted in Figure 1.

3. Present Study

The air-dried and powdered aerial parts of *E. adenophorum* (12 kg) were finely ground and immersed in petroleum ether (25 L) at room temperature for 48 h. This extraction was conducted thrice under identical conditions. The combined extract was evaporated under reduced pressure to obtain an oily residue (162 g), which was subjected to silica-gel column chromatography (CC). The column was eluted using a stepwise gradient of petroleum ether (PE) and ethyl acetate (10:0, 9:1, 8:2, 7:3, 6:4, 5:5, 0:10, v/v). The elution led to the separation into four distinct fractions labeled as Frs. I–IV). Fr. II (48 g) was fractionated on silica-gel CC with PE/acetone (from 10:1 to 2:1, v/v) as the eluent to give three sub-fractions (Frs. IIA–IIC). Fr. IIB (6.4 g) was passed through a Sephadex LH-20 column, eluted with acetone, resulting in additional sub-fractions (Fr. IIB1–Fr. IIB2). Subsequently, Fr. IIB2 (700 mg) was isolated through silica-gel CC, employing a mixture of PE and acetone (8:1, v/v) as the eluent. Further purification was achieved by subjecting it to a Sephadex LH-20 column eluted with acetone, followed by the separation using semi-preparative RP-HPLC using MeOH-H₂O (70:30, v/v) as the mobile phase. This process yielded compound 1 (*t_R* 28.9 min, 4.6 mg).

Equipment: Preparative HPLC was performed on an Ruihe with a Zorbax SB-C18 column (5 μm, 9.4 × 150 mm, Agilent, America) with a flow rate of 3.5 mL/min. The ¹H-NMR (600 MHz) and ¹³C-NMR (150 MHz) spectra were recorded on Bruker AVANCE III 600 MHz. UV spectrum was obtained on a Shimadzu 2700 or 2401 PC double-beam spectrophotometer. IR spectra were recorded on a Bruker-Tensor-27 spectrometer with KBr pellets. Mass spectra were obtained on Agilent G6230 spectrometer. Optical rotation was measured with a Horiba SEPA-300 polarimeter. X-ray analysis was conducted on a Bruker SMART APEX CCD crystallography system.

Measurement of Nitric Oxide (NO) Production: NO production was assayed in supernatants of cultured RAW264.7 cells by using an NO assays kit (Beyotime Institute of Biotechnology, Haimen, China). Cells were seeded in 96 well culture plates and pretreated with tested compounds (50 μM) and stimulated with LPS (1 lg/mL) for 24 h. The supernatant was mixed with an equal volume of Griess reagent (1% sulfanilamide, 0.1% naphthylethylenediamine dihydrochloride, and 2.5% phosphoric acid) and incubated at room temperature for 24 h [10]. The concentration of nitrite was measured by reading the absorbance at 570 nm. L-NMMA was used as a positive control [11–12].

Eupatorione A (I): Colorless needle crystal; $[\alpha]_D^{20} = -6.7$ (*c* = 0.01, MeOH); UV (MeOH): λ_{\max} (log ϵ): 251 (3.78); IR ν_{\max} (KBr): = 3449, 2958, 2924, 2852, 1716, 1645, 1454, 1383, 1150, 1075 cm⁻¹; ¹H NMR (600 MHz, acetone-*d*₆): δ (ppm) = 1.92 (3H, s, Me-1), 2.14 (3H, s, Me-3), 2.64 (1H, d, *J* = 5.0 Hz, H-6), 1.70 (1H, m, H-7), 1.89 (1H, m, H-8), 3.50 (1H, m, H-9), 2.27 (1H, m, H-10), 2.92 (1H, m, H-11a), 2.49 (1H, m, H-11b), 1.32 (3H, s, Me-14), 0.85 (3H, d, *J* = 7.1 Hz, Me-15), 4.19 (1H, s, OH); ¹³C NMR (150 MHz, acetone-*d*₆): δ (ppm) = 18.3 (CH₃, Me-15), 20.4 (CH₃, Me-3), 23.0 (CH₃, Me-1), 27.0 (CH₃, Me-14), 35.8 (CH, C-7), 45.2 (CH, C-8), 45.5 (CH₂, C-11), 48.1 (CH, C-9), 58.1 (CH, C-10), 59.1 (CH, C-6), 76.6 (C, C-13), 135.7 (C, C-4), 143.9 (C, C-2), 203.3 (C, C-5), 213.5 (C,

C-12); HRESIMS m/z 249.1487[M+H]⁺ (calcd. 249.1485 for C₁₅H₂₁O₃). The crystal data (CCDC: 2271496) of **1** see Table S1.




Figure 1. Structure of eupatorione A (**1**) isolated from *E. adenophorum*

Compound **1** was obtained as a colorless needle-shaped crystal. The IR spectrum (Figure S8) of **1** exhibited absorption bands for hydroxyl groups (3449 cm⁻¹) and α , β -unsaturated ketone ($\nu_{C=O}$ 1715, $\nu_{C=C}$ 1645, ν_{C-O} 1150 cm⁻¹). HRESIMS recorded an ion peak at m/z 249.1487 [M+H]⁺ (cal. 249.1485), its molecular formula was determined as C₁₅H₂₁O₃, signifying a total of six degrees of unsaturation. Further analysis of the ¹H and ¹³C NMR data (Figure S1 and S2), along with the assistance of DEPT experiments, revealed the presence of four methyl groups (including one bimodal methyl), one methylene group, five methine groups, five quaternary carbons (corresponding to two olefinic carbons, two ketone carbonyls, and one oxygenated quaternary carbon, respectively). The comprehensive analysis of the NMR data indicated that compound **1** should be a tricyclic natural compound. The key HMBC interactions illustrated in Figure 2 provide valuable insights into the structure of compound **1**. The correlations are as follows: correlation of Me-1 (δ_H 1.92) with C-2/C-3/C-4, Me-3 (δ_H 2.14) with C-1/C-2/C-4, Me-14 (δ_H 1.32) with C-10/C-12/C-13, Me-15 (δ_H 0.85) with C-6/C-7/C-8, H-6 (δ_H 2.64) with C-4/C-8/C-9/C-15, H-7 (δ_H 1.70) with C-5/C-9/C-10/C-11, H-8 (δ_H 1.89) with C-4/C-6/C-10/C-12/C-15, H-9 (δ_H 3.50) with C-2/C-4/C-5/C-6/C-7/C-11, H-10 (δ_H 2.27) with C-4/C-5/C-7/C-8/C-14, H-11 (δ_H 2.92, 2.49) with C-7/C-9/C-12/C-13, 13-OH (δ_H 4.19) with C-10/C-12/C-13/C-14. These correlations conclusively establish the existence of a 5/5/6 tricyclic rare skeleton for compound **1**. The apparent 1H-1H COSY correlations (Figure 2) of H-6 with H-7/H-10, H-7 with H-8/H-15, H-8 with H-9/H-11, H-10 with H-6/H-9 further confirmed the planar structure of the compound. While the ROESY spectrum only displayed limited correlations indicating the correlation of H-6 with H-9/Me-15, and Me-14 with H-9/H-10. Through the solvent volatilization, a needle-shaped crystal of **1** was obtained from a mixture of methanol and water (10:1, v/v). Subsequently, the structure, including the absolute configuration (6*S*, 7*S*, 8*S*, 9*R*, 10*S*, 13*R*), was unambiguously established through X-ray diffraction employing Cu K α radiation (Figure 3). Consequently, compound **1** has been identified as shown in Figure 1, and designated as Eupatorione A.




Figure 2. Important ¹H-¹H-COSY and HMBC correlations for **1**

Eupatorione A, an unusual sesquiterpenoid




Figure 3. X-ray crystallographic structure of compound **1**

Eupatorione A (**1**) was tested for anti-inflammatory activity by measuring the inhibitory effects on LPS-induced NO production in RAW264.7 LPS-activated macrophages. At 50 μ M, compound **1** showed relatively weak anti-inflammatory activity with an inhibition rate of 30.56%. The results are summarized in Table 1.

Table 1. Inhibitory rate of LPS-induced NO production in RAW246.7 cells treated with compound **1**

Compound	concentration (μ M)	NO Inhibitory rate (%)
1	50	30.56 \pm 0.48
L-NMMA	50	55.47 \pm 2.27

Acknowledgments

This work was supported by the National Natural Science Foundation of China (32100324, 81860758); High-level Talents Support project of Anhui University of Chinese Medicine (2023rcZD005); National Training Program of Innovation and Entrepreneurship for Undergraduates (202210628010, 202310628020X, 202310628071X), the Doctoral Scientific Research and Double High Talent Project of Xichang University (YBZ202114, LGLZ202302), the Key Research and Development Program of Science and Technology Bureau of Liangshan Yi Autonomous Prefecture (21ZDYF0019).

Supporting Information

Supporting Information accompanies this paper on <http://www.acgpubs.org/journal/records-of-natural-products>

ORCID

Hao Geng: [0000-0002-7917-6839](https://orcid.org/0000-0002-7917-6839)

Wen-Jie Gu: [0009-0000-7183-0587](https://orcid.org/0009-0000-7183-0587)

Jia-Hui Luo: [0009-0005-1696-5020](https://orcid.org/0009-0005-1696-5020)

Yong-Xun Yang: [0000-0001-5688-8991](https://orcid.org/0000-0001-5688-8991)

Yang Yu: [0000-0002-9477-2313](https://orcid.org/0000-0002-9477-2313)

References

- [1] R. A. Dixon (2001). Natural products and plant disease resistance, *Nature* **411**, 843-847.
- [2] A. A. Elkordy, R. R. Haj-Ahmad, A. S. Awaad and R. M. Zaki (2021). An overview on natural product drug formulations from conventional medicines to nanomedicines: past, present and future, *J. of Drug Deliv. Sci. Tec.* **63**, 102459.
- [3] J. Gershenzon and N. Dudareva (2007). The function of terpene natural products in the natural world, *Nat. Chem. Biol.* **3**, 408-414.

- [4] D. W. Christianson (2017). Structural and chemical biology of terpenoid cyclases. *Chem. Rev.* **117**, 11570-11648.
- [5] Y. Liu, S. H. Luo, J. Hua, D. S. Li, Y. Ling, Q. Luo and S. H. Li (2021). Characterization of defensive cadinenes and a novel sesquiterpene synthase responsible for their biosynthesis from the invasive *Eupatorium adenophorum*, *New Phytol.* **229**, 1740-1754.
- [6] L. He, J. Hou, M. Gan, J. Shi, S. Chantrapromma, H. K. Fun, I. D. Williams and H. H. Y. Sung (2008). Cadinane sesquiterpenes from the leaves of *Eupatorium adenophorum*, *J. Nat. Prod.* **71**, 1485-1488.
- [7] X. Zhao, G. W. Zheng, X. M. Niu, W. Q. Li, F. S. Wang and S. H. Li (2009). Terpenes from *Eupatorium adenophorum* and their allelopathic effects on *Arabidopsis* seeds germination, *J. Agric. Food Chem.* **57**, 478-482.
- [8] W. Shi, S. H. Luo and S. H. Li (2012). Defensive sesquiterpenoids from leaves of *Eupatorium adenophorum*, *Chin. J. Chem.* **30**, 1331-1334.
- [9] G. Simran, S. Ranabir, P. Paramita, N. Gouranga and K. D. Tarun (2022). An updated review on *Eupatorium adenophorum* Spreng. [*Ageratina adenophora* (Spreng.)]: traditional uses, phytochemistry, pharmacological activities and toxicity, *Pharmacol. Res.Mod. Chin. Med.* **2**, 100068.
- [10] L.J. Zheng, L.J. Zhang, J.Y. Huang, X.Q. Zhang, M.Q. Huang, H.Y. Xu, and L. (2022). A new alkaloid from *Ormosia hosiei* Hemsl. et Wils, *Rec. Nat. Prod.* **16(6)**, 579-584.
- [11] W. D. Xie, X. Li, J. H. Zhao, Y. H. Liu and K. H. Row (2012). Abietane diterpenoids from *Isodon inflexus*, *Phytochemistry* **81**, 153-158.
- [12] Y. X. Zhu, H. M. Yan, X. J. Wang, Z. X. Zhang, H. P. Zhang, L. S. Chai, L. Li, J. Qu and Y. Li (2019). Micranthanosides I and II, two novel 1,10-secograyanane diterpenoids and their antinociceptive analogues from the leaves and twigs of *Rhododendron micranthum*, *RSC Adv.* **9**, 18439-18450.

ACG
publications

© 2023 ACG Publications