

9,11-Cycloneren-3,7-diol: a New Cyclonerane Sesquiterpene from the Marine-Sediment-Derived Fungus *Trichoderma harzianum*

WH-22

Xin Dong , Lin-Chuan Jia , Hai-Quan Chai , Xiao-Xiao Liu 
and Lei Yang 

Qingdao Hiser Hospital Affiliated of Qingdao University (Qingdao Traditional Chinese Medicine),
Qingdao 266033, China

(Received March 19, 2024; Revised April 17, 2024; Accepted April 20, 2024)

Abstract: One new cyclonerane sesquiterpene, 9,11-cycloneren-3,7-diol (**1**), along with four known ones, 11-cycloneren-3,7,10-triol (**2**), 9-cycloneren-3,7,11-triol (**3**), 11-methoxy-9-cycloneren-3,7-diol (**4**), and cyclonerodiol (**5**), were obtained from the extract of marine-derived fungus *Trichoderma harzianum* WH-22. The structure of 9,11-cycloneren-3,7-diol (**1**) was elucidated by comprehensive spectroscopic analysis, including 1D/2D NMR and HRESIMS data. 9,11-Cycloneren-3,7-diol (**1**) exhibited weak cytotoxicity against HeLa and HepG2 cancer cell lines with IC₅₀ values of 68.2 and 59.7 μM, respectively.

Keywords: cyclonerane sesquiterpene; *Trichoderma harzianum*; structure elucidation; cytotoxic activity. © 2024 ACG Publications. All rights reserved.

1. Fungal Source

Trichoderma harzianum WH-22 was isolated from the sediments collected from the coastal zone of Weihai, China in July 2021. It was identified according to morphological characteristics and ITS regions of its rDNA, whose sequence data have been deposited at GenBank (OR534535). The fungus was preserved in Qingdao Hiser Hospital Affiliated of Qingdao University, Qingdao, China, with the number of WH-22.

2. Previous Studies

Secondary metabolites obtained from marine-derived *Trichoderma* spp. exhibited biosynthetic differences with terrestrial-derived ones. A large number of natural products with novel structures and intriguing bioactivities have been discovered from marine-derived fungi, especially for the *Trichoderma* species, which are serving as a repository of new compounds [1,2]. Among them, *T. harzianum* can produce various bioactive secondary metabolites, including terpenoids, macrolides,

*Corresponding authors: E-mail: 15269261998@126.com (X.X. Liu) ylei0128@163.com (L. Yang)

9,11-Cycloneren-3,7-diol: a new cyclonerane sesquiterpene

polyketides and so on, which are an important source for searching new drugs [3]. In addition, previous studies had reported that some compounds were isolated from *Trichoderma* spp. derived from the ocean sediments [4], but the secondary metabolites acquired from marine-sediment-derived *T. harzianum* were rarely reported [5].

3. Present Study

During our chemical investigation for functional secondary metabolites from the marine-derived fungi, one new cyclonerane sesquiterpene, 9,11-cycloneren-3,7-diol (**1**), together with four known ones, 11-cycloneren-3,7,10-triol (**2**), 9-cycloneren-3,7,11-triol (**3**), 11-methoxy-9-cycloneren-3,7-diol (**4**), and cyclonerodiol (**5**), were isolated from *T. harzianum* WH-22 by chromatographic separation and identified via spectroscopic methods. Herein, the isolation, structure elucidation, and cytotoxic activity of compound **1** are described in detail.

The mass culture was performed at 26 °C for 30 days in 60 × 1L Erlenmeyer flasks, each containing 40 g of rice and 70 mL of natural seawater. After fermentation, about 100 mL EtOAc was added into each flask to kill the fungi, and the mycelia were collected, dried, and then exhaustively extracted with EtOAc. The crude extract (80.0 g) was obtained after removing the organic solvent by evaporating under reduced pressure. The extract was subjected to silica gel column chromatography (CC) with step-gradient solvent systems involving petroleum ether (PE)/EtOAc and then CH₂Cl₂/MeOH to gain 10 fractions. Fraction 4 (6.3 g), eluted with PE/EtOAc 2:1, was further purified by CC on RP-18 (MeOH/H₂O, 7:3) and Sephadex LH-20 (MeOH) to yield 9,11-cycloneren-3,7-diol (**1**) (2.1 mg).

9,11-Cycloneren-3,7-diol (1): Colorless oil; $[\alpha]_D^{20} = -25^\circ$ ($c = 0.2$, MeOH); ¹H (500 MHz) and ¹³C (125 MHz) NMR data, see Table 1; HREIMS: m/z 239.2018 $[M + H]^+$ (calcd for C₁₅H₂₇O₂, 239.2011).

Measurement of Cytotoxicity: Cytotoxic assay toward the human tumor cell lines, including HeLa, HepG2, and MCF-7, was carried out with reported methods [6,7]. The cells were exposed to the tested compound with various concentrations (3.125, 6.25, 12.5, 25, 50, and 100 μM) in 96-well plates for 72 h, and the cell viability was determined by the CCK-8 Cell Proliferation and Cytotoxicity Assay Kit, and the absorbance was recorded by a microplate spectrophotometer at 490 nm. Epirubicin was provided as the positive control.




Figure 1. Chemical structures of compounds **1-5** isolated from *T. harzianum* WH-22

Compound **1** was acquired as colorless oil. HRESIMS analysis gave the molecular formula C₁₅H₂₆O₂, suggesting three degrees of unsaturation. The ¹H NMR spectrum exhibited notable proton signals involving four olefinic protons at $\delta = 6.21$ (br d, $J = 15.6$ Hz, H-10), 5.69 (dt, $J = 15.6$ and 7.6 Hz, H-9), 4.92 (br s, H-12a), and 4.91 (br s, H-12b), and four methyls including three methyl singlets at $\delta = 1.86$ (s, H-15), $\delta = 1.27$ (s, H-13), and $\delta = 1.16$ (s, H-14) and one methyl doublet at $\delta = 1.06$ (d, $J = 6.8$ Hz, H-1). The ¹³C NMR spectrum displayed 15 resonances, assigned to four methyls, four methylenes, four methines, and three quaternary carbons by DEPT experiments. Its NMR data was

high similar to those of 11-cycloneren-3,7,10-triol (**2**) [8], except for the presence of signals for a *trans* double bond (C-9 = C-10) and lack of signals for a methylene (C-9) and an oxygenated methine (C-10), which was validated by the COSY correlations of H-9 with H-8 and H-10 and the HMBC correlations from H-12 to C-2, C-11, and C-15. Other COSY and HMBC correlations (Figure 2), including HMBC correlations from H-13 to C-2, C-3, and C-4 and from H-14 to C-6, C-7, and C-8 and COSY correlations of H-1/H-2/H-6/H-5/H-4 and of H-8/H-9/H-10, further confirmed the planar structure of **1**. The geometry of double bond at C-9 was proposed to be *trans* by the large coupling constant ($J = 15.6$ Hz) between H-9 and H-10. H-6 was *syn* to C-1 by the NOE correlation of H-6 with H-1, while C-13 was opposite to C-1 by the NOE correlation of H-2 with H-13 (Figure 2). The relative configuration of C-14 and the absolute configuration of **1** were deduced to be the same as those of cyclonerodiol (**5**) [9,10] and 9-cycloneren-3,7,11-triol (**3**) [8], confirmed by their identical NMR data and specific optical rotation values $[\alpha]_{\text{D}}^{20} -25$ ($c = 0.2$, MeOH) for **1**; $[\alpha]_{\text{D}}^{20} -21$ ($c = 0.1$, MeOH or CDCl_3) for cyclonerodiol; $[\alpha]_{\text{D}}^{20} -22$ ($c = 0.04$, MeOH) for 9-cycloneren-3,7,11-triol].

Table 1. ^1H (500 MHz) and ^{13}C (125 MHz) NMR data of compound **1** (δ in ppm) in CDCl_3

No	δ_{H} (J in Hz)	δ_{C} , type
1	1.06, d ($J = 6.8$)	14.6, CH_3
2	1.64, m	44.5, CH
3		81.5, C
4a	1.69, m	40.5, CH_2
4b	1.59, m	
5a	1.87, m	24.5, CH_2
5b	1.61, m	
6	1.86, m	54.4, CH
7		74.9, C
8a	2.32, dd ($J = 13.9, 7.6$)	44.1, CH_2
8b	2.27, dd ($J = 13.9, 7.9$)	
9	5.69, dt (15.6, 7.6)	125.5, CH
10	6.21, br d (15.6)	137.0, CH
11		142.0, C
12a	4.92, br s	115.6, CH_2
12b	4.91, br s	
13	1.27, s	26.2, CH_3
14	1.16, s	25.6, CH_3
15	1.86, s	18.9, CH_3




Figure 2. Key ^1H - ^1H -COSY, HMBC, and NOE correlations for **1**

9,11-Cycloneren-3,7-diol: a new cyclonerane sesquiterpene

9,11-Cycloneren-3,7-diol (**1**) was evaluated for cytotoxic effect against HeLa, HepG2, and MCF-7 cell lines. The result showed that **1** exhibited weak cytotoxicity against HeLa and HepG2 cells with IC₅₀ values of 68.2 and 59.7 μM, respectively. However, **1** displayed no activity toward MCF-7 cells (IC₅₀ > 100 μM).

Table 2. Cytotoxic activity of 9,11-cycloneren-3,7-diol (IC₅₀, μM)

Compounds	HeLa	HepG2	MCF-7
1	68.2 ± 3.1	59.7 ± 4.6	– ^a
Epirubicin	5.9 ± 0.2	6.2 ± 0.7	5.8 ± 0.4

^a No cytotoxic effect at 100 μM.

Secondary metabolites from marine-sediment-derived *Trichoderma* have been rarely studied. In our chemical investigation towards the marine-sediment-derived fungus *T. harzianum* WH-22, one new cyclonerane sesquiterpene, 9,11-cycloneren-3,7-diol (**1**), and four known ones, 11-cycloneren-3,7,10-triol (**2**), 9-cycloneren-3,7,11-triol (**3**), 11-methoxy-9-cycloneren-3,7-diol (**4**), and cyclonerodiol (**5**), were isolated and elucidated. Cyclonerane sesquiterpenes were always found in some *Trichoderma* spp., such as *T. asperellum* and *T. harzianum*, and they can act as metabolite-mark of *Trichoderma*. In addition, 9,11-cycloneren-3,7-diol (**1**) was evaluated for inhibitory activity against three human tumour cell lines and exhibited cytotoxicity against HeLa and HepG2 cells. This study enriches research contents of marine natural products and makes a contribution to utilization of marine bio-resource.

Acknowledgments

This work was supported by Qingdao Science and Technology Project of Traditional Chinese Medicine (No. 2021-zyyq05).

Supporting Information

Supporting Information accompanies this paper on <http://www.acgpubs.org/journal/records-of-natural-products>

ORCID

Xin Dong: [0009-0009-3899-9351](https://orcid.org/0009-0009-3899-9351)

Lin-Chuan Jia: [0000-0002-3671-9573](https://orcid.org/0000-0002-3671-9573)

Hai-Quan Chai: [0009-0002-1526-7755](https://orcid.org/0009-0002-1526-7755)

Xiao-Xiao Liu: [0000-0001-8385-851X](https://orcid.org/0000-0001-8385-851X)

Lei Yang: [0009-0006-6525-8901](https://orcid.org/0009-0006-6525-8901)

References

- [1] A.R. Carroll, B.R. Copp, R.A. Davis, R.A. Keyzers and M.R. Prinsep (2021). Marine natural products, *Nat. Prod. Rep.* **28**, 362-413.
- [2] D.Q. Su, L.J. Ding and S. He (2018). Marine-derived *Trichoderma* species as a promising source of bioactive secondary metabolites, *Mini-Rev. Med. Chem.* **18**, 1702-1713.
- [3] R. Guo, G. Li, Z. Zhang and X. Peng (2022). Structures and biological activities of secondary metabolites from *Trichoderma harzianum*, *Mar. Drugs* **20**, 701.
- [4] Y.V. Khudyakova, M.V. Pivkin, T.A. Kuznetsova and V.I. Svetashev (2000). Fungi in sediments of the sea of Japan and their biologically active metabolites, *Microbiology* **69**, 608-611.
- [5] S.T. Fang, Y.J. Wang, X.Y. Ma, X.L. Yin and N.Y. Ji (2019). Two new sesquiterpenoids from the marine-sediment-derived fungus *Trichoderma harzianum* P1-4. *Nat. Prod. Res.* **33**, 3127-3133.

Dong *et al.*, *Rec. Nat. Prod.* (202X) X:X XX-XX

- [6] X.L. Yuan, X.Q. Li, K. Xu, X.D. Hou, Z.F. Zhang, L. Xue, X.M. Liu and P. Zhang (2020). Transcriptome profiling and cytological assessments for identifying regulatory pathways associated with diorcinol N-induced autophagy in A3 cells, *Front. Pharmacol.* **11**, 570450.
- [7] X.Z. Wang, P.L. Hou, Y.B. Qu, R.Z. Huang, Y. Feng, S.F. Zhao, Y.L. Ding and Z.X. Liao (2021). Chemical constituents of Tibetan herbal medicine *Pulicaria insignis* and their in vitro cytotoxic activities, *Rec. Nat. Prod.* **15**, 91-102.
- [8] Y.P. Song, X.H. Liu, Z.Z. Shi, F.P. Miao, S.T. Fang and N.Y. Ji (2018). Bisabolane, cyclonerane, and harziane derivatives from the marinealga-endophytic fungus *Trichoderma asperellum* cf44-2, *Phytochemistry* **152**, 45-52.
- [9] D. Laurent, N. Goasdoue, F. Kohler, F. Pellegrin and N. Platzter (1990). Characterization of cyclonerodiol isolated from corn infested by *Fusarium Moniliforme* shed.: one- and two-dimensional ¹H and ¹³C NMR study, *Magn. Reson. Chem.* **28**, 662-664.
- [10] J. Langhanki, K. Rudolph, G. Erkel and T. Opatz (2014). Total synthesis and biological evaluation of the natural product (-)-cyclonerodiol, a new inhibitor of IL-4 signaling, *Org. Biomol. Chem.* **12**, 9707-9715.

A C G
publications

© 2024 ACG Publications