

A New Sterol-Related Metabolite from the Soft Coral

Capnella imbricata

Zheng-Zhe Tsai ^{ID}^{1#}, Chih-Kai Hsu ^{ID}^{1#}, Po-Jen Chen ^{ID}², Yu-Li Chen ^{ID}^{3,4},
Su-Ying Chien ^{ID}⁵, Zhi-Hong Wen ^{ID}^{6,7}, Ping-Jyun Sung ^{ID}^{6,8,9,10,11},
Kuei-Hung Lai ^{ID}^{12,13,14*} and Hsu-Ming Chung ^{ID}^{1*}

¹Department of Applied Chemistry, National Pingtung University, Pingtung 900391, Taiwan

²Clinical Immunology Laboratory, Department of Medical Research, E-Da Hospital, I-Shou University, Kaohsiung 824005, Taiwan

³Research Center for Chinese Herbal Medicine, Graduate Institute of Healthy Industry Technology, College of Human Ecology, Chang Gung University of Science and Technology, Taoyuan 33305, Taiwan

⁴Graduate Institute of Biomedical Sciences, Graduate Institute of Natural Products, College of Medicine, Chang Gung University, Taoyuan 33305, Taiwan

⁵Instrumentation Center, National Taiwan University, Taipei 106319, Taiwan

⁶Department of Marine Biotechnology and Resources, National Sun Yat-sen University, Kaohsiung 804201, Taiwan

⁷Institute of BioPharmaceutical Sciences, National Sun Yat-sen University, Kaohsiung 804201, Taiwan

⁸National Museum of Marine Biology and Aquarium, Pingtung, 944401, Taiwan

⁹Chinese Medicine Research and Development Center, China Medical University Hospital, Taichung, 404394, Taiwan

¹⁰Graduate Institute of Natural Products, Kaohsiung Medical University, Kaohsiung, 807378, Taiwan

¹¹Fu Jen Catholic University, New Taipei City, 242062, Taiwan

¹²PhD Program in Clinical Drug Development of Herbal Medicine, College of Pharmacy, Taipei Medical University, Taipei 11031, Taiwan

¹³Graduate Institute of Pharmacognosy, College of Pharmacy, Taipei Medical University, Taipei 11031, Taiwan

¹⁴Traditional Herbal Medicine Research Center, Taipei Medical University Hospital, Taipei 11031, Taiwan

(Received February 21, 2024; Revised April 17, 2024; Accepted April 18, 2024)

Abstract: Exploratory research was carried out on the ethyl acetate extract derived from the soft coral *Capnella imbricata*, sourced from the southeastern waters of Taiwan. This investigation yielded the isolation of three sterol-related compounds (**1–3**). Among these compounds, one previously unidentified metabolite, designated as 4 β -hydroxy-24-methylene-5-cholesten-7-one (**1**), was discovered, along with two known metabolites, namely, 3 β -hydroxy-24-methylene-5-cholesten-7-one (**2**) and gorgostan-5,25-dien-3 β -ol (**3**). The structures of these isolated metabolites were determined through comprehensive spectroscopic analyses. Furthermore, the relative stereochemistry of metabolite **3** was established for the first time in this study using single-crystal X-ray diffraction analysis. The potential anti-inflammatory properties of metabolites **1–3** were evaluated by investigating their ability to suppress the expression of inducible nitric oxide synthase (iNOS) and cyclooxygenase-2 (COX-2) proteins in lipopolysaccharide (LPS)-induced RAW 264.7 macrophage cells.

Keywords: *Capnella imbricata*; sterol; X-ray; iNOS; COX-2. © 2024 ACG Publications. All rights reserved.

*Corresponding author: E-Mail: kueihunglai@tmu.edu.tw (K.-H. Lai); shiuamin@mail.nptu.edu.tw (H.M Chung)

These authors contributed equally to this work and should be considered co-first authors.

Sterols from *Capnella imbricata***1. Animal Material**

In June 2017, specimens of the soft coral *C. imbricata* were collected manually off the coast of Orchid Island (Lanyu Island), Taiwan, using self-contained underwater breathing apparatus. Following collection, these samples were carefully preserved in a freezer at $-20\text{ }^{\circ}\text{C}$ until the extraction process. To ensure proper documentation, a voucher specimen labeled with the voucher number NMMBA-TW-SI-2017-030 was deposited at the National Museum of Marine Biology and Aquarium of Taiwan.

2. Previous Studies

Over the past fifty years, soft corals classified under the genus *Capnella* have been acknowledged as abundant sources of sesquiterpenes, particularly capnellane [1–7], germacrene [8], bicyclogermacrene [8–11], xenicane diterpenes [12], and steroids [12,13]. These secondary metabolites have attracted considerable attention owing to their diverse chemical structures and potential biological activities. *Capnella imbricata* has emerged as a subject of interest due to its varied chemical composition, which yields bioactive compounds. Notably, capnellane-type [1–7] and capgermacrene-type sesquiterpenes [8–10] have been identified, exhibiting a range of biological activities including anti-inflammatory, antibacterial, and cytotoxic properties. These findings highlight the pharmaceutical potential inherent in compounds derived from *C. imbricata*.

3. Present Study

Soft coral specimens, identified as *C. imbricata* and freshly harvested from the shores of Orchid Island, underwent a preservation process involving freezing followed by freeze-drying. The resulting dried material was subsequently ground into a powder and subjected to extraction using ethyl acetate (EtOAc), resulting in the production of an extract. This extract was then subjected to further purification through column chromatography employing silica gel and high-performance liquid chromatography, ultimately leading to the isolation of three sterols: 4 β -hydroxy-24-methylene-5-cholesten-7-one (**1**), 3 β -hydroxy-24-methylene-5-cholesten-7-one (**2**), and gorgostan-5,25-dien-3 β -ol (**3**) (see Figure 1).




Figure 1. The structures of 4 β -hydroxy-24-methylene-5-cholesten-7-one (**1**), 3 β -hydroxy-24-methylene-5-cholesten-7-one (**2**), and gorgostan-5,25-dien-3 β -ol (**3**).


Table 1. ^1H and ^{13}C NMR data for sterol **1**

| Position | ^1H (J in Hz) ^a | ^{13}C ^b , type ^c |
|----------|-------------------------------------|--|
| 1a/b | 1.73 m; 2.04 m | 37.1, CH ₂ |
| 2 | 1.58 m | 32.0, CH ₂ |
| 3a/b | 1.26 m; 2.03 m | 38.5, CH ₂ |
| 4 | 4.36 s | 73.3, CH |
| 5 | | 168.3, C |
| 6 | 5.82 s | 126.3, CH |
| 7 | | 202.0, C |
| 8 | 2.53 m | 39.6, CH |
| 9 | 0.91 m | 53.6, CH |
| 10 | | 38.0, C |
| 11 | 1.50 m | 22.2, CH ₂ |
| 12a/b | 1.17 m; 2.05m | 39.6, CH ₂ |
| 13 | | 42.6, C |
| 14 | 1.35 m | 50.8, CH |
| 15 | 1.65 m | 24.1, CH ₂ |
| 16 | 1.89 m | 28.1, CH ₂ |
| 17 | 1.16 m | 55.9, CH |
| 18 | 0.75 s | 12.0, CH ₃ |
| 19 | 1.38 s | 19.5, CH ₃ |
| 20 | 1.43 m | 35.8, CH |
| 21 | 0.96 d (6.6) ^c | 18.7, CH ₃ |
| 22a/b | 1.16 m; 1.53 m | 31.0, CH ₂ |
| 23a/b | 1.86 m; 2.18 m | 34.6, CH ₂ |
| 24 | | 156.8, C |
| 25 | 2.21 m | 33.8, CH |
| 26 | 1.02 d (3.0) | 21.9, CH ₃ |
| 27 | 1.03 d (3.0) | 22.0, CH ₃ |
| 28a/b | 4.66 s; 4.72 s | 106.0, CH ₂ |

^aSpectroscopic data were recorded at 600 MHz in CDCl₃ at 25 °C.

^bSpectroscopic data was recorded at 150 MHz in CDCl₃ at 25 °C.

^cAttached protons were deduced by HSQC experiments.

**Figure 2.** Key COSY (—), HMBC (⤵), and NOESY (⋯) correlations of sterol **1**.

The ^1H - ^1H correlation spectroscopy (COSY) experiment conducted on sterol **1** facilitated the identification of two spin systems: H₂-1/H₂-2/H₂-3/H-4; H-8/H-9/H₂-11/H₂-12; H-8/H-14/H₂-15/H₂-16/H-17/H-20/H₂-22/H₂-23; H-20/H₃-21 and H₃-26/H-25/H₃-27 (refer to Figure 2). These spin systems were elucidated with the aid of a heteronuclear multiple-bond coherence (HMBC) experiment. Analysis via HMBC revealed correlations between protons and carbons, including H-6/C-4, C-10; H₃-18/C-12, C-13, C-14, C-17; H₃-19/C-1, C-5, C-9, C-10; H₃-21/C-17, C-20, C-22; H₃-26/C-24, C-25, C-27 and H₃-27/C-24, C-25, C-26 (see Figure 2), facilitating the establishment of the primary carbon skeleton of sterol **1**.

Sterols from *Capnella imbricata*

The relative configuration of sterol **1** was determined through observations in a nuclear Overhauser effect spectroscopy (NOESY) experiment (Figure 2). The NOESY correlations between H₃-18/H-20 and H₃-19/H-8 confirmed that these protons were on the same face, leading to their assignment as β-protons. The β-orientation of OH-4 at C-4 was established because H₃-19 showed no correlation with H-4, and a NOESY correlation was observed between H-4/H₂-3 and H-6. By comparing the ¹H NMR data between sterol **1** and the established sterol, 4β-Hydroxy-5-cholesten-7-one [14], we have successfully elucidated the relative configuration of sterol **1**. The stereochemistry analysis indicates that OH-6 is positioned in the β-orientation in sterol **1** (Figure 3).




Figure 3. Comparison of the ¹H NMR chemical shifts in sterol **1**, with those of the known 4β-Hydroxy-5-cholesten-7-one.

In our current study, we not only discovered a new sterol within the soft coral *C. imbricata* but also recognized two previously documented sterols, **2–3**, observed for the first time in this species. By comparing their spectroscopic characteristics with published data, the known sterols **2–3** were identified as 3β-hydroxy-24-methylene-5-cholesten-7-one (**2**) [15] and gorgostan-5,25-dien-3β-ol (**3**) [16].

The structural elucidation of sterol **3** was achieved using single-crystal X-ray diffraction, employing two Bruker D8 Venture diffractometers equipped with dual sources. Sterol **3** was obtained in crystalline form from n-hexane/acetone (11:1) as colorless prisms, adopting a monoclinic crystal system within the P₂₁ space group. The crystal dimensions were measured as 0.517 x 0.175 x 0.066 mm³, with lattice parameters a = 9.5322(4) Å, b = 7.4733(3) Å, c = 36.7188(15) Å, and a volume of 2615.65(19) Å³. The refinement process involved collecting all 22,514 reflections, with the structural model solved using direct methods and refined via a full-matrix least-squares procedure. The refined model yielded final R1 = 0.1095 and wR2 = 0.2894 values for 10,421 observed reflections [*I* > 2σ(*I*)], employing 602 variable parameters. The absolute configuration of Sterol 3 was reliably determined, with Flack's parameter calculated as -0.1(10). A depiction of the structure is provided in the Oak Ridge Thermal Ellipsoid Plot (ORTEP) diagram (see Figure 4). Access to full crystallographic data is available in the CIF file CCDC 2313253, containing supplementary information and can be obtained at no cost from the CCDC website via www.ccdc.cam.ac.uk/structures/.




Figure 4. ORTEP revealing the structure of gorgostan-5,25-dien-3β-ol (**3**)

As part of an *in vitro* assessment of anti-inflammatory activity, Western blot analysis was employed to evaluate the expression of pro-inflammatory iNOS and COX-2 proteins in LPS-stimulated RAW264.7 macrophage cells (refer to Table 2). Sterols **1–3** demonstrated a moderate

reduction in iNOS levels (ranging from 64.409% to 77.200%) at a concentration of 10 μ M, compared to control cells stimulated solely with LPS, without eliciting any cytotoxic effects (data not shown). Moreover, Sterol **3**, when administered at 10 μ M, displayed a weak inhibitory effect (90.365%) on COX-2 protein expression.

Table 2. Effects of sterols **1–3** (at 10 μ M) on LPS-induced iNOS and COX-2 protein expressions in macrophages

| Compounds | iNOS | COX-2 | β -actin |
|------------------|---------------------|---------------------|----------------------|
| vehicle | 1.588 \pm 1.097 | 0.788 \pm 0.586 | 113.619 \pm 13.449 |
| LPS | 100.00 \pm 10.472 | 100.00 \pm 8.441 | 105.505 \pm 5.533 |
| 1 | 64.409 \pm 5.534 | 100.234 \pm 5.171 | 108.451 \pm 10.141 |
| 2 | 77.200 \pm 9.145 | 101.334 \pm 5.930 | 95.592 \pm 9.602 |
| 3 | 73.820 \pm 6.169 | 90.365 \pm 9.600 | 116.316 \pm 12.860 |
| Dex ^a | 63.300 \pm 1.794 | 25.677 \pm 2.626 | 99.454 \pm 13.284 |

^aDexamethasone (DEX, 10 μ M) was used as a positive control.

Acknowledgments

This research was supported by the grants from the National Pingtung University and the Ministry of Science and Technology, Taiwan (Grant Nos: MOST 110-2320-B-153-001) awarded to Hsu-Ming Chung; by the grant from the National Science and Technology Council (MOST 110-2320-B-038-034, MOST 111-2320-B-038-040-MY3; MOST 111-2321-B-255-001), Ministry of Education (DP2-110-21121-01-N-12-03), and Taipei Medical University (TMU109-AE1-B15) awarded to Kuei-Hung Lai. The authors would like to thank Hsiao-Ching Yu and Chao-Lien Ho (The High Valued Instrument Center, National Sun Yat-sen University) for the mass (MS 006500) and NMR (NMR 001100) data collection (NSTC 112-2740-M-110-002); and to the Instrumentation Center, National Taiwan University for providing X-ray facilities (NSTC 112-2731-M-002-001, XRD000200).

Supporting Information

Supporting Information accompanies this paper on <http://www.acgpubs.org/journal/records-of-natural-products>

ORCID

Zheng-Zhe Tsai: [0009-0005-9704-6631](https://orcid.org/0009-0005-9704-6631)

Chih-Kai Hsu: [0009-0003-4853-1695](https://orcid.org/0009-0003-4853-1695)

Po-Jen Chen: [0000-0002-0135-8798](https://orcid.org/0000-0002-0135-8798)

Yu-Li Chen: [0000-0003-2392-1702](https://orcid.org/0000-0003-2392-1702)

Su-Ying Chien: [0009-0008-0931-3149](https://orcid.org/0009-0008-0931-3149)

Zhi-Hong Wen: [0000-0003-4411-044X](https://orcid.org/0000-0003-4411-044X)

Ping-Jyun Sung: [0000-0002-6644-9236](https://orcid.org/0000-0002-6644-9236)

Kuei-Hung Lai: [0000-0001-7220-6356](https://orcid.org/0000-0001-7220-6356)

Hsu-Ming Chung: [0000-0003-1472-9268](https://orcid.org/0000-0003-1472-9268)

References

- [1] Y. M. Sheikh, G. Singy, M. Kaisin, H. Eggert, C. Djerassi, B. Tursch, D. Dalozze and J. C. Braekman (1976). Terpenoids-LXXI. Chemical studies of marine invertebrates-XIV. Four representatives of a novel sesquiterpene class-the capnellane skeleton, *Tetrahedron* **32**, 1171–1178.
- [2] Y. M. Sheikh, C. Djerassi, J. C. Braekman, D. Dalozze, M. Kaisin, B. Tursch and R. Karlsson (1977). Terpenoids-LXXII. Chemical studies of marine invertebrates-XXVI. $\Delta^{9(12)}$ -Capnellene-3 β ,8 β ,10 α ,14-tetrol, a novel polyoxygenated sesquiterpene from the alcyonarian *Capnella imbricata*, *Tetrahedron* **33**, 2115–2117.

Sterols from *Capnella imbricata*

- [3] M. Kaisin, Y. M. Sheikh, L. J. Durham, C. Djerassi, B. Tursch, D. Daloze and J.C. Braekman (1974). Capnellane—a new tricyclic sesquiterpene skeleton from the soft coral *Capnella imbricata*, *Tetrahedron Lett.* **15**, 2239–2242.
- [4] E. Ayanoglu, T. Gebreyesus, C. N. Beechan, C. Djerassi and M. Kaisin (1978). Terpenoids LXXV. $\Delta^{9(12)}$ -capnellene, a new sesquiterpene hydrocarbon from the soft coral *Capnella imbricata*, *Tetrahedron Lett.* **19**, 1671–1674.
- [5] M. Kaisin, J. C. Braekman, D. Daloze and B. Tursch (1985). Novel acetoxycapnellenes from the alcyonacean *Capnella imbricata*, *Tetrahedron* **41**, 1067–1072.
- [6] L. A. Morris, M. Jaspars, K. Adamson, S. Woods and H. M. Wallace (1998). The capnellenes revisited: New structures and new biological activity, *Tetrahedron* **54**, 12953–12958.
- [7] C. H. Chang, Z. H. Wen, S. K. Wang and C. Y. Duh (2008). Capnellenes from the Formosan soft coral *Capnella imbricata*, *J. Nat. Prod.* **71**, 619–621.
- [8] C. S. Phan, T. Kamada, T. Ishii, T. Hamada and C. S. Vairappan (2019). Cytotoxic sesquiterpenoids from soft coral *Capnella imbricata*, *Nat. Prod. Commun.* **14**, 1–4.
- [9] C. S. Phan and C. S. Vairappan (2017). Capgermacrenes D-G, new sesquiterpenoids from a Bornean soft coral, *Capnella imbricata*, *Nat. Prod. Res.* **31**, 742–748.
- [10] T. Ishii, C. S. Phan, T. Kamada and C. S. Vairappan (2016). Capgermacrene C, a new sesquiterpenoid from a Bornean soft coral, *Capnella* sp., *Nat. Prod. Commun.* **11**, 1065–1066.
- [11] C. S. Phan, S. Y. Ng, E. A. Kim, Y. J. Jeon, K. Palaniveloo and C. S. Vairappan (2015). Capgermacrenes A and B, bioactive secondary metabolites from a Bornean soft coral, *Capnella* sp., *Mar. Drugs* **13**, 3103–3115.
- [12] G. J. Hooper and M. T. Davies-Coleman (1995). New metabolites from the South African soft coral *Capnella thyrsoidea*, *Tetrahedron* **51**, 9973–9984.
- [13] A. D. Wright, E. Goclik, G. M. König and G. M. König (2003). Oxygenated analogues of gorgosterol and ergosterol from the soft coral *Capnella lacertiliensis*, *J. Nat. Prod.* **66**, 157–160.
- [14] L. Kohout and J. Fajkoš, Acetolysis of 4 β -methanesulphonyloxy-6 β ,7 $\alpha\beta$ -cyclo-B-homo-5 α -cholestane* (1979). *Collect. Czechoslov. Chem. Commun.* **44**, 3308–3320.
- [15] J. A. Findlay and A. D. Patil (1985). Novel sterols from the finger sponge *Haliclona oculata*, *Can. J. Chem.* **63**, 2406–2410.
- [16] M. E. F. Hegazy, T. A. Mohamed, A. I. Elshamy, A. A. Hassanien, N. S. Abdel-Azim, M. A. Shreadah, I. I. Abdel-gawad, E. M. Elkady and P. W. Paré (2016). A new steroid from the Red Sea soft coral *Lobophytum lobophytum*, *Nat. Prod. Res.* **30**, 340–344.

ACG
publications

© 2024 ACG Publications