

Supporting Information

Rec. Nat. Prod. 10:2 (2016) 154-167

Phytotoxic and Insecticidal Properties of Essential Oils and Extracts of Four *Achillea* Species

Ahmet Çakır^{1*}, Hakan Özer², Tuba Aydın³, Şaban Kordali⁴, Aysema
Tazegül Çavuşoğlu⁵, Tülay Akçin⁶, Ebru Mete⁷ and Adnan Akçin⁸

¹*Kilis 7 Aralık University, Faculty of Science and Literature, Department of Chemistry,
79000, Kilis-Türkiye*

²*Atatürk University, Faculty of Agriculture, Department of Field and Crops, 25240, Erzurum-
Türkiye*

³*Ağrı İbrahim Çeçen University, Faculty of Pharmacy, 04100, Ağrı-Türkiye*

⁴*Atatürk University, Faculty of Agriculture, Department of Plant Protection, 25240, Erzurum-
Türkiye*

⁵*Directorate of Food, Agriculture and Livestock of Tortum District, Erzurum, Türkiye*

⁶*Ondokuz Mayıs University, Faculty of Science and Literature, Department of Biology,
Samsun-Türkiye*

⁷*Atatürk University, Faculty of Science, Department of Chemistry, 25240, Erzurum-Türkiye*

⁸*Amasya University, Faculty of Science and Literature, Department of Biology, 05100,
İpekköy, Amasya-Türkiye*

Table of Contents	Page
Figure S1: Mortality (%) of adults of <i>L. decemlineata</i> after treatment with the essential oils of <i>Achillea</i> species, according to dose exposure and treatment	2-3
Figure S2. Mortality (%) of adults of <i>L. decemlineata</i> after treatment with the essential oils of <i>Achillea</i> species, according to dose exposure and treatment	4






Figure S1. Mortality (%) of adults of *L. decemlineata* after treatment with the essential oils of *Achillea* species, according to dose exposure and treatment






Figure S2. Mortality (%) of adults of *L. decemlineata* after treatment with the essential oils of *Achillea* species, according to dose exposure and treatment