

# Supporting Information

*Rec. Nat. Prod.* (2020) 14:3 219-224

## Chemical Composition, Antibacterial and Cytotoxic Activities of the Essential Oil from *Ficus tikoua* Bur.

Minyi Tian<sup>1,2</sup>, Xiaoge Zhao<sup>2</sup>, Xianghuan Wu<sup>2</sup>, Yi Hong<sup>1</sup>, Qi Chen<sup>1</sup>,

Xiongli Liu<sup>2</sup> and Ying Zhou<sup>2,3\*</sup>

<sup>1</sup> Key laboratory of Plant Resource Conservation and Germplasm Innovation in Mountainous Region (Ministry of Education), Collaborative Innovation Center for Mountain Ecology & Agro-Bioengineering (CICMEAB), College of Life Sciences/Institute of Agro-bioengineering, Guizhou University, Guiyang 550025, P. R. China

<sup>2</sup> Guizhou Engineering Center for Innovative Traditional Chinese Medicine and Ethnic Medicine, Guizhou University, Guiyang 550025, P. R. China

<sup>3</sup> College of Pharmacy, Guizhou University of Traditional Chinese Medicine, Guiyang, 550025, P. R. China

Table of Contents	Page
<b>S1: Experimental Details</b>	2
S1.1: Essential Oil Extraction	2
S1.2: Gas Chromatography (GC) Analysis of Essential Oil	2
S1.3: Gas Chromatography-Mass Spectrometry (GC/MS) Analysis of Essential Oil	2
S1.4: Identification of Chemical Constituents	2
S1.5: Antibacterial Activity Test	2
S1.6: Cytotoxic Activity Test	3
S1.7: Statistical Analysis	4
References	4

\* Corresponding author: [yingzhou71@yeah.net](mailto:yingzhou71@yeah.net) (Y. Zhou)

## **S1: Experimental Details**

### *S1.1: Essential Oil Extraction*

*F. tikoua* was collected from Guizhou Province of China in September 2018. Dry whole plant was obtained by air-dried in the dark at room temperature. The dry whole plant of *F. tikoua* was cut into pieces and placed in a Clevenger-type apparatus and submitted to hydrodistillation (4 h). Essential oil were dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered and stored at 4°C.

### *S1.2: Gas Chromatography (GC) Analysis of Essential Oil*

Analysis of the essential oil was carried out by an Agilent 6890 gas chromatograph system coupled to a flame ionization detector (FID) and equipped with a HP-5MS capillary column (60 m × 0.25 mm × 0.25 μm film thickness). The injection volume was 2 μL (split ratio at 1:10) with helium as the carrier gas (flow rate: 1 mL/min). GC oven temperature was kept at 70°C for 2 min then increased to 180°C at 2°C per min, and programmed to 310°C at a rate of 10°C/min and then finally kept at 310°C for 14 min. The injector temperature was set at 250°C.

### *S1.3: Gas Chromatography-Mass Spectrometry (GC/MS) Analysis of Essential Oil*

The GC-MS analysis was performed on Agilent 6890 GC system fitted with an Agilent 5975C MS and equipped with HP-5MS fused silica column. GC parameters were the same as above. Mass spectra were operated in electron ionization mode at 70 eV. Mass range was from *m/z* 29 to 500 amu.

### *S1.4: Identification of Chemical Constituents*

The percentage of chemical component of the essential oil was calculated by the peak area normalization method. The constituents of the essential oil were identified by their retention time, retention indices relative to n-alkanes (C<sub>8</sub>–C<sub>30</sub>), and as well as by comparison of their mass spectra with those listed in literature, NIST 14 and Wiley 275 databases [1-6].

### *S1.5: Antibacterial Activity Test*

The antibacterial test were performed with six strains of *Enterococcus faecalis* (ATCC 19433), *Staphylococcus aureus* (ATCC 6538P), *Bacillus subtilis* (ATCC 6633), *Pseudomonas aeruginosa* (ATCC 9027), *Escherichia coli* (CICC 10389) and *Proteus vulgaris* (ACCC 11002).

Antibacterial activities of the essential oil and palmitic acid were assessed using the disc agar diffusion method with minor modification [7]. The essential oil and palmitic acid solutions were prepared by diluting with ethyl acetate (100 mg/mL). Filter paper discs (6 mm in diameter) were impregnated with 20 μL of the sample solution and incubated for 24 h at 37°C. The

inhibition zone (including the 6 mm disk) was measured and recorded after the incubation time. All the tests were repeated in triplicate. The result of antibacterial activity test is given in Table S1.

The MIC and MBC values of the essential oil and palmitic acid were determined by the microplate dilution method with slight modification [8]. The essential oil and palmitic acid were initially diluted in DMSO, and later in Mueller-Hinton Broth. The 100  $\mu$ L of twofold dilution of tested sample was transferred into each well and the inoculum was added to all wells. The final concentration of the bacterial cells in the wells was approximately  $5 \times 10^5$  CFU/mL. The 96-well plates were incubated for 24 h at 37°C. 10  $\mu$ L of resazurin aqueous solution (0.01%) was added to 96-well plates as an indicator of microbial growth by detecting the reduction of blue dye resazurin to pink resorufin. The 96-well plates were incubated at 37°C for 2 h in the dark. The MIC was defined as the lowest concentration of sample when the resazurin color changed. To obtain the MBC value, 10  $\mu$ L samples were obtained from the wells (no color change) and subcultured in Mueller-Hinton agar. The MBC was determined as the lowest concentration without any microbial growth after 24 h at 37 °C [9]. Each test was repeated in triplicate. The result of antibacterial activity test is given in Table 2.

**Table S1.** Diameter of the inhibition zones of *F. tikoua* essential oil and palmitic acid using the disc agar diffusion method

Microorganisms	Diameter of the inhibition zones (mm) <sup>a</sup>		
	Essential oil	Palmitic acid	Streptomycin
<b>Gram positive</b>			
<i>Enterococcus faecalis</i> ATCC 19433	9.19 $\pm$ 1.03	8.24 $\pm$ 0.08	8.44 $\pm$ 0.37
<i>Staphylococcus aureus</i> ATCC 6538P	10.59 $\pm$ 0.36	7.12 $\pm$ 0.45	16.83 $\pm$ 0.78
<i>Bacillus subtilis</i> ATCC 6633	10.33 $\pm$ 1.81	6.75 $\pm$ 0.18	17.07 $\pm$ 2.08
<b>Gram negative</b>			
<i>Pseudomonas aeruginosa</i> ATCC 9027	9.17 $\pm$ 1.07	8.88 $\pm$ 0.34	11.52 $\pm$ 3.32
<i>Escherichia coli</i> CICC 10389	9.79 $\pm$ 1.52	8.31 $\pm$ 0.36	15.05 $\pm$ 0.71
<i>Proteus vulgaris</i> ACCC 11002	7.89 $\pm$ 0.49	8.06 $\pm$ 0.57	15.71 $\pm$ 0.77

<sup>a</sup>The diameter of the inhibition zones (mm) were measured including the diameter of the disk (6 mm). The sample solution: Essential oil and palmitic acid were diluted with ethyl acetate, at a concentration of 100 mg/mL, (tested volume: 20  $\mu$ L); Positive control: Streptomycin (tested volume: 20  $\mu$ L, 100  $\mu$ g/mL).

### SI.6: Cytotoxic Activity Test

Human lung adenocarcinoma (A549), non-small cell lung cancer (NCI-H1299), prostatic carcinoma (PC-3), leukemic (K562) cell lines, and normal human fetal lung fibroblasts cell line (MRC-5) were maintained in RPMI 1640 medium (2 mM glutamine, 10% fetal bovine serum, 100 U/mL penicillin and 100 U/mL streptomycin) and incubated in a humidified incubator at 37°C with 5% CO<sub>2</sub> atmosphere. The cytotoxic activity was evaluated by MTT assay with slight modification [10]. The cells were seeded at a density of  $5 \times 10^3$  cells per well in 80  $\mu$ L of culture medium and incubated for 24 h before treatment. The essential oil and palmitic acid were dissolved in DMSO, and afterwards serially double diluted with culture medium. The diluted sample solution (20  $\mu$ L) was added to each well and incubated for 24, 48

and 72 h. After incubation, the medium was removed and 10  $\mu$ L of MTT (5 mg/mL in PBS) was added to each well and incubated for 4 h under the same culture conditions. Then, the MTT was removed and 150  $\mu$ L DMSO was added to each well to solubilize the formazan crystals. After shaking for 10 min at room temperature, the optical density was measured at 490 nm using a microplate spectrophotometer (Bio-Rad Model 680, Hercules, CA, USA). The cytotoxic activity was expressed as the concentration of the essential oil and palmitic acid inhibiting cell growth by 50% (IC<sub>50</sub>). All experiments were performed in triplicate. The result of cytotoxic activity test is given in Table 3 in main text.

### SI.7: Statistical Analysis

The results of the tests were carried out in triplicate and expressed as the means  $\pm$  SD. SPSS software (version 19.0) was used for statistical analysis. Data were compared by oneway analysis of variance (ANOVA) using Tukey's multiple range tests. Differences were considered to be significant at  $p < 0.05$  level.

## References

- [1] R. P. Adams (2001). Identification of essential oil components by gas chromatography/quadrupole mass spectroscopy, *Allured Publ. Corp. Carol Stream, IL*.
- [2] S. M. Sun, G. H. Chung and T. S. Shin (2012). Volatile compounds of the green alga, *Capsosiphon fulvescens*, *J. Appl. Phycol.* **24**, 1003-1013.
- [3] J. A. Pino, R. Marbot and C. Vázquez (2004). Volatile flavor constituents of Karanda (*Carissa carandas* L.) fruit, *J. Essent. Oil Res.* **16**, 432-434.
- [4] P. Evangelia, V. Constantinos, C. Maria and T. Olga (2017). Study of volatile components of *Acacia farnesiana* Willd. flowers, *Rec. Nat. Prod.* **11(5)**, 474-478.
- [5] N. Radulovic, M. Dekic, Z. S. Radic and R. Palic (2011). Chemical composition and antimicrobial activity of the essential oils of *Geranium columbinum* L. and *G. lucidum* L. (*Geraniaceae*), *Turk. J. Chem.* **35**, 499-512.
- [6] C. Demetzos, D. Angelopoulou and D. Perdetzoglou (2002). A comparative study of the essential oils of *Cistus salviifolius* in several populations of Crete (Greece), *Biochem. Syst. Ecol.* **30**, 651-665.
- [7] H. Y. Zhang, Y. Gao and P. X. Lai (2017). Chemical composition, antioxidant, antimicrobial and cytotoxic activities of essential oil from *Premna microphylla* Turczaninow, *Molecules* **22**, 381.
- [8] J. N. Eloff (1998). A sensitive and quick microplate method to determine the minimal inhibitory concentration of plant extracts for bacteria, *Planta Med.* **64**, 711-713.
- [9] M. Mzid, S. Ben Khedir, M. Ben Salem, W. Regaieg and T. Rebai (2017). Antioxidant and antimicrobial activities of ethanol and aqueous extracts from *Urtica urens*, *Pharm. Biol.* **55(1)**, 775-781.
- [10] T. Mosmann (1983). Rapid colorimetric assay for cellular growth and survival: Application to proliferation and cytotoxicity assays, *J. Immunol. Methods* **65**, 55-63.