Supporting Information

J.Chem. Metrol. 14:1 (2020) 1-11

Rapid and easy method for simultaneous measurement of widespread 27 compounds in natural products and foods

Özge Tokul-Ölmez, Bihter Şahin, Cansel Çakır and Mehmet Öztürk*

Muğla Sıtkı Koçman University, Faculty of Science, Department of Chemistry, 48121 Menteşe, Muğla, Türkiye

Table of Contents	Page
Figure S1: The Chemical structures of the reference compounds.	2

© 2020 ACG Publications. All rights reserved.

^{*} Corresponding author E-Mail: mehmetozturk@mu.edu.tr

Figure S1. The Chemical structures of the reference compounds. fumaric acid (1), gallic acid (2), trans-aconitic acid (3), p-benzoquinone (4), pyrocatechol (5), protocatechuic acid (6), unknown (7), 2-(4-hydroxyphenyl) ethanol (8), 4-hydroxybenzoic acid (9), (+)-catechin (10), methyl-p-benzoquinone (11), 6,7-dihydroxy coumarin (12), vanillic acid (13), caffeic acid (14), vanillin (15), 2,4-dihydroxybenzoic acid (16), chlorogenic acid (17), p-coumaric acid (18), ferulic acid (19), coumarin (20), trans-2-hydroxycinnamic acid (21), rutin (22), ellagic acid (23), trans-cinnamic acid (24), naringenin (25), quercetin (26), rosmarinic acid (27), chrysin (28)