

Supporting Information

Rec. Nat. Prod. X:X (2021) XX-XX

Constituents of the Flower of *Maxillaria tenuifolia* and Their Anti-Diabetic Activity

Chia-Ying Li ^{1*}

¹ Department of Applied Chemistry, National Pingtung University, Pingtung City, Pingtung County 900393, Taiwan

Table of Contents		Page
Figure S1: ¹ H-NMR (400 MHz, Acetone-d ₆) spectrum of 1 (3,4-dihydroxy benzoic acid methyl ester)		2
Figure S2: ¹ H-NMR (400 MHz, Acetone-d ₆) spectrum of 2 (flavanthridin)		3
Figure S3: ¹ H-NMR (400 MHz, DMSO-d ₆) spectrum of 3 (vanillic acid)		4
Figure S4: ¹ H-NMR (400 MHz, DMSO-d ₆) spectrum of 4 (mangiferin)		5
Figure S5: ¹³ C-NMR (100 MHz, DMSO-d ₆) spectrum of 4 (mangiferin)		6




Figure S1: $^1\text{H-NMR}$ (400 MHz, Acetone- d_6) spectrum of **1** (3,4-dihydroxy benzoic acid methyl ester)




Figure S2: $^1\text{H-NMR}$ (400 MHz, Acetone- d_6) spectrum of **2** (flavanthridin)




Figure S3: $^1\text{H-NMR}$ (400 MHz, DMSO-d_6) spectrum of **3** (vanillic acid)




Figure S4: $^1\text{H-NMR}$ (400 MHz, DMSO-d_6) spectrum of **4** (mangiferin)




Figure S5: ^{13}C -NMR (100 MHz, DMSO-d_6) spectrum of **4** (mangiferin)