

Supporting Information

Rec. Nat. Prod. X:X (202X) XX-XX

Secondary Metabolites from Marine-Derived Fungus

Aspergillus carneus GXIMD00519

Chun-Ju Lu[#], Zhen-Zhou Tang[#], Zhi-Wei Su, Hai-Yan Li, Geng-Si Zhang,

Cheng-Hai Gao¹, Yong-Hong Liu^{1*} and Xin-Ya Xu^{1*}

*Institute of Marine Drugs, Guangxi University of Chinese Medicine, Nanning, Guangxi 530200,
P.R.China*

Table of Contents	Page
Figure S1: HR-ESI-MS spectrum of 1 (carneusin A)	2
Figure S2: ¹ H-NMR (600 MHz, DMSO- <i>d</i> ₆) spectrum of 1 (carneusin A)	3
Figure S3: ¹³ C-NMR (150 MHz, DMSO- <i>d</i> ₆) spectrum of 1 (carneusin A)	4
Figure S4: HSQC spectrum of 1 (carneusin A)	5
Figure S5: HSQC spectrum of 1 (carneusin A) (From δ_c 18 ppm to δ_c 28 ppm)	6
Figure S6: HSQC spectrum of 1 (carneusin A) (From δ_c 100 ppm to δ_c 116 ppm)	7
Figure S7: HMBC spectrum of 1 (carneusin A)	8
Figure S8: HMBC spectrum of 1 (carneusin A) (From δ_c 15 ppm to δ_c 70 ppm)	9
Figure S9: HMBC spectrum of 1 (carneusin A) (From δ_c 100 ppm to δ_c 120 ppm)	10
Figure S10: HMBC spectrum of 1 (carneusin A) (From δ_c 125 ppm to δ_c 195 ppm)	11
Figure S11: ¹ H- ¹ H COSY spectrum of 1 (carneusin A)	12
Figure S12: NOESY spectrum of 1 (carneusin A)	13
Figure S13: HR-ESI-MS spectrum of 2 (carneusin B)	14
Figure S14: ¹ H-NMR (600 MHz, DMSO- <i>d</i> ₆) spectrum of 2 (carneusin B)	15
Figure S15: ¹³ C-NMR (150 MHz, CDCl ₃) spectrum of 2 (carneusin B)	16
Figure S16: HSQC spectrum of 2 (carneusin B)	17
Figure S17: HMBC spectrum of 2 (carneusin B)	18
Figure S18: HMBC spectrum of 2 (carneusin B) (From δ_c 5 ppm to 45 ppm)	19
Figure S19: HMBC spectrum of 2 (carneusin B) (From δ_c 120 ppm to 190 ppm)	20
Table S1: Relative free energies ^a and equilibrium populations ^b of conformers for (1'S, 4'S, 5'S)- 1 ^c	21
Figure S20: . The optimized structures (left) and the calculated CD spectra of conformers (1'S, 4'S, 5'S)- 1 in MeOH at M06-2X/def2TZVP level (right). $\sigma = 0.22$ eV	21
Table S2: Relative free energies ^a and equilibrium populations ^b of conformers for (6S)- 2 ^c	22
Figure S21: The optimized structures (left) and the calculated CD spectra of conformers (6S)- 2 in MeOH at M06-2X/def2TZVP level (right) . $\sigma = 0.3$ eV	23
Figure S22: DP4+ probabilities (%) for conformers (6S)- 2 (isomer 1) and (6R)- 2 (isomer 2)	24
Figure S23: The Scifinder searching results of compound 1	25
Table S3: The structural comparison of similar compounds with 1	26
Figure S24: The Scifinder searching results of compound 2	27
Table S4: The structural comparison of similar compounds with 2	28
Table S5: The NMR data comparison of similar compounds with 1	29




Figure S1: HR-ESI-MS spectrum of **1** (carneusin A)




Figure S2: $^1\text{H-NMR}$ (600 MHz, $\text{DMSO-}d_6$) spectrum of **1** (carneusin A)




Figure S3: ^{13}C -NMR (150 MHz, $\text{DMSO-}d_6$) spectrum of **1** (carnesin A)




Figure S4: HSQC spectrum of **1** (carnesuin A)




Figure S5: HSQC spectrum of **1** (carnesuin A) (From $\delta_{\text{C}} 18$ ppm to $\delta_{\text{C}} 28$ ppm)




Figure S6: HSQC spectrum of **1** (carneusin A) (From δ_{C} 100 ppm to δ_{C} 116 ppm)




Figure S7: HMBC spectrum of **1** (carnesuin A)




Figure S8: HMBC spectrum of **1** (carnesuin A) (From δ_C 15 ppm to δ_C 70 ppm)




Figure S9: HMBC spectrum of **1** (carnesuin A) (From δ_C 100 ppm to δ_C 120 ppm)




Figure S10: HMBC spectrum of **1** (carnesuin A) (From δ_C 125 ppm to δ_C 195 ppm)




Figure S11: ^1H - ^1H COSY spectrum of **1** (carnesuin A)




Figure S12: NOESY spectrum of **1** (carneusin A)




Figure S13: HR-ESI-MS spectrum of **2** (carneusin B)




Figure S14: $^1\text{H-NMR}$ (600 MHz, $\text{DMSO-}d_6$) spectrum of **2** (carnesin B)




Figure S15: ^{13}C -NMR (150 MHz, CDCl_3) spectrum of **2** (carneusin B)




Figure S16: HSQC spectrum of **2** (carnesin B)




Figure S17: HMBC spectrum of **2** (carneusin B)




Figure S18: HMBC spectrum of **2** (carnesuin B) (From δ_C 5 ppm to 45 ppm)




Figure S19: HMBC spectrum of **2** (carnesin B) (From δ_C 120 ppm to 190 ppm)

Table S1: Relative free energies^a and equilibrium populations^b of conformers for (1'S, 4'S, 5'S)-1^c

conformer	ΔG	P (%)
1a	0.00	100.00

^a B3LYP/6-31G(d), in kcal/mol. ^b From ΔG values at 298.15 K. ^c in MeOH, no imaginary frequency






Figure S20: . The optimized structures (left) and the calculated CD spectra of conformers (1'S 4'S, 5'S)-1 in MeOH at M06-2X/def2TZVP level (right). $\sigma = 0.22$ eV



Table S2: Relative free energies^a and equilibrium populations^b of conformers for (6*S*)-**2**^c

conformer	ΔG	P (%)
2a	0.00	69.13
2b	1.03	11.98
2c	1.15	9.86
2d	1.20	9.03



^a B3LYP/6-31G(d), in kcal/mol. ^b From ΔG values at 298.15 K. ^c in MeOH, no imaginary frequency





(6S)-2a



(6S)-2b



(6S)-2c



(6S)-2d

Figure S21: The optimized structures (left) and the calculated CD spectra of conformers (6S)-2 in MeOH at M06-2X/def2TZVP level (right). $\sigma = 0.3$ eV






		DP4+	 89.54%	 10.46%
Nuclei	sp2?	experimental	Isomer 1	Isomer 2
C	x	174.92	173.925	173.405
C		36.16	40.254	39.546
C		34.62	36.852	35.959
C	x	179	182.141	181.509
C		15.92	15.798	15.255
C	x	129.45	134.359	134.334
C	x	162.3	168.816	168.733
C	x	128.81	106.88	106.844
C		52.72	53.321	53.359

Figure S22: DP4+ probabilities (%) for conformers (6*S*)-**2** (isomer 1) and (6*R*)-**2** (isomer 2)

<p>Score: 95</p> <p>1.</p> <p>28458-23-3</p>  <p>Absolute stereochemistry.</p> <p>C₂₀ H₁₆ O₈ 2,6-Epoxy-2H-anthra[2,3-b]oxocin-8,13-dione, 3,4,5,6-tetrahydro-5,7,9,11-tetrahydroxy-2-methyl-, (2S,5S,6R)-</p> <p>Key Physical Properties:</p> <p>Molecular Weight 384.34</p> <p>Melting Point (Experimental) Value: 203-206 °C</p> <p>Boiling Point (Predicted) Value: 659.2±55.0 °C Condition: Press: 760 Torr</p> <p>Density (Predicted) Value: 1.676±0.06 g/cm³ Condition: Temp: 20 °C Press: 760 Torr</p> <p>pKa (Predicted) Value: 5.70±0.60 Condition: Most Acidic Temp: 25 °C</p> <p>Related Info: ~ 55 References Reactions ~ 2 Commercial Sources Spectra Experimental Properties</p>	<p>Score: 95</p> <p>2.</p> <p>93922-50-0</p>  <p>Absolute stereochemistry.</p> <p>C₂₀ H₁₆ O₈ 2,6-Epoxy-2H-anthra[2,3-b]oxocin-8,13-dione, 3,4,5,6-tetrahydro-5,7,9,11-tetrahydroxy-2-methyl-, [2S-(2α,5β,6α)]- (9CI)</p> <p>Key Physical Properties:</p> <p>Molecular Weight 384.34</p> <p>Boiling Point (Predicted) Value: 659.2±55.0 °C Condition: Press: 760 Torr</p> <p>Density (Predicted) Value: 1.676±0.06 g/cm³ Condition: Temp: 20 °C Press: 760 Torr</p> <p>pKa (Predicted) Value: 5.70±0.60 Condition: Most Acidic Temp: 25 °C</p> <p>Related Info: ~ 1 References</p>	<p>Score: 95</p> <p>3.</p> <p>99528-66-2</p>  <p>Relative stereochemistry.</p> <p>C₂₀ H₁₆ O₈ 2,6-Epoxy-2H-anthra[2,3-b]oxocin-8,13-dione, 3,4,5,6-tetrahydro-5,7,9,11-tetrahydroxy-2-methyl-, (2R,5R,6S)-rel-</p> <p>Key Physical Properties:</p> <p>Molecular Weight 384.34</p> <p>Boiling Point (Predicted) Value: 659.2±55.0 °C Condition: Press: 760 Torr</p> <p>Density (Predicted) Value: 1.676±0.06 g/cm³ Condition: Temp: 20 °C Press: 760 Torr</p> <p>pKa (Predicted) Value: 5.70±0.60 Condition: Most Acidic Temp: 25 °C</p> <p>Related Info: ~ 2 References Reactions</p>
--	--	--












<p>Score: 95</p> <p>4.</p> <p>99528-67-3</p>  <p>Relative stereochemistry.</p> <p>C₂₀ H₁₆ O₈ 2,6-Epoxy-2H-anthra[2,3-b]oxocin-8,13-dione, 3,4,5,6-tetrahydro-5,7,9,11-tetrahydroxy-2-methyl-, (2R,5S,6S)-rel-</p> <p>Key Physical Properties:</p> <p>Molecular Weight 384.34</p> <p>Boiling Point (Predicted) Value: 659.2±55.0 °C Condition: Press: 760 Torr</p> <p>Density (Predicted) Value: 1.676±0.06 g/cm³ Condition: Temp: 20 °C Press: 760 Torr</p> <p>pKa (Predicted) Value: 5.70±0.60 Condition: Most Acidic Temp: 25 °C</p> <p>Related Info: ~ 1 References Reactions Spectra</p>	<p>Score: 93</p> <p>5.</p> <p>14016-29-6</p>  <p>Absolute stereochemistry.</p> <p>C₂₀ H₁₆ O₇ 2,6-Epoxy-2H-anthra[2,3-b]oxocin-8,13-dione, 3,4,5,6-tetrahydro-7,9,11-trihydroxy-2-methyl-, (2S,6S)-</p> <p>Key Physical Properties:</p> <p>Molecular Weight 368.34</p> <p>Melting Point (Experimental) Value: 288-293 °C</p> <p>Boiling Point (Predicted) Value: 602.9±55.0 °C Condition: Press: 760 Torr</p> <p>Density (Predicted) Value: 1.585±0.06 g/cm³ Condition: Temp: 20 °C Press: 760 Torr</p> <p>pKa (Predicted) Value: 5.83±0.40 Condition: Most Acidic Temp: 25 °C</p> <p>Related Info: ~ 201 References Reactions ~ 7 Commercial Sources Experimental Properties</p>	<p>Score: 93</p> <p>6.</p> <p>79896-28-9</p>  <p>C₂₀ H₁₆ O₇ 2,6-Epoxy-2H-anthra[2,3-b]oxocin-8,13-dione, 3,4,5,6-tetrahydro-7,9,11-trihydroxy-2-methyl-</p> <p>Key Physical Properties:</p> <p>Molecular Weight 368.34</p> <p>Melting Point (Experimental) Value: 282-288 °C (decomp) Condition: Solv: chloroform (67-66-3)</p> <p>Boiling Point (Predicted) Value: 602.9±55.0 °C Condition: Press: 760 Torr</p> <p>Density (Predicted) Value: 1.585±0.06 g/cm³ Condition: Temp: 20 °C Press: 760 Torr</p> <p>pKa (Predicted) Value: 5.83±0.40 Condition: Most Acidic Temp: 25 °C</p> <p>Related Info: ~ 16 References Reactions ~ 3 Commercial Sources Spectra Experimental Properties</p>
---	--	--

Figure S23 : The Scifinder searching results of compound 1

Table S3 : The structural comparison of similar compounds with **1**

No.	Similarity score	Similar chemical structure and CAS number	compound 1
1	95	 CAS NO. 28458-23-3 Absolute stereochemistry	
2	95	 CAS NO. 93922-50-0 Absolute stereochemistry	
3	95	 CAS NO. 99528-66-2 Relative stereochemistry	
4	95	 CAS NO. 99528-67-3 Relative stereochemistry	







<p>Score: 91 1. 824393-57-9</p>  <p>Double bond geometry as shown., Rotation (-), Absolute stereochemistry.</p> <p>C₁₀ H₁₅ N O₅ 2-Butenoic acid, 4-[[[(1S)-1-(methoxycarbonyl)-2-methylpropyl]amino]-4-oxo-, (2Z)-</p> <p>Key Physical Properties: Molecular Weight 229.23 Boiling Point (Predicted) Value: 432.1±45.0 °C Condition: Press: 760 Torr Density (Predicted) Value: 1.193±0.06 g/cm³ Condition: Temp: 20 °C Press: 760 Torr pKa (Predicted) Value: 2.78±0.25 Condition: Most Acidic Temp: 25 °C</p> <p>Related Info: ~ 3 References Reactions</p>	<p>Score: 91 2. 2165716-67-4</p>  <p>Absolute stereochemistry., Double bond geometry as shown.</p> <p>C₁₀ H₁₅ N O₅ 2-Butenoic acid, 4-[[[(1S)-1-(methoxycarbonyl)-2-methylpropyl]amino]-4-oxo-, (2E)-</p> <p>Key Physical Properties: Molecular Weight 229.23 Boiling Point (Predicted) Value: 432.1±45.0 °C Condition: Press: 760 Torr Density (Predicted) Value: 1.193±0.06 g/cm³ Condition: Temp: 20 °C Press: 760 Torr pKa (Predicted) Value: 3.47±0.10 Condition: Most Acidic Temp: 25 °C</p> <p>Related Info: ~ 0 References ~ 1 Commercial Sources</p>	<p>Score: 91 3. 2307909-20-0</p>  <p>Absolute stereochemistry., Double bond geometry unknown.</p> <p>C₁₀ H₁₅ N O₅ INDEX NAME NOT YET ASSIGNED</p> <p>Key Physical Properties: Molecular Weight 229.23 Boiling Point (Predicted) Value: 432.1±45.0 °C Condition: Press: 760 Torr Density (Predicted) Value: 1.193±0.06 g/cm³ Condition: Temp: 20 °C Press: 760 Torr pKa (Predicted) Value: 3.47±0.10 Condition: Most Acidic Temp: 25 °C</p> <p>Related Info: ~ 0 References ~ 1 Commercial Sources</p>
<p>Score: 90 4. 65306-00-5</p>  <p>C₉ H₁₅ N O₅ Butanoic acid, 4-[[[(1-carboxylethyl)amino]-3-methyl-4-oxo-, 1-methyl ester</p> <p>Key Physical Properties: Molecular Weight 217.22 Boiling Point (Predicted) Value: 438.1±30.0 °C Condition: Press: 760 Torr Density (Predicted) Value: 1.194±0.06 g/cm³ Condition: Temp: 20 °C Press: 760 Torr pKa (Predicted) Value: 3.52±0.10 Condition: Most Acidic Temp: 25 °C</p> <p>Related Info: ~ 2 References ~ 2 Commercial Sources</p>	<p>Score: 90 5. 124326-04-1</p>  <p>Double bond geometry as shown., Absolute stereochemistry.</p> <p>C₈ H₁₁ N O₅ 2-Butenoic acid, 4-[[[(1S)-2-methoxy-1-methyl-2-oxoethyl]amino]-4-oxo-, (2Z)-</p> <p>Key Physical Properties: Molecular Weight 201.18 Boiling Point (Predicted) Value: 429.0±45.0 °C Condition: Press: 760 Torr Density (Predicted) Value: 1.268±0.06 g/cm³ Condition: Temp: 20 °C Press: 760 Torr pKa (Predicted) Value: 2.79±0.25 Condition: Most Acidic Temp: 25 °C</p> <p>Related Info: ~ 5 References Reactions ~ 1 Commercial Sources</p>	<p>Score: 90 6. 124326-05-2</p>  <p>Double bond geometry as shown., Absolute stereochemistry.</p> <p>C₈ H₁₁ N O₅ 2-Butenoic acid, 4-[[[(2-methoxy-1-methyl-2-oxoethyl)amino]-4-oxo-, [S:(E)]-, (9CI)</p> <p>Key Physical Properties: Molecular Weight 201.18 Boiling Point (Predicted) Value: 429.0±45.0 °C Condition: Press: 760 Torr Density (Predicted) Value: 1.268±0.06 g/cm³ Condition: Temp: 20 °C Press: 760 Torr pKa (Predicted) Value: 3.47±0.10 Condition: Most Acidic Temp: 25 °C</p> <p>Related Info: ~ 1 References ~ 5 Commercial Sources</p>

Figure S24 : The Scifinder searching results of compound 2

Table S4 : The structural comparison of similar compounds with **2**







No.	Similarity score	Chemical structure and CAS number	Chemical structure of compound 2
1	91	 <p>CAS NO. 824393-57-9</p> <p>Absolute stereochemistry</p>	
2	91	 <p>CAS NO. 2165716-67-4</p> <p>Absolute stereochemistry</p>	
3	91	 <p>CAS NO. 2307909-20-0</p> <p>Absolute stereochemistry</p>	

Table S5: The NMR data comparison of similar compounds with **1**

Position	Compound 1 ^a		nidurufin ^{a, b} (CAS NO. 28458-23-3)		2'-epinidurufin ^c (CAS NO. 93922-50-0)	
	δ_{H}	δ_{C} (mult.)	δ_{H}	δ_{C} (mult.)	δ_{H}	δ_{C} (mult.)
1	-	158.1 (C)	-	159.8 (C)	NMR data is not available	
2	-	116.1 (C)	-	115.0 (C)		
3	-	158.7 (C)	-	158.4 (C)		
4	6.85, <i>s</i>	107.2 (CH)	6.85, <i>s</i>	107.4 (CH)		
4a	-	133.0 (C)	-	134.6 (C)		
5	6.94, <i>d</i> , <i>J</i> = 2.4	109.4 (CH)	6.97, <i>s</i>	108.9 (CH)		
6	-	166.2 (C)	-	164.2 (C)		
7	6.43, <i>d</i> , <i>J</i> = 2.4	107.9 (CH)	6.49, <i>s</i>	108.0 (CH)		
8	-	164.4 (C)	-	165.2 (C)		
8a	-	108.0 (C)	-	108.5 (C)		
9	-	188.3 (C)	-	188.6 (C)		
9a	-	108.4 (C)	-	108.2 (C)		
10	-	180.6 (C)	-	180.5 (C)		
10a	-	134.6 (C)	-	133.1 (C)		
1'	5.09, <i>d</i> , <i>J</i> = 3.0	65.7 (CH)	5.36, overlapped	70.7 (CH)		
2'	2.32, <i>ddt</i> , <i>J</i> = 17.3, 8.3, 3.8 1.45, <i>d</i> , <i>J</i> = 13.6	21.9 (CH ₂)	3.75, <i>m</i>	63.5 (CH)		
3'	1.64, <i>d</i> , <i>J</i> = 13.0 1.53, <i>td</i> , <i>J</i> = 13.0, 3.2	23.5 (CH ₂)	2.16, <i>m</i> 1.82, <i>m</i>	30.2 (CH ₂)		
4'	3.55, <i>t</i> , <i>J</i> = 2.8	66.7 (CH)	1.57, <i>m</i>	22.7 (CH)		
5'	-	102.6 (C)	-	101.5 (C)		
6'	1.49, <i>s</i>	24.0 (CH ₃)	1.54, <i>s</i>	27.2 (CH ₃)		

^a in DMSO-*d*₆.^b reference: X. W. Luo, H. M. Lu, X. Q. Chen, X. F. Zhou, C. H. Gao and Y. H. Liu (2020). Secondary metabolites and their biological activities from the sponge derived fungus *Aspergillus versicolor*, *Chem. Nat. Comp.* **56**, 716-719.^c reference: R. A. Murphy Jr and M. P. Cava (1984). Stereochemistry of nidurufin: synthesis of 6,8-dideoxynidurufin and 6,8-dideoxyepinidurufin, *J. Am. Chem. Soc.* **106**, 7630-7632.