Supporting Information

Org. Commun. 16:1 (2023) 1-10

Molecular docking and antioxidant activity studies of imidodithiocarbonate derivatives containing pyrimidine

Elif Korkusuz, Yusuf Sert², Emine Kılıçkaya Selvi¹,

Hava Aydın³, İrfan Koca⁴ and İsmail Yıldırım³

 ¹ Mustafa Cikrikcioglu Vocational College, Kayseri University, 38039, Kayseri, Türkiye
² Department of Physics, Faculty of Arts and Sciences, Yozgat Bozok University, 66900, Yozgat, Türkiye

³ Department of Chemistry, Faculty of Sciences, Erciyes University, Kayseri 38039, Türkiye ⁴Department of Chemistry, Faculty of Arts and Sciences, Yozgat Bozok University, 66900, Yozgat, Türkiye

Table of Contents	Page
Figure S1: FT-IR Spectrum of 3a	2
Figure S2: ¹ H-NMR (400 MHz, CDCl ₃) Spectrum of 3a	3
Figure S3: ¹³ C-NMR (100 MHz, CDCl3) Spectrum of 3a	4
Figure S4: FT-IR Spectrum of 3b	5
Figure S5: ¹ H-NMR (400 MHz, CDCl ₃) Spectrum of 3b	6
Figure S6: ¹³ C-NMR (100 MHz, CDCl3) Spectrum of 3b	7
Figure S7: FT-IR Spectrum of 3c	8
Figure S8: ¹ H-NMR (400 MHz, CDCl ₃) Spectrum of 3c	9
Figure S9: ¹³ C-NMR (100 MHz, CDCl3) Spectrum of 3c	10




Figure S1: FT-IR Spectrum of 3a




Figure S2: ¹H-NMR (400 MHz, CDCl₃) Spectrum of 3a




Figure S3: ¹³C-NMR (100 MHz, CDCl₃) Spectrum of 3a




Figure S4: FT-IR Spectrum of 3b




Figure S5: ¹H-NMR (400 MHz, CDCl₃) Spectrum of 3b




Figure S6: ¹³C-NMR (100 MHz, CDCl3) Spectrum of 3b




Figure S7: FT-IR Spectrum of 3c




Figure S8: ¹H-NMR (400 MHz, CDCl₃) Spectrum of 3c




Figure S9: ¹³C-NMR (100 MHz, CDCl3) Spectrum of 3c