

Supporting Information

J. Chem. Metrol. 18:1 (2024) 1-9

Development and application of RP–HPLC method for estimation and evaluation of stability of brinzolamide in ophthalmic formulation

Mehul Patel¹, Ritu Dwivedi¹, Umang Shah¹
and Heta Kachhiya²

¹*Department of Pharmaceutical Chemistry and Analysis, Ramanbhai Patel College of Pharmacy, Charotar University of Science and Technology, CHARUSAT Campus, Changa, Anand, Gujarat-388421, India*

²*Department of Pharmaceutical Chemistry and Analysis, Indukaka Ipcowala College of Pharmacy, CVM University, Beyond GIDC, P.B. No. 53, Vitthal Udyognagar- 388 121, Gujarat, India*

Table of Contents	Page
Figure S1: Structure of brinzolamide (BZA).	2
Figure S2: Overlay chromatogram of brinzolamide (5 to 30µg/mL)	3
Figure S3: Peak purity analysis of brinzolamide	4
Figure S4: Chromatogram of brinzolamide degraded with acid degradation	5
Figure S5: Chromatogram of brinzolamide degraded with base degradation	6
Figure S6: Chromatogram of brinzolamide degraded with oxidation degradation	7
Figure S7: Chromatogram of brinzolamide degraded with photolytic degradation	8
Figure S8: Chromatogram of brinzolamide degraded with thermal degradation	9




Figure S1 : Structure of brinzolamide (BZA).





Figure S2 : Overlay chromatogram of brinzolamide (5 to 30µg/mL)



Impurity :Not Detected
Peak purity index : 0.999737
Single point threshold : 0.998355
Minimum peak purity index : 1382

Figure S3 : Peak purity analysis of brinzolamide




Figure S4 : Chromatogram of brinzolamide degraded with acid degradation




Figure S5 : Chromatogram of brinzolamide degraded with base degradation




Figure S6 : Chromatogram of brinzolamide degraded with oxidation degradation




Figure S7 : Chromatogram of brinzolamide degraded with photolytic degradation




Figure S8 : Chromatogram of brinzolamide degraded with thermal degradation