

Supporting Information

Org. Commun. 17:4 (2024) 265-271

Heat gun-assisted rapid and efficient synthesis of tetra-substituted Zinc(II) phthalocyanines functionalized with alkylthio or alkoxy chains

Hasan Yavuz Gören^{1,2}, Gizem Gümüştöz Çelik^{1*} and Ayşe Gül Gürek¹

¹ Department of Chemistry, Faculty of Natural Sciences, Gebze Technical University, 41400, Gebze, Kocaeli, Türkiye

² Techno Fuel and Chemical Technologies, Technopark R&D Park No:20 Gebze, Kocaeli, Türkiye

Table of Contents	Page
Figure S1: IR ATR spectra for compound 1a.	2
Figure S2: ¹ H NMR spectra for compound 1a.	2
Figure S3: ¹³ C NMR spectra for compound 1a.	3
Figure S4: Ms spectra for compound 1a.	3
Figure S5: IR ATR spectra for compound 2a.	4
Figure S6: Ms spectra for compound 2a.	4
Figure S7: IR ATR spectra for compound 3a.	5
Figure S8: Ms spectra for compound 3a.	5
Figure S9: IR ATR spectra for compound 4a.	6
Figure S10: Ms spectra for compound 4a.	6
Figure S11: IR ATR spectra for compound 1b.	7
Figure S12: MALDI-TOF spectra for compound 1b.	7
Figure S13: IR ATR spectra for compound 2b.	8
Figure S14: MALDI-TOF spectra for compound 2b (Matrix DIT).	8
Figure S15: IR ATR spectra for compound 3b.	9
Figure S16: MALDI-TOF spectra for compound 3b (Matrix DIT).	9
Figure S17: IR ATR spectra for compound 4b.	10
Figure S18: MALDI-TOF spectra for compound 4b (Matrix DIT).	10
Figure S19: a) UV-Vis graph in the range of 500-800 nm obtained at different concentrations of compound 2b, b) A graph plotting concentration versus area 2b.	11
Figure S20: a) UV-Vis graph in the range of 500-800 nm obtained at different concentrations of compound 3b, b) A graph plotting concentration versus area 3b.	12
Figure S21: a) UV-Vis graph in the range of 500-800 nm obtained at different concentrations of compound 4b, b) A graph plotting concentration versus area 4b.	13

* Corresponding authors: E-Mail: gizemgumusgozc@gtu.edu.tr




Figure S1: IR ATR spectra for compound 1a.




Figure S2: ¹H NMR spectra for compound 1a.




Figure S3: ^{13}C NMR spectra for compound **1a**.




Figure S4: Ms spectra for compound **1a**.




Figure S5: IR ATR spectra for compound 2a.




Figure S6: Ms spectra for compound 2a.




Figure S7: IR ATR spectra for compound 3a.




Figure S8: Ms spectra for compound 3a.




Figure S9: IR ATR spectra for compound **4a**.




Figure S10: Ms spectra for compound **4a**.




Figure S11: IR ATR spectra for compound **1b**.




Figure S12: MALDI-TOF spectra for compound **1b**.




Figure S13: IR ATR spectra for compound **2b**.




Figure S14: MALDI-TOF spectra for compound **2b** (Matrix DIT).




Figure S15: IR ATR spectra for compound **3b**.




Figure S16: MALDI-TOF spectra for compound **3b** (Matrix DIT).




Figure S17: IR ATR spectra for compound **4b**.




Figure S18: MALDI-TOF spectra for compound **4b** (Matrix DIT).




Figure S19: a) UV-Vis graph in the range of 500-800 nm obtained at different concentrations of compound **2b**, b) A graph plotting concentration versus area **2b**.




Figure S20: a) UV-Vis graph in the range of 500-800 nm obtained at different concentrations of compound 3b, b) A graph plotting concentration versus area 3b.




Figure S21: a) UV-Vis graph in the range of 500-800 nm obtained at different concentrations of compound **4b**, b) A graph plotting concentration versus area **4b**.